

DFG-Schwerpunktprogramm 1324

„Extraktion quantifizierbarer Information aus komplexen Systemen“

On the Power of Function Values for the Approximation Problem in Various Settings

E. Novak, H. Woźniakowski

Preprint 67



Edited by

AG Numerik/Optimierung
Fachbereich 12 - Mathematik und Informatik
Philipps-Universität Marburg
Hans-Meerwein-Str.
35032 Marburg


DFG-Schwerpunktprogramm 1324

„Extraktion quantifizierbarer Information aus komplexen Systemen“

On the Power of Function Values for the Approximation Problem in Various Settings

E. Novak, H. Woźniakowski

Preprint 67



The consecutive numbering of the publications is determined by their chronological order.

The aim of this preprint series is to make new research rapidly available for scientific discussion. Therefore, the responsibility for the contents is solely due to the authors. The publications will be distributed by the authors.

On the power of function values for the approximation problem in various settings

Erich Novak*

Mathematisches Institut, Universität Jena
Ernst-Abbe-Platz 2, 07740 Jena, Germany
email: novak@mathematik.uni-jena.de

Henryk Woźniakowski†

Department of Computer Science, Columbia University,
New York, NY 10027, USA, and
Institute of Applied Mathematics, University of Warsaw
ul. Banacha 2, 02-097 Warszawa, Poland
email: henryk@cs.columbia.edu

September 2, 2010

Abstract

This¹ is an expository paper on approximating functions from general Hilbert or Banach spaces in the worst case, average case and randomized settings with error measured in the L_p sense. We define the *power function* as the ratio between the best rate of convergence of algorithms that use function values over the best rate of convergence of algorithms that use arbitrary linear functionals for a worst possible Hilbert or Banach space for which the problem of approximating functions is well defined. Obviously, the power function takes values at most one. If these values are

*This author was partially supported by the DFG-Priority Program 1324

†This author was partially supported by the National Science Foundation.

¹Mathematics Subject Classification. Primary 41A25; Secondary 41A46, 65Y20; keywords: degree of approximation, widths and sampling numbers, complexity of numerical algorithms.

one or close to one than the power of function values is the same or almost the same as the power of arbitrary linear functionals. We summarize and supply a few new estimates on the power function. We also indicate eight open problems related to the power function since this function has not yet been studied for many cases. We believe that the open problems will be of interest to a general audience of mathematicians.

1 Introduction

This is an expository paper on the problem of approximating functions from general Hilbert or Banach spaces, which has been thoroughly studied in many books and papers. This problem has many variants depending on how we measure the error of such approximations (algorithms). A popular choice is to take the norm of an L_p space and all values of $p \in [1, \infty]$ have been considered. Furthermore, the error of algorithms can be defined in the worst case, average case or randomized setting. For the worst and average case settings, we consider deterministic algorithms. The worst case error is defined as the maximal error over the unit ball of a given space whereas the average case error is defined as the average error over the whole space with respect to a given measure. The usual choice is a zero mean Gaussian measure. For the randomized setting we consider randomized algorithms and the error is defined as the maximal expected error over the unit ball of a given space. Here, the expected error is given with respect to a probability distribution of randomized elements.

We approximate functions f by algorithms that use information about f given by finitely many linear functionals of f . Information is called *linear* if we can choose arbitrary linear functionals, and it is called *standard* if only function values may be used. Clearly, linear information is at least as powerful as standard information. For many applications, only standard information is available. But even in this case, it is a good idea to study linear information and learn how difficult is the function approximation problem. For example, if we can prove that even for linear information the problem is too difficult then, obviously, the same also holds for standard information. On the other hand, all positive results for linear information do not have to hold for standard information.

The main question addressed in this expository paper is to study the power of standard information or equivalently the power of function values. We want to know how much we lose if function values are used instead of linear information. Or more optimistically, we ask when the power of standard information is the same or nearly the same as the power of linear information. Such questions have been addressed in a number of papers and we will refer to them in the course of this paper. It has been usually done for specific spaces and only a few papers addressed these questions for some classes of spaces.

Our approach is a little more general and we want to verify the power of function val-

ues/standard information for all Hilbert or Banach spaces for which the problem of function approximation is well defined. More precisely, we define the *power function*²

$$\ell^{\text{sett-x}} : (0, \infty) \times [1, \infty] \rightarrow [0, 1].$$

Here $\text{sett} \in \{\text{wor}, \text{ran}, \text{avg}\}$ denotes the setting we use for the error definition. Hence, *wor* stands for the worst case setting, *ran* for the randomized setting, and *avg* for the average case setting. The second superscript $x \in \{H, B\}$ tells us if we consider only Hilbert spaces ($x = H$) or if we allow all Banach spaces ($x = B$).

We now explain the meaning of the value

$$\ell^{\text{sett-x}}(r, p).$$

The first argument r means that the n th minimal error behaves like n^{-r} if we use linear information. Since $r > 0$, we consider Hilbert or Banach spaces which admit convergence, and furthermore they admit a polynomial rate of convergence of the minimal errors. The second argument p denotes the use of the norm of L_p . The value $\ell^{\text{sett-x}}(r, p)$ is defined as r^{-1} times the best rate of convergence we obtain using function values for a worst possible choice of a Hilbert or Banach space. That is why $\ell^{\text{sett-x}}(r, p) \leq 1$, and the larger $\ell^{\text{sett-x}}(r, p)$ the better. Hence, if we have

$$\ell^{\text{sett-x}}(r, p) = 1$$

then the power of standard information is the same as the power of linear information. On the other hand, if we have

$$\ell^{\text{sett-x}}(r, p) = 0$$

then the power of standard information is zero as compared to the power of linear information. Finally, if we have

$$\ell^{\text{sett-x}}(r, p) \in (0, 1)$$

then we know qualitatively how much we may lose by using function values.

The concept of the power function seems to be new. For many values of p , especially when $p \neq 2$, this function has not yet been studied. This is especially the case for the randomized and average case settings. That is why we indicate eight open problems related to the power function with the hope that many mathematicians will be interested in solving them and advancing our knowledge about the power of function values.

In this paper we tried to summarize and supply a few new estimates on the power function. We now briefly indicate a few results presented in the paper.

²We needed to find a good one-letter name for the power function. Since in English and in Polish this would indicate the letter “ p ” which is already used as the parameter of the L_p space, we turn to German and use the word “Leistung”. That is why the letter ℓ denotes the power function.

In the worst case setting for the Hilbert case and $p = 2$, we conclude from [6, 8] that

$$\begin{aligned}\ell^{\text{wor-H}}(r, 2) &= 0 && \text{for all } r \in (0, \tfrac{1}{2}], \\ \ell^{\text{wor-H}}(r, 2) &\in \left[\frac{2r}{2r+1}, 1 \right] && \text{for all } r \in (\tfrac{1}{2}, \infty).\end{aligned}$$

Hence, the power of function values is zero for $r \leq 1/2$, and almost the same as the power of linear information for large r . One of the main open problem is to verify whether $\ell^{\text{wor-H}}(r, 2) = 1$ for all $r > 1/2$.

Staying with the worst case and Hilbert spaces but with $p \neq 2$, we conclude from [18] that

$$\ell^{\text{wor-H}}(r, p) = 0 \quad \text{for all } r \in \left(0, \min\left(\frac{1}{p}, \frac{1}{2}\right)\right].$$

For $r > \min(1/p, 1/2)$, we do not know anything about the values of $\ell^{\text{wor-H}}(r, p)$ except the case $p = \infty$ for which we know from [12] that

$$\ell^{\text{wor-H/B}}(r, \infty) \geq 1 - \frac{1}{r}.$$

Again for large r , the power of standard information is almost the same as the power of linear information.

For the worst case and the Banach case we have

$$\begin{aligned}\ell^{\text{wor-B}}(r, p) &= 0 && \text{for all } r \in (0, 1] \text{ and } p \in [1, 2], \\ \ell^{\text{wor-B}}(r, p) &= 0 && \text{for all } r \in (0, \tfrac{1}{2} + \tfrac{1}{p}] \text{ and } p \in (2, \infty), \\ \ell^{\text{wor-B}}(r, p) &\leq 1 - \frac{1}{r} \left(1 - \frac{1}{p}\right) && \text{for all } r > 1 \text{ and } p \in [1, 2], \\ \ell^{\text{wor-B}}(r, p) &\leq 1 - \frac{1}{2r} && \text{for all } r > 1 \text{ and } p \in [2, \infty), \\ 1 - \frac{1}{r} \leq \ell^{\text{wor-B}}(r, \infty) &\leq 1 - \frac{1}{2r} && \text{for all } r > 1.\end{aligned}$$

Even though we do not know much about the power function in this case, we can conclude that the Hilbert and Banach cases are different since

$$\ell^{\text{wor-B}}(r, 2) < \ell^{\text{wor-H}}(r, 2) \quad \text{for all } r \in (\tfrac{1}{2}, \infty).$$

Surprisingly enough, for the randomized setting with the Hilbert case and for the average case setting with the Hilbert or Banach case we have complete knowledge about the power function for $p = 2$ due to [28] and [5]. More precisely, we know that

$$\ell^{\text{ran-H}}(r, 2) = \ell^{\text{avg-H/B}}(r, 2) = 1 \quad \text{for all } r > 0.$$

More estimates of the power function can be found in the successive sections.

2 The worst case setting

Let F be a Hilbert or Banach space of functions, defined on a set Ω , such that the linear functionals $f \mapsto f(x)$ are continuous for all $x \in \Omega$. We assume that $F \subset L_p$ and that the embedding $I : F \rightarrow L_p$ is continuous³, where $I(f) = f$. We write H instead of F if F is a Hilbert space.

Let (c_n) be a sequence of nonnegative numbers. Assume first that (c_n) converges to zero. We define its (polynomial) rate of convergence $r(c_n)$ by

$$r(c_n) = \sup\{\beta \geq 0 \mid \lim_{n \rightarrow \infty} c_n n^\beta = 0\}.$$

If (c_n) is not convergent to zero we set $r(c_n) = 0$. Then $r(c_n)$ is well defined for all nonnegative sequences (c_n) . For example, the rate of convergence of $n^{-\alpha}$ is $\max(0, \alpha)$.

We approximate functions from F using finitely many arbitrary linear functionals $L \in F^*$ or function values $f(x)$ for some $x \in \Omega$. We define the error of such approximations by taking the worst case setting with respect to the L_p norm. The norm of L_p is denoted by $\|\cdot\|_p$.

We define two classes Λ^{all} and Λ^{std} of information evaluations. We have $\Lambda^{\text{std}} \subseteq \Lambda^{\text{all}} = F^*$ and Λ^{std} consists of linear functionals of the form $L_x(f) = f(x)$ for all $f \in F$, where $x \in \Omega$. We approximate functions from F by algorithms $A_n : F \rightarrow L_p$ given by

$$A_n(f) = \phi_n(L_1(f), L_2(f), \dots, L_n(f)),$$

where n is a nonnegative integer, $\phi_n : \mathbb{R}^n \rightarrow L_p$ is an arbitrary mapping, and $L_j \in \Lambda$, where $\Lambda \in \{\Lambda^{\text{all}}, \Lambda^{\text{std}}\}$. The choice of L_j can be adaptive, that is, it may depend on the already computed values $L_1(f), L_2(f), \dots, L_{j-1}(f)$. For $n = 0$, the mapping A_n is a constant element of the space L_p . More details can be found in e.g., [14, 21].

Hence, we consider algorithms that use n linear functionals either from the class Λ^{std} or from the class Λ^{all} . We define the minimal errors as follows.

Definition 1. For $n = 0$ and $n \in \mathbb{N} := \{1, 2, \dots\}$, let

$$e_n^{\text{all-wor}}(F, L_p) = \inf_{A_n \text{ with } L_j \in \Lambda^{\text{all}}} \sup_{\|f\|_F \leq 1} \|f - A_n(f)\|_p$$

and

$$e_n^{\text{std-wor}}(F, L_p) = \inf_{A_n \text{ with } L_j \in \Lambda^{\text{std}}} \sup_{\|f\|_F \leq 1} \|f - A_n(f)\|_p.$$

³We do not specify Ω or the underlying measure of L_p since they can be arbitrary.

For $n = 0$, it is easy to see that the best algorithm is $A_0 = 0$ and we obtain

$$e_0^{\text{all-wor}}(F, L_p) = e_0^{\text{std-wor}}(F, L_p) = \sup_{\|f\|_F \leq 1} \|f\|_p = \sup_{\|f\|_F \leq 1} \|I(f)\|_p = \|I\|.$$

This is the initial error that can be achieved without computing any linear functional on the functions f . Clearly,

$$e_n^{\text{all-wor}}(F, L_p) \leq e_n^{\text{std-wor}}(F, L_p) \quad \text{for all } n \in \mathbb{N}.$$

The sequences $(e_n^{\text{all-wor}}(F, L_p))$ and $(e_n^{\text{std-wor}}(F, L_p))$ are both non-increasing but not necessarily convergent to zero.

We want to compare the rates of convergence

$$r^{\text{all-wor}}(F, L_p) = r(e_n^{\text{all-wor}}(F, L_p)) \quad \text{and} \quad r^{\text{std-wor}}(F, L_p) = r(e_n^{\text{std-wor}}(F, L_p)).$$

In particular, we would like to know if it is possible that the sequence $(e_n^{\text{all-wor}}(F, L_p))$ converges much faster than the sequence $(e_n^{\text{std-wor}}(F, L_p))$. In many cases it is much easier to analyze the sequence $(e_n^{\text{all-wor}}(F, L_p))_{n \in \mathbb{N}}$. Then it is natural to ask what can be said about the sequence $(e_n^{\text{std-wor}}(F, L_p))_{n \in \mathbb{N}}$.

The main question addressed in this paper is to find or estimate the *power* function defined as $\ell^{\text{wor-x}} : (0, \infty) \times [1, \infty] \rightarrow [0, 1]$ by

$$\ell^{\text{wor-x}}(r, p) := \inf_{F: r^{\text{all-wor}}(F, L_p) = r} \frac{r^{\text{std-wor}}(F, L_p)}{r},$$

where $x \in \{H, B\}$ and indicates that the infimum is taken over all Hilbert spaces ($x = H$) or over all Banach spaces ($x = B$) continuously embedded in L_p for which function values are continuous linear functionals and the rate of convergence is r when we use arbitrary linear functionals.

It is easy to show, and it will be shown later, that the set of spaces F for which $r^{\text{all-wor}}(F, L_p) = r$ is not empty and therefore $\ell^{\text{wor-x}}$ is well defined. Obviously, $\ell^{\text{wor-x}}(r, p) \in [0, 1]$, as already claimed. The power function $\ell^{\text{wor-x}}$ measures the ratio between the best rates of convergence between approximations based on function values and on arbitrary linear functionals for a worst possible Hilbert or Banach space.

Suppose now that we take the minimal $n = n^{\text{wor-all/std}}(\varepsilon, F, L_p)$ for which the minimal worst case error is ε or $\varepsilon \|I\|$. Assume for simplicity that

$$e_n^{\text{all-wor}}(F, L_p) = n^{-r} \quad \text{and} \quad e_n^{\text{std-wor}}(F, L_p) = n^{-\alpha}$$

for some positive $\alpha = r^{\text{std-wor}}(F, L_p) \leq r$. Then

$$n^{\text{wor-all}}(\varepsilon, F, L_p) = \lceil \varepsilon^{-1/r} \rceil \quad \text{and} \quad n^{\text{wor-std}}(\varepsilon, F, L_p) = \lceil \varepsilon^{-1/\alpha} \rceil.$$

Clearly,

$$\lim_{\varepsilon \rightarrow 0} \frac{\ln n^{\text{wor-all}}(\varepsilon, F, L_p)}{\ln n^{\text{wor-std}}(\varepsilon, F, L_p)} = \frac{\alpha}{r} \geq \ell^{\text{wor-x}}(r, p).$$

Hence, if $\ell^{\text{wor-x}}(r, p) = 1$ then function values are as powerful as arbitrary linear functionals. On the other hand, the smaller $\ell^{\text{wor-x}}(r, p)$ the less powerful are function values as compared to arbitrary linear functionals. If $\ell^{\text{wor-x}}(r, p) = 0$ then the polynomial behavior of $n^{\text{all}}(\varepsilon, F, L_p)$ in ε^{-1} can be drastically changed for $n^{\text{std}}(\varepsilon, F, L_p)$.

Remark 1. It is well known that in some cases we can restrict ourselves only to linear algorithms. This holds when $p = \infty$ or when F is a Hilbert space. Then the corresponding infima for the minimal worst case errors are attained by

$$A_n(f) = \sum_{j=1}^n L_j(f) h_j$$

for some $L_j \in \Lambda \in \{\Lambda^{\text{std}}, \Lambda^{\text{all}}\}$ and $h_j \in L_p$. Much more about the existence of linear optimal error algorithms can be found in e.g., [14].

2.1 Double Hilbert Case

In this subsection we consider the approximation problem defined over a Hilbert space with the error measured also in the Hilbert space L_2 . That is why the name of this subsection is the double Hilbert case. Approximation in the L_2 norm for Hilbert spaces has been studied in many papers. For our problem the most relevant papers are [6], [8] and [27].

Assume that H is a Hilbert space of functions defined on a set Ω . Since we assume that function values are continuous this means that H is a reproducing kernel Hilbert space, $H = H(K)$, where K is defined on $\Omega \times \Omega$. Let $p = 2$ and $L_2 = L_2(\Omega, \mu)$ be the space of μ -square integrable functions with a measure μ on Ω . Since the embedding $I : H(K) \rightarrow L_2(\Omega, \mu)$ is continuous we have

$$\int_{\Omega} |f(t)|^2 d\mu(t) < \infty \quad \text{for all } f \in H(K).$$

In particular, we can take $f = K(\cdot, t)$ for arbitrary $t \in \Omega$, since such a function f belongs to $H(K)$. Therefore $W = I^*I : H(K) \rightarrow H(K)$, where I^* is defined by $\langle g, I(f) \rangle_{L_2(\Omega, \mu)} = \langle I^*(g), f \rangle_{H(K)}$ for all $f \in H(K)$ and $g \in L_2(\Omega, \mu)$, is given by

$$W(f)(x) = \int_{\Omega} K(x, t) f(t) d\mu(t) \quad \text{for all } f \in H(K).$$

Clearly, the operator W is self-adjoint and semi positive definite. It is well known that $\lim_n e_n^{\text{wor-all}}(H, L_2) = 0$ iff W is compact. Unfortunately, in general, W does not have to be compact and therefore $e_n^{\text{wor-all}}(H, L_2)$ does not have to go to zero. In fact, the sequence $e_n^{\text{wor-all}}(H, L_2)$ can be an arbitrary non-increasing sequence as the following example shows.

Example 1 (Arbitrary Sequence $e_n^{\text{wor-all}}(H, L_2)$).

Let $(\alpha_n)_{n \in \mathbb{N}}$ be an arbitrary non-increasing sequence of nonnegative numbers. Define k^* as the number of positive α_n . If all α_n are positive we formally set $k^* = \infty$. If k^* is finite let $\mathbb{N}_{k^*} = \{1, 2, \dots, k^*\}$, otherwise let $\mathbb{N}_{k^*} = \mathbb{N}$.

For $k \in \mathbb{N}_{k^*}$, take arbitrary disjoint nonempty intervals I_k such that $\bigcup_{k \in \mathbb{N}_{k^*}} I_k = [0, 1]$, and define the functions $e_k : [0, 1] \rightarrow \mathbb{R}$ by

$$e_k = \frac{\sqrt{\alpha_k}}{\sqrt{|I_k|}} 1_{I_k},$$

where $|I_k|$ denotes the Lebesgue measure of I_k . That is, $e_k(x) = \sqrt{\alpha_k/|I_k|}$ for $x \in I_k$ and $e_k(x) = 0$ for $x \notin I_k$.

Define the Hilbert space $H = \overline{\text{span}}\{e_k \mid k \in \mathbb{N}_{k^*}\}$ equipped with the inner product such that $\langle e_k, e_j \rangle_H = \delta_{k,j}$ for all $k, j \in \mathbb{N}_{k^*}$. This means that H is the space of piecewise constant functions $f : [0, 1] \rightarrow \mathbb{R}$ such that

$$f = \sum_{k=1}^{k^*} a_k e_k \quad \text{with} \quad a_k = \langle f, e_k \rangle_H \quad \text{and} \quad \|f\|_h = \left(\sum_{k=1}^{k^*} a_k^2 \right)^{1/2} < \infty.$$

The Hilbert space H has the reproducing kernel

$$K(x, y) = \sum_{k=1}^{k^*} e_k(x) e_k(y) \quad \text{for all} \quad x, y \in [0, 1].$$

Indeed, first of all note that K is well defined since for all x and y the last series has at most one nonzero term. Then $\langle K(\cdot, y_i), K(\cdot, y_j) \rangle_H = K(y_i, y_j)$, and

$$0 \leq \left\| \sum_{j=1}^m a_j K(\cdot, y_j) \right\|_H^2 = \sum_{i,j=1}^m a_i a_j K(y_i, y_j).$$

This shows that the matrix $(K(y_i, y_j))_{i,j=1,2,\dots,m}$ is symmetric and semi positive definite for all m and y_j . Clearly,

$$\langle f, K(\cdot, y) \rangle_H = \sum_{k=1}^{k^*} a_k e_k(y) = f(y),$$

and this completes the proof that K is the reproducing kernel of H .

Let $L_2 = L_2([0, 1])$ be the usual space of square Lebesgue integrable functions. Note that

$$\|e_k\|_2 = \frac{\alpha_k}{\sqrt{|I_k|}} \left(\int_{I_k} dt \right)^{1/2} = \alpha_k.$$

Therefore for any $f \in H$ we have

$$\|I(f)\|_2 = \|f\|_2 = \left(\sum_{k=1}^{k^*} \alpha_k^2 \alpha_k^2 \right)^{1/2} \leq \alpha_1 \|f\|_H.$$

The last bound is sharp, and therefore $\|I\| = \alpha_1$ showing that H is continuously embedded in L_2 . The operator W takes now the form

$$W(f) = \sum_{k=1}^{k^*} \langle f, e_k \rangle_2 e_k.$$

Note that $W(e_k) = \|e_k\|_2^2 e_k = \alpha_k^2 e_k$. This means that (α_k^2, e_k) are the eigenpairs of W and

$$W(f) = \sum_{k=1}^{k^*} \alpha_k^2 \langle f, e_k \rangle_H e_k.$$

It is well known that

$$e_n^{\text{wor-all}}(H, L_2) = \alpha_{n+1} \quad \text{for all } k = 0, 1, \dots$$

This proves that the behavior of $e_n^{\text{wor-all}}(H, L_2)$ can be arbitrary and, in general, we do not have convergence of $e_n^{\text{wor-all}}(H, L_2)$ to zero. Clearly, W is compact iff $\lim_n \alpha_n = 0$.

On the other hand, this example also shows that for a given $\beta \geq 0$ we can define a sequence α_k such that $r^{\text{all-wor}}(H, L_2) = \beta$. Indeed, it is enough to take $\alpha_k = k^{-\beta}$. \square

We discuss the power function $\ell^{\text{wor-H}}$. We now assume that $r^{\text{all-wor}}(H, L_2) = r > 0$, i.e., in particular we assume that the operator W is compact. Then W has eigenpairs (λ_j, η_j) ,

$$W(\eta_j) = \lambda_j \eta_j \quad \text{for all } j = 1, 2, \dots,$$

with $\langle \eta_j, \eta_k \rangle_H = \delta_{j,k}$. Without loss of generality we can order the eigenvalues λ_j such that $\lambda_1 \geq \lambda_2 \geq \dots$. For all $f \in H$ we have

$$\langle f, \eta_k \rangle_2 = \langle I(f), I(\eta_k) \rangle_2 = \langle f, W(\eta_k) \rangle_H = \lambda_k \langle f, \eta_k \rangle_H.$$

In particular, letting $f = \eta_j$, we conclude that the functions η_j are also orthogonal in the space L_2 .

It is well known that

$$e_n^{\text{wor-all}}(H, L_2) = \sqrt{\lambda_{n+1}} \quad \text{for all } n \in \mathbb{N}.$$

If $(e_n^{\text{wor-all}}(H, L_2))$ is convergent to zero then the same also holds for function values, i.e., $(e_n^{\text{wor-std}}(H, L_2))$ is also convergent. Indeed, we can reason as in Section 10.4 of [14] that all linear functionals can be approximated with an arbitrarily small error when we use function values, and then it is enough to remember that the error $\sqrt{\lambda_{n+1}}$ is achieved by a linear algorithm that uses n linear functionals $\langle f, \eta_j \rangle_{H(K)}$.

We have

$$\text{trace}(W) := \sum_{j=1}^{\infty} \lambda_j = \int_{\Omega} K(x, x) \, d\mu(x) = \sum_{n=0}^{\infty} [e_n^{\text{wor-all}}(H, L_2)]^2$$

and this is finite if $r^{\text{all-wor}}(H, L_2) > \frac{1}{2}$. If $r^{\text{all-wor}}(H, L_2) = \frac{1}{2}$ then $\sum_{n=0}^{\infty} [e_n^{\text{wor-all}}(H, L_2)]^2$ may be finite or infinite, and if $r^{\text{all-wor}}(H, L_2) < \frac{1}{2}$ then $\sum_{n=0}^{\infty} [e_n^{\text{wor-all}}(H, L_2)]^2$ is infinite.

We can now apply the result from [27] which states that $\int_{\Omega} K(x, x) \, d\mu(t) < \infty$ implies

$$r^{\text{std-wor}}(H, L_2) \geq r^{\text{all-wor}}(H, L_2) - \frac{1}{2}.$$

The result from [8] states that $r^{\text{all-wor}}(H, L_2) = r > \frac{1}{2}$ implies

$$r^{\text{std-wor}}(H, L_2) \geq r - \frac{r}{2r+1} = \frac{2r^2}{2r+1}.$$

The case $\sum_{n=0}^{\infty} [e_n^{\text{wor-all}}(H, L_2)]^2 = \infty$ was studied in [6]. It was shown that for any $r \in [0, \frac{1}{2}]$ there is a Hilbert space H such that

$$r^{\text{all-wor}}(H, L_2) = r \quad \text{and} \quad r^{\text{std-wor}}(H, L_2) = 0.$$

These results give us the following bounds on the power function $\ell^{\text{wor-H}}(\cdot, 2)$.

Theorem 1 ([6, 8]).

$$\begin{aligned} \ell^{\text{wor-H}}(r, 2) &= 0 && \text{for all } r \in (0, \frac{1}{2}], \\ \ell^{\text{wor-H}}(r, 2) &\in \left[\frac{2r}{2r+1}, 1 \right] && \text{for all } r \in (\frac{1}{2}, \infty). \end{aligned}$$

Although, we do not know the power function $\ell^{\text{wor-H}}(\cdot, 2)$ exactly, we know that there is a jump at $\frac{1}{2}$ since $\ell^{\text{wor-H}}(r, 2) \geq 1/2$ for all $r > 1/2$. Note also that for large r , the values of $\ell^{\text{wor-H}}(r, 2)$ are close to 1. This means that the power of function values for $r \in (0, \frac{1}{2})$ is zero, and is almost optimal for large r .

The problem of finding the exact values of $\ell^{\text{wor-H}}(r, 2)$ for $r > \frac{1}{2}$ is one of the main open problems in the worst case setting. We know that many people, including two of us, spent a lot of time trying to solve this problem but so far in vain. That is why we propose an open problem with the hope that it will be soon solved by the reader.

Open Problem 1. Suppose that $r > \frac{1}{2}$. Is it true that

$$\ell^{\text{wor-H}}(r, 2) = 1?$$

If not, what are the values of $\ell^{\text{wor-H}}(r, 2)$? □

The rate of convergence neglects to distinguish sequences that differ by a power of logarithms of n . Indeed, for $c_n = n^{-r}$ and $b_n = n^{-r}[\ln(n+1)]^\beta$ for a positive r and an arbitrary β we have $r(c_n) = r(b_n) = r$ independently of β . Obviously, for some standard spaces we would like to know not only the rate but also a power of logarithms. We discuss this point in the next example, where we use the notation $c_n \asymp b_n$ iff there exist positive numbers a_1 and a_2 such that $a_1 \leq c_n/b_n \leq a_2$ for sufficiently large n .

Example 2 (Sobolev spaces, $p = 2$).

a) For the standard Sobolev spaces $W_2^s([0, 1]^d)$ with an arbitrary $s > 0$, which measures the total smoothness of functions, it is well known that

$$e_n^{\text{all-wor}}(W_2^s([0, 1]^d), L_2) \asymp n^{-s/d}.$$

Of course, in general, function values are not well defined in $W_2^s([0, 1]^d)$. We must assume the embedding condition $2s > d$ and then function values are well defined and they are continuous linear functionals. Furthermore, it is known that

$$e_n^{\text{all-wor}}(W_2^s([0, 1]^d), L_2) \asymp e_n^{\text{std-wor}}(W_2^s([0, 1]^d), L_2) \asymp n^{-s/d},$$

see, e.g., [14] for a survey of such results.

b) For the Sobolev spaces $W_2^{r, \text{mix}}([0, 1]^d)$ with $r > 0$, which measures the smoothness of functions with respect to each variable, it is known that

$$e_n^{\text{all-wor}}(W_2^{r, \text{mix}}([0, 1]^d), L_2) \asymp n^{-r}(\log n)^{(d-1)r},$$

see, e.g., [1, 11, 17, 19, 21, 26], where this result can be found in various generality.

For function values, we must assume that $r > 1/2$, and then the best upper bound is

$$e_n^{\text{std-wor}}(W_2^{r,\text{mix}}([0, 1]^d), L_2) = \mathcal{O}(n^{-r}(\log n)^{(d-1)(r+1/2)}),$$

see [17, 19, 22].

It is *not* known whether this extra power $(d-1)/2$ for logarithms is needed. It would be very interesting to verify whether

$$e_n^{\text{all-wor}}(W_2^{r,\text{mix}}([0, 1]^d), L_2) \asymp e_n^{\text{std-wor}}(W_2^{r,\text{mix}}([0, 1]^d), L_2)$$

holds also for this example. □

The examples in [6] use very irregular sequences $(e_n^{\text{all-wor}}(H, L_2))$ and hence do not exclude a positive answer of the question in the next open problem.

Open Problem 2. Assume that $e_n^{\text{all-wor}}(H, L_2) \asymp n^{-r} [\ln(n+1)]^\beta$ with arbitrary $r > 0$ and $\beta \in \mathbb{R}$. Is it true that this implies

$$e_n^{\text{std-wor}}(H, L_2) \asymp e_n^{\text{all-wor}}(H, L_2)?$$

2.2 Single Hilbert Case

In this short subsection we mostly consider the approximation problem defined over a Hilbert space with the error measured in the non-Hilbert space L_p for $p \neq 2$. That is why the name of this subsection is the single Hilbert case.

We report a recent result of Tandetzky [18] who considers the approximation problem for arbitrary $p \in [1, \infty)$. He proved that for any $r \in (0, \min(\frac{1}{p}, \frac{1}{2})]$ there exists a Hilbert space H continuously embedded in $L_p = L_p([0, 1])$ such that

$$r^{\text{all-wor}}(H, L_p) = r \quad \text{and} \quad r^{\text{std-wor}}(H, L_p) = 0.$$

This result obviously implies that the power function is zero over $(0, \min(\frac{1}{p}, \frac{1}{2})]$. It seems to us that no example is known in the literature for a Hilbert space for which $e_n^{\text{all-wor}}(H, L_p)$ tends to zero faster than the sequence $e^{\text{std-wor}}(H, L_p)$ with the additional assumption that $r^{\text{all-wor}}(H, L_p) > \min(\frac{1}{p}, \frac{1}{2})$. This implies that we do not know the behavior of the power function over $(\min(\frac{1}{p}, \frac{1}{2}), \infty)$. We summarize our partial knowledge of the power function in the following theorem.

Theorem 2 ([18]). Let $p \neq 2$.

$$\begin{aligned} \ell^{\text{wor-H}}(r, p) &= 0 & \text{for all } r \in (0, \min(\frac{1}{p}, \frac{1}{2})], \\ \ell^{\text{wor-H}}(r, p) &\in [0, 1] & \text{for all } r \in (\min(\frac{1}{p}, \frac{1}{2}), \infty). \end{aligned}$$

Only for the case $p = \infty$ we know a little more about the behavior of the power function. In this case the rates are related as explained in the following theorem.

Theorem 3 ([12]). Let F be a Hilbert or a Banach space. Then

$$e_n^{\text{std-wor}}(F, L_\infty) \leq (1 + n) e_n^{\text{all-wor}}(F, L_\infty) \quad \text{for all } n \in \mathbb{N}. \quad (1)$$

This inequality follows from Proposition 1.2.5, page 16, in [12], where it is stated for the Kolmogorov widths and also applies to the linear or Gelfand widths.

The inequality (1) cannot be improved even if we assume that F is a Hilbert space. This follows from the following example.

Example 3. Take $F = H = \mathbb{R}^{n+1}$. That is $f \in H$ is now defined on $\{1, 2, \dots, n+1\}$ and can be identified with $f = [f_1, f_2, \dots, f_{n+1}]$ with $f_i = f(i)$. The space H is equipped with the inner product

$$\langle f, g \rangle_H = \left[\sum_{i=1}^{n+1} f_i \right] \left[\sum_{i=1}^{n+1} g_i \right] + \varepsilon \sum_{i=1}^{n+1} f_i g_i \quad \text{for all } f, g \in H.$$

The unit ball of H is now

$$B = \{f \in \mathbb{R}^{n+1} \mid [\sum_{i=1}^{n+1} f_i]^2 + \varepsilon \sum_{i=1}^{n+1} f_i^2 \leq 1\}.$$

Then for $\varepsilon \rightarrow 0$, we obtain

$$e_n^{\text{std-wor}}(F, L_\infty) \geq 1.$$

Indeed, knowing $f(x_i)$ for $i = 1, 2, \dots, n$, with $x_i \in \{1, 2, \dots, n+1\}$, we take f such that $f(x_i) = 0$. Then at least one component of f from the unit ball is free and can be taken as $\pm 1/\sqrt{1 + \varepsilon}$. This proves that the worst case error of any algorithm is at least $1/\sqrt{1 + \varepsilon}$ which in the limit as ε goes to zero is 1.

Consider the information

$$N(f) = [f_1 - f_2, f_2 - f_3, \dots, f_n - f_{n+1}] \quad \text{for all } f \in H.$$

It is known that the minimal error of all algorithms that use N is the supremum of $\|f\|_H$ for $f \in B$ and $N(f) = 0$. Observe that $N(f) = 0$ implies that $f = [c, c, \dots, c]$. Next, $f \in B$ implies that

$$c^2 \leq \frac{1 + \varepsilon/(n+1)}{(n+1)^2}.$$

Hence, again for $\varepsilon \rightarrow 0$, we obtain $e_n^{\text{all-wor}}(F, L_\infty) \leq 1/(n+1)$. \square

We now conclude a partial behavior of the power function $\ell^{\text{wor-H}}(\cdot, \infty)$ from Theorem 3. Let $r^{\text{all-wor}}(F, L_\infty) = r > 1$. Then the inequality (1) implies that

$$r^{\text{std-wor}}(F, L_\infty) \geq r - 1.$$

Thus, Theorem 3 implies the following behavior of the power function for $p = \infty$.

Theorem 4.

$$\begin{aligned} \ell^{\text{wor-H/B}}(r, \infty) &\in [0, 1] && \text{for all } r \in (0, 1], \\ \ell^{\text{wor-H/B}}(r, \infty) &\in \left[\frac{r-1}{r}, 1 \right] && \text{for all } r > 1. \end{aligned}$$

Hence, for both $p = 2$ and $p = \infty$, we see that for large r , the power function is almost one.

We want to guess the behavior of the power function for $r > \min(\frac{1}{p}, \frac{1}{2})$. It can be helpful to see the actual rates of convergence for some standard spaces. In particular, for $p = \infty$, the rates are known for Sobolev spaces.

Example 4 (Sobolev spaces, $p = \infty$).

a) For the Sobolev spaces $W_2^s([0, 1]^d)$ and an arbitrary s for which $2s > d$, it is well known that

$$e_n^{\text{all-wor}}(W_2^s([0, 1]^d), L_\infty) \asymp e_n^{\text{std-wor}}(W_2^s([0, 1]^d), L_\infty) \asymp n^{-s/d+1/2},$$

see, e.g., [14].

b) For the Sobolev spaces $W_2^{s,\text{mix}}([0, 1]^d)$ with $s > 1/2$, it is known that

$$e_n^{\text{all-wor}}(W_2^{s,\text{mix}}([0, 1]^d), L_\infty) \asymp e_n^{\text{std-wor}}(W_2^{s,\text{mix}}([0, 1]^d), L_\infty) \asymp n^{-s+1/2}(\log n)^{(d-1)s},$$

see [20]. □

Hence, at least for the standard Sobolev spaces the rates are the same even up to logarithmic factors. This again suggests that the power function can be just one over $(\min(\frac{1}{p}, \frac{1}{2}), \infty)$. This leads us to the next open problem.

Open Problem 3. Verify whether it is true that for all $p \in [1, \infty]$ we have

$$\ell^{\text{wor-H}}(r, p) = \begin{cases} 0 & \text{for all } r \in (0, \min(\frac{1}{p}, \frac{1}{2})], \\ 1 & \text{for all } r \in (\min(\frac{1}{p}, \frac{1}{2}), \infty). \end{cases}$$

□

We end this section with a remark on the rates of convergence for different p .

Remark 2. It is interesting to compare the sequences

$$e_n^{\text{all-wor}}(H, L_p) \quad \text{and/or} \quad e_n^{\text{std-wor}}(H, L_p)$$

for the same H but different p . The following example shows that, in general, there exists *no* relation between these sequences. Some relations do exist as shown in [7] but under some additional assumptions about H . The following example shows that some assumptions on H are indeed needed, otherwise the worst can happen.

Take $L_2 = L_2([0, 1])$, $L_\infty = L_\infty([0, 1])$ and assume that $[0, 1]$ is the disjoint union of intervals I_k with positive length λ_k , so that $\sum_{k=1}^{\infty} \lambda_k = 1$. Assume also that

$$\lambda_1 \geq \lambda_2 \geq \dots$$

and put $e_k = 1_{I_k}$. We define a Hilbert space H by its unit ball

$$B = \left\{ \sum_{k=1}^{\infty} \alpha_k e_k \mid \sum_{k=1}^{\infty} \frac{\alpha_k^2}{\gamma_k^2} \leq 1 \right\},$$

where

$$\gamma_1 \geq \gamma_2 \geq \dots > 0 \quad \text{with} \quad \lim_{k \rightarrow \infty} \gamma_k = 0.$$

Hence for $f = \sum_{k=1}^{\infty} \alpha_k e_k \in H$ we obtain

$$\|f\|_H^2 = \sum_{k=1}^{\infty} \frac{\alpha_k^2}{\gamma_k^2} \quad \text{and} \quad \|f\|_2^2 = \sum_{k=1}^{\infty} \alpha_k^2 \lambda_k, \quad \|f\|_\infty = \sup_k |\alpha_k|.$$

From this it is easy to conclude that the optimal approximation for L_2 as well as for L_∞ is given by

$$f = \sum_{k=1}^{\infty} \alpha_k e_k \mapsto \sum_{k=1}^n \alpha_k e_k.$$

Note that

$$\alpha_k = \langle f, e_k \rangle_H = f(x_k) \lambda_k,$$

where $x_k \in I_k$. This means that the optimal error algorithm for function values and linear functionals is the same, and therefore

$$e^{\text{all-wor}}(H, L_p) = e^{\text{std-wor}}(H, L_p) \quad \text{for} \quad p \in \{2, \infty\}.$$

However,

$$e_n^{\text{all-wor}}(H, L_\infty) = \gamma_{n+1} \quad \text{and} \quad e_n^{\text{all-wor}}(H, L_2) = \gamma_{n+1} \sqrt{\lambda_{n+1}}.$$

Since $\{\gamma_n\}$ and $\{\lambda_n\}$ are not related, it is easy to get an example with

$$r^{\text{all-wor}}(H, L_\infty) = 0 \quad \text{but} \quad r^{\text{all-wor}}(H, L_2) = \infty.$$

Hence, in general, the worst difference between the minimal rates for L_2 and L_∞ approximation can happen.

2.3 Banach Case

In this subsection we study the approximation problem defined over a Banach space that is continuously embedded in L_p . As always we assume that function evaluations are continuous functionals. We establish some bounds on the power functions by recalling known results for Sobolev spaces.

Example 5 (Sobolev spaces, $1 \leq p < \infty$).

For the Sobolev space $W_p^s([0, 1]^d)$ for an arbitrary $s > 0$, it is known that

$$e_n^{\text{all-wor}}(W_p^s([0, 1]^d), L_p) \asymp n^{-s/d}.$$

Function values are well defined in $W_p^s([0, 1]^d)$ only if the embedding condition $s/d > 1/p$ or $s = d$ and $p = 1$ holds. However, we may use the approach suggested in [2] that allows us to consider the case without this embedding condition. Namely, we limit ourselves only to continuous functions by taking

$$F = W_p^s([0, 1]^d) \cap C([0, 1]^d)$$

with norm

$$\|f\|_F = \|f\|_{W_p^s([0, 1]^d)} + \|f\|_{C([0, 1]^d)}.$$

Here, $C([0, 1]^d)$ is the space of continuous functions equipped with the max norm. Then F is a Banach space for which function values are well defined and correspond to linear continuous functionals. Then for $s/d \leq 1/p$ and $s/d < 1$ in the case $p = 1$, respectively, it was shown in [2] that

$$e_n^{\text{std-wor}}(F, L_p) \asymp 1.$$

□

The last example implies that

$$\begin{aligned}\ell^{\text{wor-B}}(r, p) &= 0 & \text{for all } r \in (0, 1/p] \text{ and } 1 < p < \infty, \\ \ell^{\text{wor-B}}(r, 1) &= 0 & \text{for all } r \in (0, 1).\end{aligned}$$

We now show that $\ell^{\text{wor-B}}(r, p) = 0$ over larger domains of r for a given p by recalling other results for Sobolev spaces.

Example 6 (Sobolev space $W_1^s([0, 1]^d)$, $1 \leq p < \infty$).

Consider the approximation problem for the Sobolev space $W_1^s([0, 1]^d)$ with error measured in $L_p = L_p([0, 1]^d)$. This problem is well defined and convergent for the class Λ^{all} if we assume that $s/d > 1 - 1/p$.

For $p \in [1, 2]$ we have

$$e_n^{\text{all-wor}}(W_1^s([0, 1]^d), L_p) \asymp n^{-s/d},$$

whereas for $p \in [2, \infty)$ we have

$$e_n^{\text{all-wor}}(W_1^s([0, 1]^d), L_p) \asymp n^{-s/d+1/2-1/p},$$

see e.g., [24]. The last relation also holds for $p = \infty$ as will be needed later.

The same results are also valid for the space $F = W_1^s([0, 1]^d) \cap C([0, 1]^d)$ with the norm

$$\|f\|_F = \|f\|_{W_1^s([0, 1]^d)} + \|f\|_{C([0, 1]^d)}.$$

For the space F we can consider function values for all $s/d > 1 - 1/p$. For $s/d \leq 1$ we have

$$e_n^{\text{std-wor}}(F, L_p) \asymp 1.$$

□

Let $p \in [1, 2]$. The last example implies that

$$\ell^{\text{wor-B}}(r, p) = 0 \quad \text{for all } r \in \left(1 - \frac{1}{p}, 1\right].$$

For $p = 1$, the last interval is $(0, 1]$. For $p \in (1, 2]$ we showed before that $\ell^{\text{wor-B}}(r, p) = 0$ for all $r \in (0, 1/p]$. Since $(0, 1/p] \cup (1 - 1/p, 1] = (0, 1]$ we obtain

$$\ell^{\text{wor-B}}(r, p) = 0 \quad \text{for all } r \in (0, 1] \text{ and } p \in [1, 2].$$

Let $p \in [2, \infty)$. The last example implies that

$$\ell^{\text{wor-B}}(r, p) = 0 \quad \text{for all } r \in \left(\frac{1}{2}, \frac{1}{2} + \frac{1}{p}\right].$$

To show that $\ell^{\text{wor-B}}(r, p) = 0$ also for $p \in [2, \infty)$ and $r \in (0, \frac{1}{2}]$ we increase the space $F = W_1^s([0, 1]) \cap C([0, 1])$ with the norm

$$\|f\|_F = \|f\|_{W_1^s([0,1])} + \|f\|_{C([0,1])}$$

from above (for $d = 1$) even more by adding functions from a Hölder class C^α , where $0 < \alpha \leq 1/2$. Hence we take the space

$$\tilde{F} = F + C^\alpha$$

with the norm

$$\|f\|_{\tilde{F}} := \inf\{\|g\|_F + \|h\|_{C^\alpha} \mid f = g + h, g \in F, h \in C^\alpha\}.$$

Since the unit ball of \tilde{F} is larger than that of F we still have $e_n^{\text{std-wor}}(\tilde{F}, L_p) \asymp 1$ for $s \leq 1$. It is well known that $e_n^{\text{all-wor}}(C^\alpha([0, 1]), L_p) \asymp n^{-\alpha}$ and the same holds for \tilde{F} if $\alpha \leq s - 1/2 + 1/p$. Hence for $p \in [2, \infty)$ we obtain

$$\ell^{\text{wor-B}}(r, p) = 0 \quad \text{for all} \quad r \in \left(0, \frac{1}{2} + \frac{1}{p}\right].$$

We learnt some properties of the power function by using known results for Sobolev spaces in the case $s/d \leq 1/p_1$ so that function values did not supply even convergence. Since we needed to assume that $s/d > 1/p_1 - 1/p$, the case $p = \infty$ could not be covered.

We now recall some results for Sobolev spaces when the embedding condition is satisfied and when there is a difference in the convergence rates between function values and arbitrary linear functionals.

Example 7 (Sobolev space $W_1^s([0, 1]^d)$, $1 \leq p \leq \infty$).

Consider the approximation problem for the Sobolev space $W_1^s([0, 1]^d)$ with error measured in L_p . We now assume that $s/d \geq 1$. Then function values are well defined and are continuous linear functionals. Furthermore,

$$e_n^{\text{std-wor}}(W_1^s([0, 1]^d), L_p) \asymp n^{-s/d+1-1/p},$$

see, e.g., the survey of such results in Section 4.2.4 of [14] or [23, 24]. □

The last two examples imply the following estimates of the power function. For all $r > 1$ and $p \in [1, 2]$ we have

$$\ell^{\text{wor-B}}(r, p) \leq 1 - \frac{1}{r} \left(1 - \frac{1}{p}\right),$$

and for all $r > 1$ and $p \in [2, \infty]$ we have

$$\ell^{\text{wor-B}}(r, p) \leq 1 - \frac{1}{2r}.$$

We summarize the properties of the power function established in this section in the following theorem. The only case where we have a positive lower bound is the case $p = \infty$, see Theorem 4.

Theorem 5.

$$\begin{aligned} \ell^{\text{wor-B}}(r, p) &= 0 && \text{for all } r \in (0, 1] \text{ and } p \in [1, 2], \\ \ell^{\text{wor-B}}(r, p) &= 0 && \text{for all } r \in (0, \frac{1}{2} + \frac{1}{p}] \text{ and } p \in (2, \infty), \\ \ell^{\text{wor-B}}(r, p) &\leq 1 - \frac{1}{r} \left(1 - \frac{1}{p}\right) && \text{for all } r > 1 \text{ and } p \in [1, 2], \\ \ell^{\text{wor-B}}(r, p) &\leq 1 - \frac{1}{2r} && \text{for all } r > 1 \text{ and } p \in [2, \infty), \\ 1 - \frac{1}{r} \leq \ell^{\text{wor-B}}(r, \infty) &\leq 1 - \frac{1}{2r} && \text{for all } r > 1. \end{aligned}$$

It is interesting to notice that although we do not know the exact values of the power functions in the Hilbert and Banach cases, we can check that they are different at least for $p = 2$. Indeed, from Theorems 1 and 5 we have

$$\begin{aligned} \ell^{\text{wor-B}}(r, 2) &= \ell^{\text{wor-H}}(r, 2) && \text{for all } r \in (0, \frac{1}{2}], \\ \ell^{\text{wor-B}}(r, 2) = 0 &< \frac{1}{2} \leq \ell^{\text{wor-H}}(r, 2) && \text{for all } r \in (\frac{1}{2}, 1], \\ \ell^{\text{wor-B}}(r, 2) \leq 1 - \frac{1}{2r} &< \frac{2r}{2r+1} \leq \ell^{\text{wor-H}}(r, 2) && \text{for all } r \in (1, \infty). \end{aligned}$$

This shows that at least for $p = 2$ the power of function values for the Hilbert case is larger than for the Banach case for all $r > \frac{1}{2}$.

Obviously, it would be desirable to find the exact values of the power function $\ell^{\text{wor-B}}(r, p)$ for all $r \in (0, \infty)$ and $p \in [1, \infty]$. However, it could be a very difficult problem. Hence, as maybe a less difficult problem, we would like to check the following property of the power function.

Open Problem 4. For $p \in [1, \infty]$, find the supremum $a^*(p)$ of a for which

$$\ell^{\text{wor-B}}(r, p) = 0 \quad \text{for all } r \in (0, a].$$

We only know that $a^*(p) \geq 1$ for all $p \in [1, \infty)$.

We already indicated that the power functions for the Hilbert and Banach cases are different for $p = 2$. It would be of interest to check if this holds for all p .

Open Problem 5. Find all $p \in [1, \infty]$ for which

$$\ell^{\text{wor-B}}(\cdot, p) \neq \ell^{\text{wor-H}}(\cdot, p).$$

Similar as in Example 3 we present an example of a Banach space F where the ratio

$$\frac{e_n^{\text{std-wor}}(F, L_p)}{e_n^{\text{all-wor}}(F, L_p)}$$

is large for $p > 1$ and a fixed n .

Example 8. Take $F = \ell_1^{n+1}$, i.e., $F = \mathbb{R}^{n+1}$ with the ℓ_1 norm. Then we obtain

$$e_n^{\text{std-wor}}(F, L_p) = (n+1)^{1-1/p} e_n^{\text{all-wor}}(F, L_p), \quad (2)$$

since $e_n^{\text{std-wor}}(F, L_p) = 1$ and $e_n^{\text{all-wor}}(F, L_p) = (n+1)^{1/p-1}$. The upper bound in the last statement follows again with the information $N(x) = (x_2 - x_1, x_3 - x_2, \dots, x_{n+1} - x_n)$ while the lower bound follows from the fact that the unit ball of ℓ_1^{n+1} contains a ℓ_p^{n+1} ball of radius $(n+1)^{1/p-1}$.

Again this ratio $(n+1)^{1-1/p}$ as in (2) can be obtained with a Hilbert space and actually we can take the same spaces as in Example 3, i.e., we define in $H = \mathbb{R}^{n+1}$ the scalar product

$$\langle f, g \rangle_H = \left[\sum_{i=1}^{n+1} f_i \right] \left[\sum_{i=1}^{n+1} g_i \right] + \varepsilon \sum_{i=1}^{n+1} f_i g_i \quad \text{for all } f, g \in H,$$

and consider the limit where $\varepsilon > 0$ tends to zero. □

We end this section with another open problem.

Open Problem 6. Find the supremum of $e_n^{\text{std-wor}}(F, L_p)/e_n^{\text{all-wor}}(F, L_p)$ over all Banach and/or Hilbert spaces. So far we know that

$$\sup_F \frac{e_n^{\text{std-wor}}(F, L_p)}{e_n^{\text{all-wor}}(F, L_p)} \geq (n+1)^{1-1/p}, \quad (3)$$

and equality holds if $p = \infty$.

3 The randomized setting

We approximate the embedding operator $I : F \rightarrow L_p$ in the randomized setting. We now briefly define this setting. The reader may find more on this subject, e.g., in [14, 15, 21].

We approximate I by algorithms A_n that use n values of linear functionals on the average and each linear functional is chosen randomly with respect to a probability distribution.

More precisely, the algorithm A_n is of the following form

$$A_n(f, \omega) = \phi_{n, \omega} \left(L_{1, \omega_1}(f), L_{2, \omega_2}(f), \dots, L_{n(\omega), \omega_{n(\omega)}}(f) \right). \quad (4)$$

Here $\omega = [\omega_1, \omega_2, \dots]$, and the linear functionals L_{j, ω_j} are random functionals distributed according to a probability distribution on elements ω_j which may depend on j as well as on the values already computed, i.e., on $L_{i, \omega_i}(f)$ for $i = 1, 2, \dots, j - 1$. The mapping $\phi_{n, \omega} : \mathbb{R}^{n(\omega)} \rightarrow L_p$ is a random mapping, and

$$\mathbb{E}_\omega n(\omega) \leq n.$$

We also allow adaptive choices of the functionals L_{j, ω_j} . That is, L_{j, ω_j} may depend on the already selected functionals and the values $L_{1, \omega_1}(f), L_{2, \omega_2}(f), \dots, L_{j-1, \omega_{j-1}}(f)$.

Without loss of generality, we assume that $A_n(f, \cdot)$ is measurable, and define the randomized error of A_n as

$$e^{\text{ran}}(A_n) = \sup_{\|f\|_F \leq 1} \left(\mathbb{E}_\omega \|I(f) - A_n(f, \omega)\|_p^2 \right)^{1/2}.$$

Again we compare such algorithms with algorithms that are based on function values, i.e., each L_{j, ω_j} is now of the form $L_{j, \omega_j}(f) = f(t_{j, \omega_j})$ and

$$A_n(f, \omega) = \phi_{n, \omega} \left(f(t_{1, \omega_1}), f(t_{2, \omega_2}), \dots, f(t_{n(\omega), \omega_{n(\omega)}}) \right). \quad (5)$$

Hence, we consider algorithms that use n linear functionals either from the class Λ^{std} or the class Λ^{all} . We define the minimal errors as follows.

Definition 2. For $n \in \mathbb{N}_0$, let

$$e_n^{\text{all-ran}}(F, L_p) = \inf \left\{ e^{\text{ran}}(A_n) \mid L_j \in \Lambda^{\text{all}} \text{ and } A_n \text{ as in (4)} \right\},$$

and

$$e_n^{\text{std-ran}}(F, L_p) = \inf \left\{ e^{\text{ran}}(A_n) \mid L_j \in \Lambda^{\text{std}} \text{ and } A_n \text{ as in (5)} \right\}.$$

For $n = 0$, it is easy to see that the best algorithm is $A_0 = 0$ and obtain

$$e_0^{\text{all-ran}}(F, L_p) = e_0^{\text{std-ran}}(F, L_p) = \sup_{\|f\|_F \leq 1} \|f\|_p = \sup_{\|f\|_F \leq 1} \|I(f)\|_p = \|I\|.$$

This is the initial error that can be achieved without computing any linear functional on the functions f . Clearly,

$$e_n^{\text{all-ran}}(F, L_p) \leq e_n^{\text{std-ran}}(F, L_p) \quad \text{for all } n \in \mathbb{N}.$$

The sequences $(e_n^{\text{all-ran}}(F, L_p))$ and $(e_n^{\text{std-ran}}(F, L_p))$ are both non-increasing but not necessarily convergent to zero.

As in the worst case setting, we want to compare the rates of convergence

$$r^{\text{all-ran}}(F, L_p) = r(e_n^{\text{all-ran}}(F, L_p)) \quad \text{and} \quad r^{\text{std-ran}}(F, L_p) = r(e_n^{\text{std-ran}}(F, L_p)).$$

In particular, we would like to know if it is possible that the sequence $(r^{\text{all-ran}}(F, L_p))$ converges much faster than the sequence $(r^{\text{std-ran}}(F, L_p))$. The main question addressed in this section is to find or estimate the *power* function defined as $\ell^{\text{ran-x}} : (0, \infty) \times [1, \infty] \rightarrow [0, 1]$ by

$$\ell^{\text{ran-x}}(r, p) := \inf_{F: r^{\text{all-ran}}(F, L_p) = r} \frac{r^{\text{std-ran}}(F, L_p)}{r},$$

where $x \in \{H, B\}$ and indicates that the infimum is taken over all Hilbert spaces ($x = H$) or over all Banach spaces ($x = B$) continuously embedded in L_p and the rate of convergence is r when we use arbitrary linear functionals. In the randomized setting we do *not* need to assume that function values are continuous linear functionals.

3.1 Double Hilbert Case

In this subsection we consider the approximation problem defined over a Hilbert space with the error measured also in the Hilbert space L_2 . It may be surprising but the results in the double Hilbert case are complete due to [28], and there is no need to discuss different cases depending on the values of r .

Theorem 6 ([28]). Let $I : H \rightarrow L_2(\Omega)$ be a continuous embedding from a Hilbert space H into $L_2(\Omega)$. Then

$$r^{\text{all-ran}}(H, L_2) = r^{\text{std-ran}}(H, L_2).$$

Therefore

$$\ell^{\text{ran-H}}(r, 2) = 1 \quad \text{for all } r > 0.$$

We add that it was known before, see [13, 25], that also

$$r^{\text{all-ran}}(H, L_2) = r^{\text{all-wor}}(H, L_2).$$

This means that the power of function values in the randomized setting is the same as the power of arbitrary linear functionals in the worst case setting, which in turn is the same as in the randomized setting.

3.2 Other Cases

For $p > 2$, we know examples from the literature where the rate $r^{\text{all-ran}}(H, L_p)$ is larger than the rate $r^{\text{std-ran}}(H, L_p)$. Namely take $I : W_2^r([0, 1]) \rightarrow L_p([0, 1])$. Then with Λ^{all} one can achieve the order n^{-r} (with additional log terms in the case $p = \infty$, but the order is still r), see [10]. For Λ^{std} the optimal order is $n^{-r+1/2-1/p}$, see [2]. The authors of [2, 10] studied the case of integer r , but the results can be extended via interpolation to all $r > 1$. Therefore we obtain

$$\ell^{\text{ran-H}}(r, p) \leq \frac{r - 1/2 + 1/p}{r} \quad \text{if } r \geq 1 \text{ and } p > 2.$$

We summarize these estimates of the power function in the following theorem.

Theorem 7. Let $p > 2$. Then

$$\ell^{\text{ran-H}}(r, p) \leq 1 - \frac{1/2 - 1/p}{r} \quad \text{for all } r \geq 1.$$

Sobolev embeddings in the randomized setting were studied by several authors, including [2, 3, 4, 10, 12, 21, 25]. For our purpose, the most important papers are [2, 10] and the paper [3] for the interpolation argument.

For the embedding $I : W_2^r([0, 1]) \rightarrow L_\infty([0, 1])$ the rate is improved by $1/2$ if we switch from the class Λ^{std} to the class Λ^{all} . This gap of $1/2$ is the largest possible under some additional conditions, see [7, 9]. Let us add in passing that the same gap of $1/2$ appears for Λ^{all} between the worst case and the randomized setting.

The Hilbert case for $p \in [1, 2)$ as well as the Banach case for all $p \in [1, \infty]$ have not yet been studied. We pose this as an open problem.

Open Problem 7. Study the power function in the randomized setting for the Hilbert case with $p \in [1, 2)$ and for the Banach case for all $p \in [1, \infty]$. In particular, determine the supremum $a^*(p)$ of a for which

$$\ell^{\text{ran-H/B}}(r, p) = 0 \quad \text{for all } r \in (0, a].$$

4 The average case setting with a Gaussian measure

In the average case setting we assume that $I : F \rightarrow L_p(\Omega)$ is continuously embedded and function evaluations are continuous functionals on F . As far as we know, only the case $p = 2$ was studied and we report the known results from [5] for this case.

We assume that F is a separable Hilbert/Banach space equipped with a zero mean Gaussian measure μ . As in the worst case setting, we consider deterministic algorithms, and due to general results, see [21], it is enough to compare linear algorithms

$$A_n(f) = \sum_{k=1}^n L_k(f)g_k \quad \text{and} \quad A_n(f) = \sum_{k=1}^n f(x_k)g_k,$$

where $g_k \in L_2(\Omega)$. The average case error of an algorithm is defined by

$$e^{\text{avg}}(A) = \left(\int_F \|f - A(f)\|_p^2 d\mu(f) \right)^{1/p}.$$

As in the other settings, we define the minimal n th average case errors $e_n^{\text{all-avg}}(F, L_p)$, $e_n^{\text{std-avg}}(F, L_p)$ and the power function $\ell^{\text{avg-H/B}}$. That is, for

$$r^{\text{all/std-avg}}(F, L_p) = r(e_n^{\text{all/std-avg}}(F, L_p))$$

we have

$$\ell^{\text{avg-x}}(r, p) := \inf_{F: r^{\text{all-avg}}(F, L_p) = r} \frac{r^{\text{std-avg}}(F, L_p)}{r}.$$

As always, $x \in \{H, B\}$ and we take the infimum over separable Hilbert ($x = H$) or Banach ($x = B$) spaces equipped with zero mean Gaussian measures that are continuously embedded in L_p and for which function values are continuous linear functionals as well as the rate of convergence is r when arbitrary linear functionals are used.

As already mentioned, results are known only for $p = 2$. Then the cases of the Hilbert and Banach spaces are the same due to the presence of Gaussian measures. This follows from the fact that even if F is a separable Banach space then the minimal errors for the class Λ^{all} depend on the Gaussian measure $\nu = \mu I^{-1}$ given by

$$\nu(A) = \mu(\{f \in F \mid I(f) \in A\})$$

for a Borel set A of L_2 . The measure ν is also a zero mean Gaussian measure whose covariance operator $C_\nu : L_2 \rightarrow L_2$ is given by

$$\langle C_\nu f_1, f_2 \rangle_{L_2} = \int_{L_2} \langle f, f_1 \rangle_{L_2} \langle f, f_2 \rangle_{L_2} d\nu(f) \quad \text{for all } f_1, f_2 \in L_2.$$

The operator C_ν is self adjoint, semi positive definite, compact and has a finite trace. That is, its ordered eigenvalues λ_j have a finite sum. It is known that

$$e_n^{\text{all-avg}}(F, L_2) = \left(\sum_{j=n+1}^{\infty} \lambda_j \right)^{1/2}.$$

Similarly as in the randomized setting for the double Hilbert space, the results on the power function are complete and there is no need to discuss different cases of r .

Theorem 8 ([5]). Let $I : F \rightarrow L_2(\Omega)$ be a continuous embedding from a separable Banach space F equipped with a zero mean Gaussian measure μ into the $L_2(\Omega)$. Then

$$r^{\text{all-avg}}(F, L_2) = r^{\text{std-avg}}(F, L_2).$$

Therefore

$$\ell^{\text{avg-H/B}}(r, 2) = 1 \quad \text{for all } r > 0.$$

Of course it would be interesting to study the power function for other values of p . This is posed as our last open problem.

Open Problem 8. Study the power function in the average case setting for $p \neq 2$. In particular, verify whether a similar result as Theorem 8 holds.

Acknowledgment

We appreciate comments on this paper from Stefan Heinrich, Anargyros Papageorgiou, Joseph F. Traub and Grzegorz W. Wasilkowski. We are especially thankful to Stefan Heinrich for pointing out some errors in our previous manuscript.

References

- [1] E. M. Galeev, Linear widths of Hölder-Nikolskii classes of periodic functions of several variables, *Math. Notes* **59**, 133–146, 1996.
- [2] S. Heinrich, Randomized approximation of Sobolev embeddings, in: *Monte Carlo and Quasi-Monte Carlo Methods 2006*, A. Keller, S. Heinrich, H. Niederreiter (eds.), 445–459, Springer, Berlin, 2008.
- [3] S. Heinrich, Randomized approximation of Sobolev embeddings II, *J. Complexity* **25**, 455–472, 2009.

- [4] S. Heinrich, Randomized approximation of Sobolev embeddings III, *J. Complexity* **25**, 473–507, 2009.
- [5] F. Hickernell, G. W. Wasilkowski and H. Woźniakowski, Tractability of linear multivariate problems in the average case setting, in: *Monte Carlo and Quasi-Monte Carlo Methods 2006*, A. Keller, S. Heinrich, H. Niederreiter (eds.), 461–494, Springer, Berlin, 2008.
- [6] A. Hinrichs, E. Novak and J. Vybíral, Linear information versus function evaluations for L_2 -approximation, *J. Approx. Th.* **153**, 97–107, 2008.
- [7] F. Y. Kuo, G. W. Wasilkowski and H. Woźniakowski, Multivariate L_∞ approximation in the worst case setting over reproducing kernel Hilbert spaces, *J. Approx. Th.* **152**, 135–160, 2008.
- [8] F. Y. Kuo, G. W. Wasilkowski and H. Woźniakowski, On the power of standard information for multivariate approximation in the worst case setting, *J. Approx. Th.* **158**, 97–125, 2009.
- [9] F. Y. Kuo, G. W. Wasilkowski and H. Woźniakowski, On the power of standard information for L_∞ approximation in the randomized case setting, *BIT Numer. Math.* **49**, 543–564, 2009.
- [10] P. Mathé, Random approximation of Sobolev embeddings, *J. Complexity* **7**, 261–281, 1991.
- [11] C. A. Micchelli and G. Wahba, Design problems for optimal surface interpolation, in *Approximation Theory and Applications*, Z. Ziegler ed., pp. 329–347, Academic Press, New York, 1981.
- [12] E. Novak, *Deterministic and Stochastic Error Bounds in Numerical Analysis*, LNIM **1349**, Springer-Verlag, Berlin, 1988.
- [13] E. Novak, Optimal linear randomized methods for linear operators in Hilbert spaces, *J. Complexity* **8**, 22–36, 1992.
- [14] E. Novak and H. Woźniakowski, *Tractability of Multivariate Problems*, Volume I: Linear Information, European Math. Soc., Zürich, 2008.
- [15] E. Novak and H. Woźniakowski, *Tractability of Multivariate Problems*, Volume II: Standard Information for Functionals, EMS, Zürich, 2010, to appear.
- [16] A. Pietsch, *Operator Ideals*, North Holland, 1980.
- [17] W. Sickel and T. Ullrich, Spline interpolation on sparse grids, to appear in *Applicable Analysis*.
- [18] R. Tandetzky, Approximation of functions from a Hilbert space using function values or general linear information, in progress, 2010.

- [19] V. N. Temlyakov, *Approximation of Periodic Functions*, Nova Science, New York, 1993.
- [20] V. N. Temlyakov, On approximate recovery of functions with bounded mixed derivative. *J. Complexity* **9**, 41–59, 1993.
- [21] J. F. Traub, G. W. Wasilkowski and H. Woźniakowski, *Information-Based Complexity*, Academic Press, 1988.
- [22] H. Triebel, *Bases in Function Spaces, Sampling, Discrepancy, Numerical Integration*, EMS Publ. House, Zürich, 2010.
- [23] J. Vybíral, Sampling numbers and function spaces, *J. Complexity* **23**, 773–792, 2008.
- [24] J. Vybíral, Widths of embeddings in function spaces, *J. Complexity* **24**, 545–570, 2008.
- [25] G. W. Wasilkowski, Randomization for continuous problems, *J. Complexity* **5**, 195–218, 1989.
- [26] H. Woźniakowski, Tractability and strong tractability of multivariate tensor product problems, *J. of Computing and Information* **4**, 1–19, 1994.
- [27] G. W. Wasilkowski and H. Woźniakowski, On the power of standard information for weighted approximation, *Found. Comput. Math.* **1**, 417–434, 2001.
- [28] G. W. Wasilkowski and H. Woźniakowski, The power of standard information for multivariate approximation in the randomized case setting, *Math. Comp.* **76**, 965–988, 2007.

Preprint Series DFG-SPP 1324

<http://www.dfg-spp1324.de>

Reports

- [1] R. Ramlau, G. Teschke, and M. Zhariy. A Compressive Landweber Iteration for Solving Ill-Posed Inverse Problems. Preprint 1, DFG-SPP 1324, September 2008.
- [2] G. Plonka. The Easy Path Wavelet Transform: A New Adaptive Wavelet Transform for Sparse Representation of Two-dimensional Data. Preprint 2, DFG-SPP 1324, September 2008.
- [3] E. Novak and H. Woźniakowski. Optimal Order of Convergence and (In-) Tractability of Multivariate Approximation of Smooth Functions. Preprint 3, DFG-SPP 1324, October 2008.
- [4] M. Espig, L. Grasedyck, and W. Hackbusch. Black Box Low Tensor Rank Approximation Using Fibre-Crosses. Preprint 4, DFG-SPP 1324, October 2008.
- [5] T. Bonesky, S. Dahlke, P. Maass, and T. Raasch. Adaptive Wavelet Methods and Sparsity Reconstruction for Inverse Heat Conduction Problems. Preprint 5, DFG-SPP 1324, January 2009.
- [6] E. Novak and H. Woźniakowski. Approximation of Infinitely Differentiable Multivariate Functions Is Intractable. Preprint 6, DFG-SPP 1324, January 2009.
- [7] J. Ma and G. Plonka. A Review of Curvelets and Recent Applications. Preprint 7, DFG-SPP 1324, February 2009.
- [8] L. Denis, D. A. Lorenz, and D. Trede. Greedy Solution of Ill-Posed Problems: Error Bounds and Exact Inversion. Preprint 8, DFG-SPP 1324, April 2009.
- [9] U. Friedrich. A Two Parameter Generalization of Lions' Nonoverlapping Domain Decomposition Method for Linear Elliptic PDEs. Preprint 9, DFG-SPP 1324, April 2009.
- [10] K. Bredies and D. A. Lorenz. Minimization of Non-smooth, Non-convex Functionals by Iterative Thresholding. Preprint 10, DFG-SPP 1324, April 2009.
- [11] K. Bredies and D. A. Lorenz. Regularization with Non-convex Separable Constraints. Preprint 11, DFG-SPP 1324, April 2009.

- [12] M. Döhler, S. Kunis, and D. Potts. Nonequispaced Hyperbolic Cross Fast Fourier Transform. Preprint 12, DFG-SPP 1324, April 2009.
- [13] C. Bender. Dual Pricing of Multi-Exercise Options under Volume Constraints. Preprint 13, DFG-SPP 1324, April 2009.
- [14] T. Müller-Gronbach and K. Ritter. Variable Subspace Sampling and Multi-level Algorithms. Preprint 14, DFG-SPP 1324, May 2009.
- [15] G. Plonka, S. Tenorth, and A. Iske. Optimally Sparse Image Representation by the Easy Path Wavelet Transform. Preprint 15, DFG-SPP 1324, May 2009.
- [16] S. Dahlke, E. Novak, and W. Sickel. Optimal Approximation of Elliptic Problems by Linear and Nonlinear Mappings IV: Errors in L_2 and Other Norms. Preprint 16, DFG-SPP 1324, June 2009.
- [17] B. Jin, T. Khan, P. Maass, and M. Pidcock. Function Spaces and Optimal Currents in Impedance Tomography. Preprint 17, DFG-SPP 1324, June 2009.
- [18] G. Plonka and J. Ma. Curvelet-Wavelet Regularized Split Bregman Iteration for Compressed Sensing. Preprint 18, DFG-SPP 1324, June 2009.
- [19] G. Teschke and C. Borries. Accelerated Projected Steepest Descent Method for Nonlinear Inverse Problems with Sparsity Constraints. Preprint 19, DFG-SPP 1324, July 2009.
- [20] L. Grasedyck. Hierarchical Singular Value Decomposition of Tensors. Preprint 20, DFG-SPP 1324, July 2009.
- [21] D. Rudolf. Error Bounds for Computing the Expectation by Markov Chain Monte Carlo. Preprint 21, DFG-SPP 1324, July 2009.
- [22] M. Hansen and W. Sickel. Best m-term Approximation and Lizorkin-Triebel Spaces. Preprint 22, DFG-SPP 1324, August 2009.
- [23] F.J. Hickernell, T. Müller-Gronbach, B. Niu, and K. Ritter. Multi-level Monte Carlo Algorithms for Infinite-dimensional Integration on \mathbb{R}^N . Preprint 23, DFG-SPP 1324, August 2009.
- [24] S. Dereich and F. Heidenreich. A Multilevel Monte Carlo Algorithm for Lévy Driven Stochastic Differential Equations. Preprint 24, DFG-SPP 1324, August 2009.
- [25] S. Dahlke, M. Fornasier, and T. Raasch. Multilevel Preconditioning for Adaptive Sparse Optimization. Preprint 25, DFG-SPP 1324, August 2009.

- [26] S. Dereich. Multilevel Monte Carlo Algorithms for Lévy-driven SDEs with Gaussian Correction. Preprint 26, DFG-SPP 1324, August 2009.
- [27] G. Plonka, S. Tenorth, and D. Roşca. A New Hybrid Method for Image Approximation using the Easy Path Wavelet Transform. Preprint 27, DFG-SPP 1324, October 2009.
- [28] O. Koch and C. Lubich. Dynamical Low-rank Approximation of Tensors. Preprint 28, DFG-SPP 1324, November 2009.
- [29] E. Faou, V. Gradinaru, and C. Lubich. Computing Semi-classical Quantum Dynamics with Hagedorn Wavepackets. Preprint 29, DFG-SPP 1324, November 2009.
- [30] D. Conte and C. Lubich. An Error Analysis of the Multi-configuration Time-dependent Hartree Method of Quantum Dynamics. Preprint 30, DFG-SPP 1324, November 2009.
- [31] C. E. Powell and E. Ullmann. Preconditioning Stochastic Galerkin Saddle Point Problems. Preprint 31, DFG-SPP 1324, November 2009.
- [32] O. G. Ernst and E. Ullmann. Stochastic Galerkin Matrices. Preprint 32, DFG-SPP 1324, November 2009.
- [33] F. Lindner and R. L. Schilling. Weak Order for the Discretization of the Stochastic Heat Equation Driven by Impulsive Noise. Preprint 33, DFG-SPP 1324, November 2009.
- [34] L. Kämmerer and S. Kunis. On the Stability of the Hyperbolic Cross Discrete Fourier Transform. Preprint 34, DFG-SPP 1324, December 2009.
- [35] P. Cerejeiras, M. Ferreira, U. Kähler, and G. Teschke. Inversion of the noisy Radon transform on $SO(3)$ by Gabor frames and sparse recovery principles. Preprint 35, DFG-SPP 1324, January 2010.
- [36] T. Jahnke and T. Udrescu. Solving Chemical Master Equations by Adaptive Wavelet Compression. Preprint 36, DFG-SPP 1324, January 2010.
- [37] P. Kittipoom, G. Kutyniok, and W.-Q Lim. Irregular Shearlet Frames: Geometry and Approximation Properties. Preprint 37, DFG-SPP 1324, February 2010.
- [38] G. Kutyniok and W.-Q Lim. Compactly Supported Shearlets are Optimally Sparse. Preprint 38, DFG-SPP 1324, February 2010.
- [39] M. Hansen and W. Sickel. Best m -Term Approximation and Tensor Products of Sobolev and Besov Spaces – the Case of Non-compact Embeddings. Preprint 39, DFG-SPP 1324, March 2010.

- [40] B. Niu, F.J. Hickernell, T. Müller-Gronbach, and K. Ritter. Deterministic Multi-level Algorithms for Infinite-dimensional Integration on \mathbb{R}^N . Preprint 40, DFG-SPP 1324, March 2010.
- [41] P. Kittipoom, G. Kutyniok, and W.-Q Lim. Construction of Compactly Supported Shearlet Frames. Preprint 41, DFG-SPP 1324, March 2010.
- [42] C. Bender and J. Steiner. Error Criteria for Numerical Solutions of Backward SDEs. Preprint 42, DFG-SPP 1324, April 2010.
- [43] L. Grasedyck. Polynomial Approximation in Hierarchical Tucker Format by Vector-Tensorization. Preprint 43, DFG-SPP 1324, April 2010.
- [44] M. Hansen und W. Sickel. Best m -Term Approximation and Sobolev-Besov Spaces of Dominating Mixed Smoothness - the Case of Compact Embeddings. Preprint 44, DFG-SPP 1324, April 2010.
- [45] P. Binev, W. Dahmen, and P. Lamby. Fast High-Dimensional Approximation with Sparse Occupancy Trees. Preprint 45, DFG-SPP 1324, May 2010.
- [46] J. Ballani and L. Grasedyck. A Projection Method to Solve Linear Systems in Tensor Format. Preprint 46, DFG-SPP 1324, May 2010.
- [47] P. Binev, A. Cohen, W. Dahmen, R. DeVore, G. Petrova, and P. Wojtaszczyk. Convergence Rates for Greedy Algorithms in Reduced Basis Methods. Preprint 47, DFG-SPP 1324, May 2010.
- [48] S. Kestler and K. Urban. Adaptive Wavelet Methods on Unbounded Domains. Preprint 48, DFG-SPP 1324, June 2010.
- [49] H. Yserentant. The Mixed Regularity of Electronic Wave Functions Multiplied by Explicit Correlation Factors. Preprint 49, DFG-SPP 1324, June 2010.
- [50] H. Yserentant. On the Complexity of the Electronic Schrödinger Equation. Preprint 50, DFG-SPP 1324, June 2010.
- [51] M. Guillemard and A. Iske. Curvature Analysis of Frequency Modulated Manifolds in Dimensionality Reduction. Preprint 51, DFG-SPP 1324, June 2010.
- [52] E. Herrholz and G. Teschke. Compressive Sensing Principles and Iterative Sparse Recovery for Inverse and Ill-Posed Problems. Preprint 52, DFG-SPP 1324, July 2010.
- [53] L. Kämmerer, S. Kunis, and D. Potts. Interpolation Lattices for Hyperbolic Cross Trigonometric Polynomials. Preprint 53, DFG-SPP 1324, July 2010.

- [54] G. Kutyniok and W.-Q Lim. Shearlets on Bounded Domains. Preprint 54, DFG-SPP 1324, July 2010.
- [55] A. Zeiser. Wavelet Approximation in Weighted Sobolev Spaces of Mixed Order with Applications to the Electronic Schrödinger Equation. Preprint 55, DFG-SPP 1324, July 2010.
- [56] G. Kutyniok, J. Lemvig, and W.-Q Lim. Compactly Supported Shearlets. Preprint 56, DFG-SPP 1324, July 2010.
- [57] A. Zeiser. On the Optimality of the Inexact Inverse Iteration Coupled with Adaptive Finite Element Methods. Preprint 57, DFG-SPP 1324, July 2010.
- [58] S. Jokar. Sparse Recovery and Kronecker Products. Preprint 58, DFG-SPP 1324, August 2010.
- [59] T. Aboiyar, E. H. Georgoulis, and A. Iske. Adaptive ADER Methods Using Kernel-Based Polyharmonic Spline WENO Reconstruction. Preprint 59, DFG-SPP 1324, August 2010.
- [60] O. G. Ernst, A. Mugler, H.-J. Starkloff, and E. Ullmann. On the Convergence of Generalized Polynomial Chaos Expansions. Preprint 60, DFG-SPP 1324, August 2010.
- [61] S. Holtz, T. Rohwedder, and R. Schneider. On Manifolds of Tensors of Fixed TT-Rank. Preprint 61, DFG-SPP 1324, September 2010.
- [62] J. Ballani, L. Grasedyck, and M. Kluge. Black Box Approximation of Tensors in Hierarchical Tucker Format. Preprint 62, DFG-SPP 1324, October 2010.
- [63] M. Hansen. On Tensor Products of Quasi-Banach Spaces. Preprint 63, DFG-SPP 1324, October 2010.
- [64] S. Dahlke, G. Steidl, and G. Teschke. Shearlet Coorbit Spaces: Compactly Supported Analyzing Shearlets, Traces and Embeddings. Preprint 64, DFG-SPP 1324, October 2010.
- [65] W. Hackbusch. Tensorisation of Vectors and their Efficient Convolution. Preprint 65, DFG-SPP 1324, November 2010.
- [66] P. A. Cioica, S. Dahlke, S. Kinzel, F. Lindner, T. Raasch, K. Ritter, and R. L. Schilling. Spatial Besov Regularity for Stochastic Partial Differential Equations on Lipschitz Domains. Preprint 66, DFG-SPP 1324, November 2010.

- [67] E. Novak and H. Woźniakowski. On the Power of Function Values for the Approximation Problem in Various Settings. Preprint 67, DFG-SPP 1324, November 2010.