

DFG-Schwerpunktprogramm 1324

„Extraktion quantifizierbarer Information aus komplexen Systemen“

Piecewise Tensor Product Wavelet Bases by Extensions and Approximation Rates

N. Chegini, S. Dahlke, U. Friedrich, R. Stevenson

Preprint 101



Edited by

AG Numerik/Optimierung
Fachbereich 12 - Mathematik und Informatik
Philipps-Universität Marburg
Hans-Meerwein-Str.
35032 Marburg

DFG-Schwerpunktprogramm 1324

„Extraktion quantifizierbarer Information aus komplexen Systemen“

Piecewise Tensor Product Wavelet Bases by Extensions and Approximation Rates

N. Chegini, S. Dahlke, U. Friedrich, R. Stevenson

Preprint 101



The consecutive numbering of the publications is determined by their chronological order.

The aim of this preprint series is to make new research rapidly available for scientific discussion. Therefore, the responsibility for the contents is solely due to the authors. The publications will be distributed by the authors.

PIECEWISE TENSOR PRODUCT WAVELET BASES BY EXTENSIONS AND APPROXIMATION RATES

NABI CHEGINI, STEPHAN DAHLKE, ULRICH FRIEDRICH, ROB STEVENSON

ABSTRACT. Following [*Studia Math.*, 76(2) (1983), pp. 1–58 and 95–136] by Z. Ciesielski and T. Figiel and [*SIAM J. Math. Anal.*, 31 (1999), pp. 184–230] by W. Dahmen and R. Schneider, by the application of extension operators we construct a basis for a range of Sobolev spaces on a domain Ω from corresponding bases on subdomains that form a non-overlapping decomposition. As subdomains, we take hypercubes, or smooth parametric images of those, and equip them with tensor product wavelet bases. We prove approximation rates from the resulting piecewise tensor product basis that are independent of the spatial dimension of Ω . For two- and three dimensional polytopes we show that the solution of Poisson type problems satisfies the required regularity condition. The dimension independent rates will be realized numerically in linear complexity by the application of the adaptive wavelet-Galerkin scheme.

1. INTRODUCTION

Let $\Omega = \cup_{k=0}^N \Omega_k \subset \mathbb{R}^n$ be a non-overlapping domain decomposition. By the use of extension operators, we will construct isomorphisms between the Cartesian product of Sobolev spaces on the subdomains, which incorporate suitable boundary conditions, and Sobolev spaces on Ω . By applying such an isomorphism to the union of Riesz bases for the Sobolev spaces on the subdomains, the result is a Riesz basis for the Sobolev space on Ω .

Since the approach can be applied recursively, to understand the construction of such an isomorphism, it is sufficient to consider the case of having two subdomains. For $i \in \{1, 2\}$, let R_i be the restriction of functions on Ω to Ω_i , let η_2 be the extension by zero of functions on Ω_2 to functions on Ω , and let E_1 be some extension of functions on Ω_1 to functions on Ω which, for some $m \in \mathbb{N}_0$, is bounded from $H^m(\Omega_1)$ to the target space $H^m(\Omega)$. Then $\begin{bmatrix} R_1 \\ R_2(\text{Id} - E_1 R_1) \end{bmatrix} : H^m(\Omega) \rightarrow H^m(\Omega_1) \times H_{0, \partial\Omega_1 \cap \partial\Omega_2}^m(\Omega_2)$ is boundedly invertible with inverse $[E_1 \quad \eta_2]$, see Figure 1 ($H_{0, \partial\Omega_1 \cap \partial\Omega_2}^m(\Omega_2)$ is the space of $H^m(\Omega_2)$ functions that vanish up to order $m - 1$ at $\partial\Omega_1 \cap \partial\Omega_2$). Consequently, if Ψ_1 is a Riesz basis for $H^m(\Omega_1)$ and Ψ_2 is a Riesz basis for $H_{0, \partial\Omega_1 \cap \partial\Omega_2}^m(\Omega_2)$, then $E_1\Psi_1 \cup \eta_2\Psi_2$ is a Riesz basis for $H^m(\Omega)$.


The principle to construct a basis for a function space on Ω by applying an isomorphism between this space and the product of corresponding function spaces

Date: September 2, 2011.

2000 Mathematics Subject Classification. 15A69, 35B65, 41A25, 41A63, 42C40, 65N12, 65T60.

Key words and phrases. Wavelets, tensor product approximation, domain decomposition, extension operators, weighted anisotropic Sobolev space, regularity, adaptive wavelet scheme, best approximation rates.

The first author was supported by the Netherlands Organization for Scientific Research (NWO) under contract no. 613.000.902.

FIGURE 1. Splitting of u into a sum of functions on the subdomains.

on non-overlapping subdomains was introduced in [CF83]. In [DS99] (see also [KS06]), this idea was revisited with the aim to practically construct such a basis for doing computations, rather than to show its existence.

In addition to the findings from [DS99], in the current work we derive precise conditions on the ordering of the subdomains so that the corresponding “true” extension operators (not the trivial zero extensions), being the building blocks of the isomorphism, actually do exist as bounded mappings. To explain this, as an example, consider the construction of a basis for $H^1(\Omega)$, where Ω is an L-shaped domain subdivided into 3 subdomains as illustrated in Figure 2. Choosing the




FIGURE 2. A non-feasible configuration

action of the extension operators as illustrated by the subsequent arrows yields an invalid configuration, i.e., no bounded operators exist. Indeed, in view of the second extension from Ω_3 to Ω , in the first step we have to construct a Riesz basis for $H_{0,\partial\Omega_2\cap\partial\Omega_3}^1(\Omega_1 \cup \Omega_2)$, meaning that the first extension should be bounded from $H^1(\Omega_1)$ to $H_{0,\partial\Omega_2\cap\partial\Omega_3}^1(\Omega_1 \cup \Omega_2)$. In view of the boundary condition incorporated in the latter space, this is, however, impossible. On the other hand, by reversing one of the two arrows, one does obtain a valid configuration.

The conditions on the directions of the arrows will be different when homogeneous boundary conditions are imposed on the outer boundary, e.g., when a basis for $H_0^1(\Omega)$ is sought.

Our main interest in the construction of a basis from bases from subdomains lies in the use of *piecewise tensor product approximation*. On \square one can construct a basis for the Sobolev space $H^m(\square)$ (or for a subspace incorporating Dirichlet boundary conditions) by taking an n -fold tensor product of a collection of univariate functions that forms a Riesz basis for $L_2(0, 1)$ as well as, properly scaled, for $H^m(0, 1)$. Thinking of a univariate wavelet basis of order $d > m$, the advantage of this approach is that the rate of nonlinear best M -term approximation of a sufficiently smooth function u is $d - m$, compared to $\frac{d-m}{n}$ for standard best M -term isotropic

(wavelet) approximation of order d on \square . The division of the “one-dimensional” rate $d - m$ by n is commonly referred to as the *curse of dimensionality*.

One may argue that for any fixed n , a rate $d - m$ can also be obtained by isotropic approximation by increasing the order from d to $nd - (n - 1)m$. Concerning the required smoothness of u , however, in the latter case it is (essentially) necessary and sufficient that for $1 \leq i \leq n$, $0 \leq k \leq m$, it holds that $\partial^\alpha \partial_i^k u \in L_p(\square)$ for $p = (d - m + \frac{1}{2})^{-1}$ and $|\alpha|_1 \leq n(d - m)$, whereas with tensor approximation the last condition reads as the much milder one $|\alpha|_\infty \leq d - m$ (cf. [Nit06, SU09] for details).

Actually, the above conditions guarantee only any rate $s < d - m$. Interpolation arguments that are used do not give a result for the “endpoint” $s = d - m$.

In any case, for dimensions $n \geq 3$, the solution of an elliptic boundary value problem of order $2m = 2$ generally does *not* satisfy the conditions such that isotropic approximation converges with the best, or any near best possible rate allowed by the polynomial order, i.e., $\frac{d-m}{n}$ for order d . In order to achieve this rate, generally anisotropic approximation is mandatory (cf. [Ape99]).

In addition to avoiding the curse of dimensionality, the possibility of anisotropic approximation is automatically included in (adaptive) tensor product approximation. In [DS10], see also [Nit05], it was shown that best approximations of u from a suitably chosen nested sequence of spaces spanned by tensor product wavelets realizes the best possible rate $d - m$, so not only any near best possible rate, when for $1 \leq i \leq n$, $0 \leq k \leq m$ and $|\alpha|_\infty \leq d - m$, $\partial^\alpha \partial_i^k u$ is in a *weighted* $L_2(\square)$ space, with weight being an n -fold product of univariate weights on $(0, 1)$ that vanish near the endpoints. Clearly, the optimal rate $d - m$ for this linear approximation scheme implies this rate for the nonlinear best M -term approximation from the tensor product basis. What is more, in [DS10] it was shown that with a sufficiently smooth right-hand side, the solution of Poisson’s problem on the n -dimensional unit cube \square satisfies this regularity condition.

In view of these results on \square , we consider a domain Ω whose closure is the union of n -cubes from $\mathbb{Z}^n + \square$, with $\square := (0, 1)^n$, or a domain Ω that is a parametric image of such a domain under a piecewise smooth, globally C^{m-1} diffeomorphism κ . We equip $H^m(\Omega)$ (or a subspace incorporating Dirichlet boundary conditions) with a Riesz basis constructed from tensor product wavelet bases of order d on the subdomains using extension operators as discussed before. Our restriction to decompositions of Ω into subdomains from a topological Cartesian grid will allow us to rely on univariate extensions. We will show the best possible approximation rate $d - m$ for any u that is *piecewise* in the aforementioned weighted Sobolev spaces. The latter turns out to be technically hard. Indeed, in order to end up with locally supported wavelets, we will apply local, scale-dependent extension operators – i.e., only wavelets that are non-zero near an interface will be extended, – which do not preserve more smoothness than essentially membership of H^m .

Furthermore, using anisotropic regularity results recently shown in [CDN10], we show that if, additionally, Ω is a two- or, more interesting, a three-dimensional polytope, then for a sufficiently smooth right-hand side, the solution of Poisson’s problem satisfies this piecewise regularity condition. For that to hold in three dimensions, it will be needed that κ is piecewise trilinear, and it may require a refinement of the initial decomposition of Ω .

Since it defines a boundedly invertible mapping from a Hilbert space, being $H_0^1(\Omega)$, to its dual, the Poisson problem is an example of a well-posed operator equation. Equipping $H_0^1(\Omega)$ with a Riesz basis constructed using extension operators from tensor product wavelet bases of order d on the subdomains, the operator equation is equivalently formulated as a boundedly invertible bi-infinite matrix vector equation. Approximate solutions produced by the adaptive wavelet-Galerkin method ([CDD01, Ste09]) were proven to converge with the best possible rate in linear complexity. We perform numerical tests in two and three dimensions with wavelets of order $d = 5$ that confirm that this rate is $d - m$.

This paper is organized as follows: In Sect. 2, we present the abstract idea behind the construction of isomorphisms between Sobolev spaces on a domain and the product of corresponding Sobolev spaces on subdomains that form a non-overlapping decomposition.

In Sect. 3, we recall results on tensor product approximation on a hypercube, and collect assumptions on the univariate wavelets, being the building blocks of the tensor product wavelets.

In Sect. 4, we consider a domain Ω that is the union of hypercubes from a Cartesian grid of hypercubes. We formulate precise conditions on the order in which univariate extensions over interfaces are applied, and which boundary conditions have to be imposed, so that the composition of these extensions is an isomorphism between a range of Sobolev spaces on Ω and the product of the corresponding Sobolev spaces on the collection of hypercubes. Equipping these hypercubes with tensor product wavelet bases, we end up with a piecewise tensor product wavelet basis on Ω .

In order to obtain locally supported primal and dual wavelets, in Sect. 5 the extension operators are replaced by scale-dependent modifications, in the sense that only wavelets with supports “near” the interfaces are extended. It is shown that approximation from the resulting piecewise tensor product basis gives rise to rates that are independent of the spatial dimension, assuming the function that is approximated satisfies some mild, piecewise weighted Sobolev smoothness conditions.

In Sect. 6, these regularity conditions are verified for the solution of Poisson’s problem with sufficiently smooth right-hand side in two and three-dimensional polytopes.

The best possible rates from the piecewise tensor product basis can be realized in linear complexity by the application of the adaptive wavelet-Galerkin scheme. In Sect. 7, we present some numerical results obtained with this scheme.

2. CONSTRUCTION OF THE ISOMORPHISMS

In an abstract setting, for a class of mappings between a Banach space and the Cartesian product of two other Banach spaces, we give conditions on such mappings to be isomorphisms. The results will be applied to construct isomorphisms between a Sobolev space on a domain and the product of Sobolev spaces on subdomains.

Proposition 2.1. *For normed linear spaces V and V_i ($i = 1, 2$), let $E_1 \in B(V_1, V)$, $\eta_2 \in B(V_2, V)$, $R_1 \in B(V, V_1)$, and $R_2 \in B(\mathfrak{S}\eta_2, V_2)$ be such that*

$$R_1 E_1 = \text{Id}, \quad R_2 \eta_2 = \text{Id}, \quad R_1 \eta_2 = 0, \quad \mathfrak{S}(\text{Id} - E_1 R_1) \subset \mathfrak{S}\eta_2.$$

Then

$$E = [E_1 \quad \eta_2] \in B(V_1 \times V_2, V) \text{ is boundedly invertible,}$$

with inverse

$$E^{-1} = \begin{bmatrix} R_1 \\ R_2(\text{Id} - E_1 R_1) \end{bmatrix}.$$

Proof. Using that $R_1 E_1 = \text{Id}$, $R_1 \eta_2 = 0$, $R_2 \eta_2 = \text{Id}$, we have

$$\begin{bmatrix} R_1 \\ R_2(\text{Id} - E_1 R_1) \end{bmatrix} [E_1 \quad \eta_2] = \begin{bmatrix} \text{Id} & 0 \\ 0 & \text{Id} \end{bmatrix},$$

and using that $\mathfrak{S}(\text{Id} - E_1 R_1) \subset \mathfrak{S}\eta_2$ and $R_2 \eta_2 = \text{Id}$, we have

$$[E_1 \quad \eta_2] \begin{bmatrix} R_1 \\ R_2(\text{Id} - E_1 R_1) \end{bmatrix} = E_1 R_1 + \eta_2 R_2(\text{Id} - E_1 R_1) = \text{Id}. \quad \square$$

In applications V (V_i) will be densely embedded in a Hilbert space H (H_i). Questions about boundedness of E or E^{-1} in dual spaces then reduce to properties of the Hilbert adjoint of E . Study of the Hilbert adjoint will also be relevant for the investigation of dual bases.

Proposition 2.2. *For Hilbert spaces H and H_i ($i = 1, 2$), let $R_i \in B(H, H_i)$, and isometries $\eta_i \in B(H_i, H)$ be such that*

$$R_i \eta_j = \delta_{ij} \quad (i, j \in \{1, 2\}), \quad H = \mathfrak{S}\eta_1 \oplus^\perp \mathfrak{S}\eta_2,$$

and let $E_1 \in B(H_1, H)$ be such that $R_1 E_1 = \text{Id}$.

Then $\eta_1 R_1 + \eta_2 R_2 = \text{Id}$, $E \in B(H_1 \times H_2, H)$ is boundedly invertible, $\eta_i^* = R_i$, and

$$E^* = \begin{bmatrix} E_1^* \\ R_2 \end{bmatrix}, \quad E^{-*} = [\eta_1 \quad (\text{Id} - \eta_1 E_1^*) \eta_2].$$

Proof. The first statement follows from $\eta_1 R_1 + \eta_2 R_2 = \text{Id}$ on $\mathfrak{S}\eta_i$. The second statement follows from Proposition 2.1 once we have verified that $\mathfrak{S}(\text{Id} - E_1 R_1) \subset \mathfrak{S}\eta_2$. Writing $(\text{Id} - E_1 R_1)x = \eta_1 x_1 + \eta_2 x_2$, and applying R_1 to both sides, we find $x_1 = 0$ as required. For any $u \in H_i$, $v \in H$,

$$\langle \eta_i u, v \rangle_H = \langle \eta_i u, \sum_j \eta_j R_j v \rangle_H = \langle \eta_i u, \eta_i R_i v \rangle_H = \langle u, R_i v \rangle_{H_i},$$

or $\eta_i^* = R_i$. Now the last statement follows from the formulas for E and E^{-1} given in Proposition 2.1. \square

Remark 2.3. The formulas for E and E^{-*} , and so those for E^{-1} and E^* are symmetric, with reversed roles of H_1 and H_2 , in the sense that with $E_2 := (\text{Id} - \eta_1 E_1^*) \eta_2$, it holds that $(\text{Id} - \eta_2 E_2^*) \eta_1 = E_1$.

Let \tilde{V} and \tilde{V}_i ($i = 1, 2$) be reflexive Banach spaces with

$$\tilde{V} \hookrightarrow H, \quad \tilde{V}_i \hookrightarrow H_i \text{ with dense embeddings.}$$

In this setting, we have that boundedness, or bounded invertibility of

$$E : \tilde{V}'_1 \times \tilde{V}'_2 \rightarrow \tilde{V}'$$

is equivalent to boundedness, or to bounded invertibility of

$$E^* : \tilde{V} \rightarrow \tilde{V}_1 \times \tilde{V}_2.$$

Proposition 2.4. *Let the assumptions of Proposition 2.2 be valid. Let*

$$R_2 \in B(\tilde{V}, \tilde{V}_2), \quad \eta_1 \in B(\tilde{V}_1, \tilde{V}), \quad E_1^* \in B(\tilde{V}, \tilde{V}_1).$$

Then $E^ \in B(\tilde{V}, \tilde{V}_1 \times \tilde{V}_2)$, and so $E \in B(\tilde{V}'_1 \times \tilde{V}'_2, \tilde{V}')$, is boundedly invertible if and only if R_2 has a right-inverse $\hat{E}_2 \in B(\tilde{V}_2, \tilde{V})$.*

Proof. The assumptions imply that $E^* \in B(\tilde{V}, \tilde{V}_1 \times \tilde{V}_2)$, and that for $E^{-*} \in B(\tilde{V}_1 \times \tilde{V}_2, \tilde{V})$ it suffices to show that $E_2 := (\text{Id} - \eta_1 E_1^*)\eta_2 \in B(\tilde{V}_2, \tilde{V})$. If the latter is true, then, since $R_2 E_2 = \text{Id}$, we can take $\hat{E}_2 = E_2$.

Conversely, let $\hat{E}_2 \in B(\tilde{V}_2, \tilde{V})$ be a right-inverse of R_2 . We have that

$$\begin{aligned} R_1(\text{Id} - E_2 R_2) &= R_1 - R_1 \eta_2 R_2 + R_1 \eta_1 E_1^* \eta_2 R_2 = R_1 + E_1^* \eta_2 R_2 \\ &= E_1^*(\eta_1 R_1 + \eta_2 R_2) = E_1^* \in B(\tilde{V}, \tilde{V}_1). \end{aligned}$$

So

$$\text{Id} - E_2 R_2 = (\eta_1 R_1 + \eta_2 R_2)(\text{Id} - E_2 R_2) = \eta_1 R_1(\text{Id} - E_2 R_2) \in B(\tilde{V}, \tilde{V}),$$

or $E_2 R_2 \in B(\tilde{V}, \tilde{V})$. But then $E_2 = E_2 R_2 \hat{E}_2 \in B(\tilde{V}_2, \tilde{V})$. \square

Finally in this section, we apply interpolation arguments to conclude boundedness of E in scales of Banach spaces.

Proposition 2.5. (a). *Let V, \underline{V} , and V_i, \underline{V}_i ($i = 1, 2$) be Banach spaces with*

$$\underline{V} \hookrightarrow V, \quad \underline{V}_i \hookrightarrow V_i \quad \text{with dense embeddings.}$$

Let the mappings (R_1, R_2, E_1, η_2) satisfy the conditions from Proposition 2.1 for both triples (V, V_1, V_2) and $(\underline{V}, \underline{V}_1, \underline{V}_2)$. Then for $s \in [0, 1]$, $q \in [1, \infty]$,

$$E \in B([V_1, \underline{V}_1]_{s,q} \times [V_2, \underline{V}_2]_{s,q}, [V, \underline{V}]_{s,q}) \text{ is boundedly invertible.}$$

(b). *Let $\tilde{V}, \underline{\tilde{V}}$, and $\tilde{V}_i, \underline{\tilde{V}}_i$ be reflexive Banach spaces, and H and H_i be Hilbert spaces ($i = 1, 2$) with*

$$\underline{\tilde{V}} \hookrightarrow \tilde{V} \hookrightarrow H, \quad \underline{\tilde{V}}_i \hookrightarrow \tilde{V}_i \hookrightarrow H_i \quad \text{with dense embeddings.}$$

Let the conditions of Proposition 2.2 be satisfied, as well as the conditions of Proposition 2.4 for both triples $(\tilde{V}, \tilde{V}_1, \tilde{V}_2)$ and $(\underline{\tilde{V}}, \underline{\tilde{V}}_1, \underline{\tilde{V}}_2)$. Then for $s \in [0, 1]$, $q \in [1, \infty]$,

$$E \in B([\tilde{V}_1, \underline{\tilde{V}}_1]_{s,q}' \times [\tilde{V}_2, \underline{\tilde{V}}_2]_{s,q}', [\tilde{V}, \underline{\tilde{V}}]_{s,q}') \text{ is boundedly invertible.}$$

3. APPROXIMATION BY TENSOR PRODUCT WAVELETS ON THE HYPERCUBE

We will study non-overlapping domain decompositions, where the subdomains are either unit n -cubes or smooth images of those. Sobolev spaces on these cubes, that appear with the construction of a Riesz basis for a Sobolev space on the domain as a whole, will be equipped with tensor product wavelet bases. From [DS10], we recall the construction of those bases, as well as results on the rate of approximation from spans of suitably chosen subsets of these bases.

For $t \in [0, \infty) \setminus (\mathbb{N}_0 + \{\frac{1}{2}\})$ and $\vec{\sigma} = (\sigma_\ell, \sigma_r) \in \{0, \dots, [t + \frac{1}{2}]\}^2$, with $\mathcal{I} := (0, 1)$, let

$$H_{\vec{\sigma}}^t(\mathcal{I}) := \{v \in H^t(\mathcal{I}) : v(0) = \dots = v^{(\sigma_\ell - 1)}(0) = 0 = v(1) = \dots = v^{(\sigma_r - 1)}(1)\}.$$

Remark 3.1. Later, we will use this definition also with \mathcal{I} reading as a general non-empty interval, with 0 and 1 reading as its left and right boundary.

For t and $\vec{\sigma}$ as above, and for $\tilde{t} \in [0, \infty) \setminus (\mathbb{N}_0 + \{\frac{1}{2}\})$ and $\vec{\sigma} = (\tilde{\sigma}_\ell, \tilde{\sigma}_r) \in \{0, \dots, [\tilde{t} + \frac{1}{2}]\}^2$, we assume *univariate wavelet* collections

$$\Psi_{\vec{\sigma}, \vec{\sigma}} := \{\psi_\lambda^{(\vec{\sigma}, \vec{\sigma})} : \lambda \in \nabla_{\vec{\sigma}, \vec{\sigma}}\} \subset H_{\vec{\sigma}}^t(\mathcal{I})$$

such that

$$(W_1) \quad \Psi_{\vec{\sigma}, \vec{\sigma}} \text{ is a Riesz basis for } L_2(\mathcal{I}),$$

$$(W_2) \quad \{2^{-|\lambda|t} \psi_\lambda^{(\vec{\sigma}, \vec{\sigma})} : \lambda \in \nabla_{\vec{\sigma}, \vec{\sigma}}\} \text{ is a Riesz basis for } H_{\vec{\sigma}}^t(\mathcal{I}),$$

where $|\lambda| \in \mathbb{N}_0$ denotes the *level* of λ . Denoting the dual basis of $\Psi_{\vec{\sigma}, \vec{\sigma}}$ for $L_2(\mathcal{I})$ as

$$\tilde{\Psi}_{\vec{\sigma}, \vec{\sigma}} := \{\tilde{\psi}_\lambda^{(\vec{\sigma}, \vec{\sigma})} : \lambda \in \nabla_{\vec{\sigma}, \vec{\sigma}}\}, \text{ furthermore we assume that}$$

$$(W_3) \quad \{2^{-|\lambda|\tilde{t}} \tilde{\psi}_\lambda^{(\vec{\sigma}, \vec{\sigma})} : \lambda \in \nabla_{\vec{\sigma}, \vec{\sigma}}\} \text{ is a Riesz basis for } H_{\vec{\sigma}}^{\tilde{t}}(\mathcal{I}),$$

and that for some

$$\mathbb{N} \ni d > t,$$

$$(W_4) \quad |\langle \tilde{\psi}_\lambda^{(\vec{\sigma}, \vec{\sigma})}, u \rangle_{L_2(\mathcal{I})}| \lesssim 2^{-|\lambda|d} \|u\|_{H^d(\text{supp } \tilde{\psi}^{(\vec{\sigma}, \vec{\sigma})})} \quad (u \in H^d(\mathcal{I}) \cap H_{\vec{\sigma}}^t(\mathcal{I})),$$

$$(W_5) \quad \rho := \sup_{\lambda \in \nabla_{\vec{\sigma}, \vec{\sigma}}} 2^{|\lambda|} \max(\text{diam supp } \tilde{\psi}_\lambda^{(\vec{\sigma}, \vec{\sigma})}, \text{diam supp } \psi_\lambda^{(\vec{\sigma}, \vec{\sigma})}) \\ \simeq \inf_{\lambda \in \nabla_{\vec{\sigma}, \vec{\sigma}}} 2^{|\lambda|} \max(\text{diam supp } \tilde{\psi}_\lambda^{(\vec{\sigma}, \vec{\sigma})}, \text{diam supp } \psi_\lambda^{(\vec{\sigma}, \vec{\sigma})}),$$

$$(W_6) \quad \sup_{j, k \in \mathbb{N}_0} \#\{|\lambda| = j : [k2^{-j}, (k+1)2^{-j}] \cap (\text{supp } \tilde{\psi}_\lambda^{(\vec{\sigma}, \vec{\sigma})} \cup \text{supp } \psi_\lambda^{(\vec{\sigma}, \vec{\sigma})}) \neq \emptyset\} < \infty.$$

The conditions (W_5) and (W_6) will be referred to by saying that *both primal and dual wavelets* are *local* or *locally finite*, respectively. For some arguments, it will be used that by increasing the coarsest scale, the constant ρ can always be assumed to be sufficiently small.

With, for $n \in \mathbb{N}$,

$$\square := \mathcal{I}^n,$$

one has $L_2(\square) = \otimes_{i=1}^n L_2(\mathcal{I})$. For

$$\boldsymbol{\sigma} = (\vec{\sigma}_i = ((\sigma_i)_\ell, (\sigma_i)_r))_{1 \leq i \leq n} \in (\{0, \dots, [t + \frac{1}{2}]\}^2)^n,$$

we define

$$H_{\boldsymbol{\sigma}}^t(\square) := H_{\vec{\sigma}_1}^t(\mathcal{I}) \otimes L_2(\mathcal{I}) \otimes \dots \otimes L_2(\mathcal{I}) \cap \dots \cap L_2(\mathcal{I}) \otimes \dots \otimes L_2(\mathcal{I}) \otimes H_{\vec{\sigma}_n}^t(\mathcal{I}),$$

which is the space of $H^t(\square)$ -functions whose normal derivatives of up to orders $(\sigma_i)_\ell$ and $(\sigma_i)_r$ vanish at the facets $\bar{\mathcal{I}}^{i-1} \times \{0\} \times \bar{\mathcal{I}}^{n-i}$ and $\bar{\mathcal{I}}^{i-1} \times \{1\} \times \bar{\mathcal{I}}^{n-i}$, respectively ($1 \leq i \leq n$) (the proof of this fact given in [DS10] for $t \in \mathbb{N}_0$ can be generalized to $t \in [0, \infty) \setminus (\mathbb{N}_0 + \frac{1}{2})$).

The *tensor product wavelet* collection

$$\Psi_{\boldsymbol{\sigma}, \vec{\sigma}} := \otimes_{i=1}^n \Psi_{\vec{\sigma}_i, \vec{\sigma}_i} = \{\psi_\lambda^{(\boldsymbol{\sigma}, \vec{\sigma})} := \otimes_{i=1}^n \psi_{\lambda_i}^{(\vec{\sigma}_i, \vec{\sigma}_i)} : \lambda \in \nabla_{\boldsymbol{\sigma}, \vec{\sigma}} := \prod_{i=1}^n \nabla_{\vec{\sigma}_i, \vec{\sigma}_i}\},$$

and its renormalized version $\{(\sum_{i=1}^n 4^{t|\lambda_i|})^{-1/2} \psi_\lambda^{(\boldsymbol{\sigma}, \vec{\sigma})} : \lambda \in \nabla_{\boldsymbol{\sigma}, \vec{\sigma}}\}$ are Riesz bases for $L_2(\square)$ and $H_{\boldsymbol{\sigma}}^t(\square)$, respectively. The collection that is dual to $\Psi_{\boldsymbol{\sigma}, \vec{\sigma}}$ reads as

$$\tilde{\Psi}_{\boldsymbol{\sigma}, \vec{\sigma}} := \otimes_{i=1}^n \tilde{\Psi}_{\vec{\sigma}_i, \vec{\sigma}_i} = \{\tilde{\psi}_\lambda^{(\boldsymbol{\sigma}, \vec{\sigma})} := \otimes_{i=1}^n \tilde{\psi}_{\lambda_i}^{(\vec{\sigma}_i, \vec{\sigma}_i)} : \lambda \in \nabla_{\boldsymbol{\sigma}, \vec{\sigma}}\},$$

and its renormalized version $\{(\sum_{i=1}^n 4^{|\lambda_i|})^{-\tilde{t}/2} \tilde{\psi}_\lambda^{(\boldsymbol{\sigma}, \vec{\sigma})} : \lambda \in \nabla_{\boldsymbol{\sigma}, \vec{\sigma}}\}$ is a Riesz basis for $H_{\vec{\sigma}}^{\tilde{t}}(\square)$.

For $\lambda \in \nabla_{\sigma, \bar{\sigma}}$, we set $|\lambda| := (|\lambda_1|, \dots, |\lambda_n|)$. As usual, for $\mathbf{j}, \mathbf{j} \in \mathbb{N}_0^n$, $|\mathbf{j}| \leq |\mathbf{j}|$ will mean that $|\mathbf{j}|_i \leq |\mathbf{j}|_i$ ($1 \leq i \leq n$), whereas $|\mathbf{j}| \geq |\mathbf{j}|$ or $|\mathbf{j}| = |\mathbf{j}|$ will mean that $|\mathbf{j}| \leq |\mathbf{j}|$ or $|\mathbf{j}| \leq |\mathbf{j}|$ and $|\mathbf{j}| \geq |\mathbf{j}|$, respectively.

For $\theta \geq 0$, the *weighted Sobolev space* $\mathcal{H}_\theta^d(\mathcal{I})$ is defined as the space of all measurable functions u on \mathcal{I} for which the norm

$$\|u\|_{\mathcal{H}_\theta^d(\mathcal{I})} := \left[\sum_{j=0}^d \int_{\mathcal{I}} |x^\theta (1-x)^\theta u^{(j)}(x)|^2 dx \right]^{\frac{1}{2}}$$

is finite. For

$$m \in \{0, \dots, [t]\},$$

we will consider the *weighted Sobolev space*

$$\mathcal{H}_{m, \theta}^d(\square) := \bigcap_{p=1}^n \otimes_{i=1}^n \mathcal{H}_{\theta - \delta_{ip} \min(m, \theta)}^d(\mathcal{I}),$$

equipped with a squared norm that is the sum over $p = 1, \dots, n$ of the squared norms on $\otimes_{i=1}^n \mathcal{H}_{\theta - \delta_{ip} \min(m, \theta)}^d(\mathcal{I})$.

Theorem 3.2 ([DS10, Thm. 4.3]). *For any $\theta \in [0, d]$, there exist a (nested) sequence $(\nabla_M^{(\sigma, \bar{\sigma})})_{M \in \mathbb{N}} \subset \nabla_{\sigma, \bar{\sigma}}$ with $\#\nabla_M^{(\sigma, \bar{\sigma})} \approx M$, such that*

$$\inf_{v \in \text{span}\{\psi_\lambda^{(\sigma, \bar{\sigma})} : \lambda \in \nabla_M^{(\sigma, \bar{\sigma})}\}} \|u - v\|_{H^m(\square)} \lesssim M^{-(d-m)} \|u\|_{\mathcal{H}_{m, \theta}^d(\square)}, \quad (u \in \mathcal{H}_{m, \theta}^d(\square) \cap H_\sigma^m(\square)),$$

where for $m = 0$, $M^{-(d-m)}$ should be read as $(\log \#M)^{(n-1)(\frac{1}{2}+d)} M^{-d}$.

The index sets $\nabla_M^{(\sigma, \bar{\sigma})}$ can be chosen to have the following multiple tree property: For any $\lambda \in \nabla_M^{(\sigma, \bar{\sigma})}$ and any $\mathbf{j} \in \mathbb{N}_0^n$ with $\mathbf{j} \leq |\lambda|$, there exists a $\mu \in \nabla_M^{(\sigma, \bar{\sigma})}$ with $|\mu| = \mathbf{j}$, and $\text{supp } \psi_\lambda^{(\sigma, \bar{\sigma})} \cap \text{supp } \psi_\mu^{(\sigma, \bar{\sigma})} \neq \emptyset$.

With the notations $u \in H_\sigma^t(\alpha + \square)$ and $u \in \mathcal{H}_{m, \theta}^d(\alpha + \square)$, we will mean that $u(\cdot + \alpha) \in H_\sigma^t(\square)$ or $u(\cdot + \alpha) \in \mathcal{H}_{m, \theta}^d(\square)$, respectively.

4. CONSTRUCTION OF RIESZ BASES BY EXTENSION

Let $\{\square_0, \dots, \square_N\}$ be a set of hypercubes in $\mathbb{Z}^n + \square$, and let $\hat{\Omega}$ be a (reference) domain (i.e., open and connected) in \mathbb{R}^n with $\cup_{k=0}^N \square_k \subset \hat{\Omega} \subset (\cup_{k=0}^N \bar{\square}_k)^{\text{int}}$, and such that $\partial \hat{\Omega}$ is the union of (closed) facets of the \square_k 's. The case $\hat{\Omega} \subsetneq (\cup_{k=0}^N \bar{\square}_k)^{\text{int}}$ corresponds to the situation that $\hat{\Omega}$ has one or more cracks. We will describe a construction of Riesz bases for Sobolev spaces on $\hat{\Omega}$ from Riesz bases for corresponding Sobolev spaces on the subdomains \square_k using extension operators. We start with giving sufficient conditions (\mathcal{D}_1) – (\mathcal{D}_5) such that suitable extension operators exist. At the end of this section, we will consider domains given as the parametric image of $\hat{\Omega}$.

We assume that there exists a sequence $(\{\hat{\Omega}_k^{(q)} : q \leq k \leq N\})_{0 \leq q \leq N}$ of sets of polytopes, such that $\hat{\Omega}_k^{(0)} = \square_k$ and where each next term in the sequence is created from its predecessor by joining two of its polytopes. More precisely, we assume that for any $1 \leq q \leq N$, there exists a $q \leq \bar{k} = \bar{k}^{(q)} \leq N$ and $q - 1 \leq k_1 = k_1^{(q)} \neq k_2 = k_2^{(q)} \leq N$ such that

$$\begin{aligned} (\mathcal{D}_1) \quad \hat{\Omega}_{\bar{k}}^{(q)} &= \left(\hat{\Omega}_{k_1}^{(q-1)} \cup \hat{\Omega}_{k_2}^{(q-1)} \setminus \partial \hat{\Omega} \right)^{\text{int}} \text{ is connected, and the interface } J := \hat{\Omega}_{\bar{k}}^{(q)} \setminus \\ & \left(\hat{\Omega}_{k_1}^{(q-1)} \cup \hat{\Omega}_{k_2}^{(q-1)} \right) \text{ is part of a hyperplane,} \end{aligned}$$

$$\begin{aligned} (\mathcal{D}_2) \quad & \{\hat{\Omega}_k^{(q)} : q \leq k \leq N, k \neq \bar{k}\} = \{\hat{\Omega}_k^{(q-1)} : q-1 \leq k \leq N, k \neq \{k_1, k_2\}\}, \\ (\mathcal{D}_3) \quad & \hat{\Omega}_N^{(N)} = \hat{\Omega}. \end{aligned}$$

For some

$$t \in [0, \infty) \setminus (\mathbb{N}_0 + \{\frac{1}{2}\}),$$

to each of the *closed* facets of all the hypercubes \square_k , we associate a number in $\{0, \dots, \lfloor t + \frac{1}{2} \rfloor\}$ indicating the order of the Dirichlet boundary condition on that facet (where a Dirichlet boundary condition of order 0 means no boundary condition). On facets on the boundary of $\hat{\Omega}$, this number can be chosen at one's convenience (it is dictated by the boundary conditions of the boundary value problem that one wants to solve on $\hat{\Omega}$), and, as will follow from the conditions imposed below, on the other facets it should be either 0 or $\lfloor t + \frac{1}{2} \rfloor$.

By construction, each facet of any $\hat{\Omega}_k^{(q)}$ is a union of some facets of the $\square_{k'}$'s, that will be referred to as subfacets. Letting each of these subfacets inherit the Dirichlet boundary conditions imposed on the $\square_{k'}$'s, we define

$$\mathring{H}^t(\hat{\Omega}_k^{(q)}),$$

and so for $k = q = N$ in particular $\mathring{H}^t(\hat{\Omega}) = \mathring{H}^t(\hat{\Omega}_N^{(N)})$, to be the closure in $H^t(\hat{\Omega}_k^{(q)})$ of the smooth functions on $\hat{\Omega}_k^{(q)}$ that satisfy these boundary conditions. Note that for $0 \leq k \leq N$, for some $\sigma(k) \in (\{0, \dots, \lfloor t + \frac{1}{2} \rfloor\}^2)^n$,

$$\mathring{H}^t(\hat{\Omega}_k^{(0)}) = \mathring{H}^t(\square_k) = H_{\sigma(k)}^t(\square_k).$$

Remark 4.1. On the intersection of facets of hypercubes $\square_{k'}$, the natural interpretation of the boundary conditions is the minimal one such that the boundary conditions on each of these facets is not violated.

The boundary conditions on the hypercubes that determine the spaces $\mathring{H}^t(\hat{\Omega}_k^{(q)})$, and the order in which polytopes are joined should be chosen such that

$$(\mathcal{D}_4) \quad \text{on the } \hat{\Omega}_{k_1}^{(q-1)} \text{ and } \hat{\Omega}_{k_2}^{(q-1)} \text{ sides of } J, \text{ the boundary conditions are of order } 0 \text{ and } \lfloor t + \frac{1}{2} \rfloor, \text{ respectively,}$$

and, w.l.o.g. assuming that $J = \{0\} \times \check{J}$ and $(0, 1) \times \check{J} \subset \Omega_{k_1}^{(q-1)}$,

$$(\mathcal{D}_5) \quad \text{for any function in } \mathring{H}^t(\hat{\Omega}_{k_1}^{(q-1)}) \text{ that vanishes near } \{0, 1\} \times \check{J}, \text{ its reflection in } \{0\} \times \mathbb{R}^{n-1} \text{ (extended with zero, and then restricted to } \hat{\Omega}_{k_2}^{(q-1)}) \text{ is in } \mathring{H}^t(\hat{\Omega}_{k_2}^{(q-1)}).$$

The condition (\mathcal{D}_5) can be formulated by saying that the order of the boundary condition at any subfacet of $\hat{\Omega}_{k_1}^{(q-1)}$ adjacent to J should not be less than this order at its reflection in J , where in case this reflection is not part of $\partial\hat{\Omega}_{k_2}^{(q-1)}$ the latter order should be read as the highest possible one $\lfloor t + \frac{1}{2} \rfloor$; and furthermore, that the order of the boundary condition at any subfacet of $\hat{\Omega}_{k_2}^{(q-1)}$ adjacent to J should not be larger than this order at its reflection in J , where in case this reflection is not part of $\partial\hat{\Omega}_{k_1}^{(q-1)}$, the latter order should be read as the lowest possible one 0. See Figure 3 for an illustration.

Given $1 \leq q \leq N$, for $i \in \{1, 2\}$, let $R_i^{(q)}$ be the *restriction* of functions on $\hat{\Omega}_{\bar{k}}^{(q)}$ to $\hat{\Omega}_{k_i}^{(q-1)}$, let $\eta_2^{(q)}$ be the *extension* of functions on $\hat{\Omega}_{k_2}^{(q-1)}$ to $\hat{\Omega}_{\bar{k}}^{(q)}$ by zero, and let $E_1^{(q)}$ be some *extension* of functions on $\hat{\Omega}_{k_1}^{(q-1)}$ to $\hat{\Omega}_{\bar{k}}^{(q)}$.




FIGURE 3. Two illustrations with (\mathcal{D}_1) – (\mathcal{D}_5) . The fat arrow indicates the action of the extension $E_1^{(q)}$.

Proposition 4.2. *Assume that*

$$E_1^{(q)} \in B(L_2(\hat{\Omega}_{k_1}^{(q-1)}), L_2(\hat{\Omega}_{k_1}^{(q)})), \quad E_1^{(q)} \in B(\hat{H}^t(\hat{\Omega}_{k_1}^{(q-1)}), \hat{H}^t(\hat{\Omega}_{k_1}^{(q)})).$$

Then for $s \in [0, 1]$

$$E^{(q)} := [E_1^{(q)} \quad \eta_2^{(q)}] \in B\left(\prod_{i=1}^2 [L_2(\hat{\Omega}_{k_i}^{(q-1)}), \hat{H}^t(\hat{\Omega}_{k_i}^{(q-1)})]_{s,2}, [L_2(\hat{\Omega}_{k_1}^{(q)}), \hat{H}^t(\hat{\Omega}_{k_1}^{(q)})]_{s,2}\right)$$

is boundedly invertible.

Proof. Taking $V^{(q)} = L_2(\hat{\Omega}_{k_1}^{(q)})$, $V_i^{(q)} = L_2(\hat{\Omega}_{k_i}^{(q-1)})$, $\underline{V}^{(q)} = \hat{H}^t(\hat{\Omega}_{k_1}^{(q)})$, $\underline{V}_i^{(q)} = \hat{H}^t(\hat{\Omega}_{k_i}^{(q-1)})$, and noting that $\mathfrak{S}(\text{Id} - E_1^{(q)} R_1^{(q)}) \subset \{u \in \hat{H}^t(\hat{\Omega}_{k_1}^{(q)}) : u = 0 \text{ on } \hat{\Omega}_{k_1}^{(q-1)}\} = \mathfrak{S}(\eta_2^{(q)}|_{\hat{H}^t(\hat{\Omega}_{k_2}^{(q-1)})})$, the result follows from an application of Proposition 2.5(a). \square

Corollary 4.3. *For E being the composition for $q = 1, \dots, N$ of the mappings $E^{(q)}$ from Proposition 4.2, trivially extended with identity operators in coordinates $k \in \{q-1, \dots, N\} \setminus \{k_1^{(q)}, k_2^{(q)}\}$, it holds that*

$$(4.1) \quad E \in B\left(\prod_{k=0}^n [L_2(\square_k), \hat{H}^t(\square_k)]_{s,2}, [L_2(\hat{\Omega}), \hat{H}^t(\hat{\Omega})]_{s,2}\right).$$

is boundedly invertible.

Under the conditions (\mathcal{D}_1) – (\mathcal{D}_5) , the extensions $E_1^{(q)}$ can be constructed (essentially) as tensor products of *univariate extensions* with identity operators in the other Cartesian directions.

Proposition 4.4. *W.l.o.g. let $J = \{0\} \times \check{J}$ and $(0, 1) \times \check{J} \subset \hat{\Omega}_{k_1}^{(q-1)}$. Let G_1 be an extension operator of functions on $(0, 1)$ to functions on $(-1, 1)$ such that*

$$G_1 \in B(L_2(0, 1), L_2(-1, 1)), \quad G_1 \in B(H^t(0, 1), H^t_{([t+\frac{1}{2}], 0)}(-1, 1)).$$

Then $E_1^{(q)}$ defined by $R_2^{(q)} E_1^{(q)}$ being the composition of the restriction to $(0, 1) \times \check{J}$, followed by an application of

$$G_1 \otimes \text{Id} \otimes \dots \otimes \text{Id},$$

followed by an extension by 0 to $\hat{\Omega}_{k_2}^{(q-1)} \setminus (-1, 0) \times \check{J}$, satisfies the assumptions made in Proposition 4.2.

Remark 4.5. The condition that an extension by G_1 vanishes up to order $\lfloor t + \frac{1}{2} \rfloor$ at -1 is fully harmless since it can easily be enforced by multiplying an extension by some smooth cut-off function. The scale-dependent extension that we will discuss in Subsection 5.1 satisfies this boundary condition automatically.

Our main interest of Corollary 4.3 lies in the following:

Corollary 4.6. *For $0 \leq k \leq N$, let Ψ_k be a Riesz basis for $L_2(\square_k)$, that renormalized in $H^t(\square_k)$, is a Riesz basis for $\hat{H}^t(\square_k) = H^t_{\sigma(k)}(\square)$. Then for $s \in [0, 1]$, and with E from Corollary 4.3, the collection $E(\prod_{k=0}^N \Psi_k)$, normalized in the corresponding norm, is a Riesz basis for $[L_2(\hat{\Omega}), \hat{H}^t(\hat{\Omega})]_{s,2}$.*

Remark 4.7. Although we allow for $t \in (0, \frac{1}{2})$, for these values of t our exposition is not very relevant. Indeed, for those t , a piecewise tensor product basis can simply be constructed as the union of the tensor product bases on the hypercubes.

To find the corresponding dual basis, we follow Section 2. Taking for $q = 1, \dots, N$,

$$H^{(q)} = L_2(\hat{\Omega}_k^{(q)}), \quad H_i^{(q)} = L_2(\hat{\Omega}_{k_i}^{(q)}),$$

and with $\eta_1^{(q)}$ being the extension of functions on $\hat{\Omega}_{k_1}^{(q-1)}$ to $\hat{\Omega}_k^{(q)}$ by zero, Proposition 2.2 shows that

$$(E^{(q)})^{-*} = [\eta_1^{(q)} \quad (\text{Id} - \eta_1^{(q)}(E_1^{(q)})^*)\eta_2^{(q)}].$$

Corollary 4.8. *In the situation of Corollary 4.6, let $\tilde{\Psi}_k$ the Riesz basis for $L_2(\square_k)$ that is dual to Ψ_k . Then $E^{-*}(\prod_{k=0}^N \tilde{\Psi}_k)$ is the Riesz basis for $L_2(\hat{\Omega})$ that is dual to $E(\prod_{k=0}^N \Psi_k)$. The operator E^{-*} is the composition for $q = 1, \dots, N$ of the mappings $(E^{(q)})^{-*}$ trivially extended with identity operators in coordinates $k \in \{q-1, \dots, N\} \setminus \{k_1^{(q)}, k_2^{(q)}\}$.*

Below we give conditions such that $E^{-*}(\prod_{k=0}^N \tilde{\Psi}_k)$, properly scaled, is a Riesz basis for a range of Sobolev spaces with positive smoothness indices, and so, equivalently, $E(\prod_{k=0}^N \Psi_k)$ to be a Riesz basis for the corresponding dual spaces.

For some $\tilde{t} \in [0, \infty) \setminus (\mathbb{N}_0 + \{\frac{1}{2}\})$, to each of the closed facets of all the hypercubes \square_k , we associate a number in $\{0, \dots, \lfloor \tilde{t} + \frac{1}{2} \rfloor\}$ indicating the order of the dual Dirichlet boundary condition on that facet. On facets on the boundary of $\hat{\Omega}$, this number can be chosen arbitrarily, where on the interior facets it is 0 or $\lfloor \tilde{t} + \frac{1}{2} \rfloor$.

We define $\hat{H}^{\tilde{t}}(\hat{\Omega}_k^{(q)})$, and so for $k = q = N$ in particular $\hat{H}^{\tilde{t}}(\hat{\Omega}) = \hat{H}^{\tilde{t}}(\hat{\Omega}_N^{(N)})$, to be the closure in $H^{\tilde{t}}(\hat{\Omega}_k^{(q)})$ of the smooth functions on $\hat{\Omega}_k^{(q)}$ that on any of its facets satisfy the boundary conditions that were imposed on each of its subfacets. Note that with some abuse of notation, even when $\tilde{t} = t$ generally $\hat{H}^{\tilde{t}}(\hat{\Omega}_k^{(q)}) \neq \hat{H}^t(\hat{\Omega}_k^{(q)})$, and that for $0 \leq k \leq N$,

$$\hat{H}^{\tilde{t}}(\hat{\Omega}_k^{(0)}) = \hat{H}^{\tilde{t}}(\square_k) = H^{\tilde{t}}_{\sigma(k)}(\square_k),$$

for some $\tilde{\sigma}(k) \in (\{0, \dots, \lfloor \tilde{t} + \frac{1}{2} \rfloor\}^2)^n$.

We make the following assumptions on the selection of the boundary conditions that determine the dual spaces $\hat{H}^{\tilde{t}}(\hat{\Omega}_k^{(q)})$:

(\mathcal{D}'_4) on the $\hat{\Omega}_{k_1}^{(q-1)}$ and $\hat{\Omega}_{k_2}^{(q-1)}$ sides of J , the boundary conditions are of order $\lfloor \tilde{t} + \frac{1}{2} \rfloor$ and 0, respectively,

and, w.l.o.g. assuming that $J = \{0\} \times \check{J}$ and $(0, 1) \times \check{J} \subset \Omega_{k_1}^{(q-1)}$,

(\mathcal{D}'_5) for any function in $\mathring{H}^{\tilde{t}}(\hat{\Omega}_{k_2}^{(q-1)})$ that vanishes near $\{-1, 0\} \times \check{J}$, its reflection in $\{0\} \times \mathbb{R}^{n-1}$ (extended with zero, and then restricted to $\hat{\Omega}_{k_1}^{(q-1)}$) is in $\mathring{H}^{\tilde{t}}(\hat{\Omega}_{k_1}^{(q-1)})$.

Proposition 4.9. *For $1 \leq q \leq N$, let the extension $E_1^{(q)}$ be of tensor product form as in Proposition 4.4 with $G_1^* \in B(H_{(0, \lfloor \tilde{t} + \frac{1}{2} \rfloor]}^{\tilde{t}}(-1, 1), H_{[\lfloor \tilde{t} + \frac{1}{2} \rfloor, \lfloor \tilde{t} + \frac{1}{2} \rfloor]}^{\tilde{t}}(0, 1))$, and let $\tilde{\Psi}_k$, properly scaled, be a Riesz basis for $\mathring{H}^{\tilde{t}}(\square_k)$. Then for $s \in [0, 1]$, $E^{-*}(\prod_{k=0}^N \tilde{\Psi}_k)$ is, properly scaled, a Riesz basis for $[L_2(\hat{\Omega}), \mathring{H}^{\tilde{t}}(\hat{\Omega})]_{s,2}$.*

Remark 4.10. The boundary conditions imposed on G_1^*u at 1 are fully harmless. The scale-dependent extension G_1 that we will discuss in Subsection 5.1 satisfies these boundary conditions automatically. On the other hand, thinking of $t \geq \tilde{t}$, the boundary conditions at 0 are, when $\tilde{t} > \frac{1}{2}$, the only properties that are not already implied by the conditions on G_1 from Proposition 4.4.

Proof. The conditions (\mathcal{D}'_4) , (\mathcal{D}'_5) imply both that $R_2^{(q)}$ has a right-inverse which is in $B(\mathring{H}^{\tilde{t}}(\Omega_{k_2}^{(q-1)}), \mathring{H}^{\tilde{t}}(\Omega_k^{(q)}))$ and $(E_1^{(q)})^* \in B(\mathring{H}^{\tilde{t}}(\Omega_k^{(q)}), \mathring{H}^{\tilde{t}}(\Omega_{k_1}^{(q-1)}))$, by the assumption on G_1^* . Since $R_2^{(q)} \in B(\mathring{H}^{\tilde{t}}(\Omega_k^{(q)}), \mathring{H}^{\tilde{t}}(\Omega_{k_2}^{(q-1)}))$, $\eta_1^{(q)} \in B(\mathring{H}^{\tilde{t}}(\Omega_{k_1}^{(q-1)}), \mathring{H}^{\tilde{t}}(\Omega_k^{(q)}))$ directly follow from (\mathcal{D}'_4) , an N -fold application of Proposition 2.4 together with the assumption on the bases $\tilde{\Psi}_k$ completes the proof. \square

To end the discussion about the stability of $E(\prod_{k=0}^N \Psi_k)$ in dual norms, we note that for $\tilde{t} < \frac{1}{2}$, which suffices for our application for solving PDEs, the conditions (\mathcal{D}'_4) , (\mathcal{D}'_5) , and those from Proposition 4.9 are void, with the exception of the very mild condition of $\tilde{\Psi}_k$, properly scaled, being a Riesz basis for $\mathring{H}^{\tilde{t}}(\square_k)$.

The construction of Riesz bases on the reference domain $\hat{\Omega}$ extends to more general domains in a standard fashion. Let Ω be the image of $\hat{\Omega}$ under a homeomorphism κ . We define the *pull-back* κ^* by $\kappa^*w = w \circ \kappa$, and so its inverse κ^{-*} , known as the *push-forward*, satisfies $\kappa^{-*}v = v \circ \kappa^{-1}$.

Proposition 4.11. *Let κ^* be boundedly invertible as a mapping both from $L_2(\Omega)$ to $L_2(\hat{\Omega})$ and from $H^t(\Omega)$ to $H^t(\hat{\Omega})$. Setting $\mathring{H}^t(\Omega) := \mathfrak{S}\kappa^{-*}|_{\mathring{H}^t(\hat{\Omega})}$, we have that $\kappa^{-*} \in B([L_2(\hat{\Omega}), \mathring{H}^t(\hat{\Omega})]_{s,2}, [L_2(\Omega), \mathring{H}^t(\Omega)]_{s,2})$ is boundedly invertible ($s \in [0, 1]$). So if Ψ is a Riesz basis for $L_2(\hat{\Omega})$ and, properly scaled, for $\mathring{H}^t(\hat{\Omega})$, then for $s \in [0, 1]$, $\kappa^{-*}\Psi$ is, properly scaled, a Riesz basis for $[L_2(\Omega), \mathring{H}^t(\Omega)]_{s,2}$.*

If $\tilde{\Psi}$ is the collection dual to Ψ , then $|\det D\kappa^{-1}(\cdot)|\kappa^{-}\tilde{\Psi}$ is the collection dual to $\kappa^{-*}\Psi$.*

5. APPROXIMATION BY –PIECEWISE– TENSOR PRODUCT WAVELETS

In the setting of Proposition 4.4, Corollary 4.6 and Proposition 4.9, writing $\square_k = \square + \alpha_k$, where $\alpha_k \in \mathbb{Z}^n$, we select the the primal and dual bases for $L_2(\square_k)$ to be

$$\Psi_{\sigma(k), \bar{\sigma}(k)}(\cdot - \alpha_k), \quad \tilde{\Psi}_{\sigma(k), \bar{\sigma}(k)}(\cdot - \alpha_k)$$

as constructed in Section 3, which, properly scaled, are Riesz bases for $H_{\sigma(k)}^t(\square_k)$ and $H_{\bar{\sigma}(k)}^{\tilde{t}}(\square_k)$, respectively.

In the setting of Proposition 4.11, for $m \in \{0, \dots, \lfloor t \rfloor\}$ and $u \in \mathring{H}^m(\Omega) := [L_2(\Omega), \mathring{H}^t(\Omega)]_{m/t, 2}$, with additionally

$$(5.1) \quad u \in \kappa^{-*} \left(\prod_{k=0}^N \mathcal{H}_{m, \theta}^d(\square_k) \right) := \{v : \Omega \rightarrow \mathbb{R} : v \circ \kappa \in \prod_{k=0}^N \mathcal{H}_{m, \theta}^d(\square_k)\},$$

we study approximation rates from $\kappa^{-*} E \left(\prod_{k=0}^N \Psi_{\sigma^{(k)}, \vec{\sigma}^{(k)}}(\cdot - \alpha_k) \right)$ in the $H^m(\Omega)$ -norm. Since, as is assumed in Proposition 4.11, $\kappa^* \in B(H^m(\Omega), H^m(\hat{\Omega}))$ is boundedly invertible, it is sufficient to study this issue for the case that $\kappa = \text{Id}$ and so $\Omega = \hat{\Omega}$.

We will apply extension operators $E_1^{(q)}$ that are built from univariate extension operators. The latter will be chosen such that the resulting primal and dual wavelets on $\hat{\Omega}$ are, restricted to each $\square_k \subset \hat{\Omega}$, tensor products of collections of univariate functions that are local and locally finite (cf. parts (1) and (2) of the forthcoming Proposition 5.4).

5.1. Construction of scale-dependent extension operators. We make the following additional assumptions on the univariate wavelets. For $\vec{\sigma} = (\sigma_\ell, \sigma_r) \in \{0, \dots, \lfloor t + \frac{1}{2} \rfloor\}^2$, $\vec{\sigma} = (\tilde{\sigma}_\ell, \tilde{\sigma}_r) \in \{0, \dots, \lfloor \tilde{t} + \frac{1}{2} \rfloor\}^2$, and with $\vec{0} := (0, 0)$,

$$(\mathcal{W}_7) \quad V_j^{(\vec{\sigma})} := \text{span}\{\psi_\lambda^{(\vec{\sigma}, \vec{\sigma})} : \lambda \in \nabla_{\vec{\sigma}, \vec{\sigma}}, |\lambda| \leq j\} \text{ is independent of } \vec{\sigma}, \text{ and } V_j^{(\vec{\sigma})} = V_j^{(\vec{0})} \cap H_{\vec{\sigma}}^t(\mathcal{I}),$$

$$(\mathcal{W}_8) \quad \nabla_{\vec{\sigma}, \vec{\sigma}} \text{ is the disjoint union of } \nabla_{\sigma_\ell, \tilde{\sigma}_\ell}^{(\ell)}, \nabla^{(I)}, \nabla_{\sigma_r, \tilde{\sigma}_r}^{(r)} \text{ such that}$$

$$(i) \quad \sup_{\lambda \in \nabla_{\vec{\sigma}, \vec{\sigma}}^{(\ell)}, x \in \text{supp } \psi_\lambda^{(\vec{\sigma}, \vec{\sigma})}} 2^{|\lambda|} |x| \lesssim \rho, \quad \sup_{\lambda \in \nabla_{\vec{\sigma}, \vec{\sigma}}^{(r)}, x \in \text{supp } \psi_\lambda^{(\vec{\sigma}, \vec{\sigma})}} 2^{|\lambda|} |1-x| \lesssim \rho,$$

$$(ii) \quad \text{for } \lambda \in \nabla^{(I)}, \psi_\lambda^{(\vec{\sigma}, \vec{\sigma})} = \psi_\lambda^{(\vec{0}, \vec{0})}, \tilde{\psi}_\lambda^{(\vec{\sigma}, \vec{\sigma})} = \tilde{\psi}_\lambda^{(\vec{0}, \vec{0})}, \text{ and the extensions of } \psi_\lambda^{(\vec{0}, \vec{0})} \text{ and } \tilde{\psi}_\lambda^{(\vec{0}, \vec{0})} \text{ by zero are in } H^t(\mathbb{R}) \text{ and } H^{\tilde{t}}(\mathbb{R}), \text{ respectively.}$$

$$(\mathcal{W}_9) \quad \begin{cases} \text{span}\{\psi_\lambda^{(\vec{0}, \vec{0})}(1-\cdot) : \lambda \in \nabla^{(I)}, |\lambda| = j\} = \text{span}\{\psi_\lambda^{(\vec{0}, \vec{0})} : \lambda \in \nabla^{(I)}, |\lambda| = j\}, \\ \text{span}\{\psi_\lambda^{(\sigma_\ell, \sigma_r), (\tilde{\sigma}_\ell, \tilde{\sigma}_r)}(1-\cdot) : \lambda \in \nabla_{\sigma_\ell, \tilde{\sigma}_\ell}^{(\ell)}, |\lambda| = j\} = \\ \quad \text{span}\{\psi_\lambda^{(\sigma_r, \sigma_\ell), (\tilde{\sigma}_r, \tilde{\sigma}_\ell)} : \lambda \in \nabla_{\sigma_r, \tilde{\sigma}_r}^{(r)}, |\lambda| = j\}, \end{cases}$$

$$(\mathcal{W}_{10}) \quad \begin{cases} \psi_\lambda^{(\vec{\sigma}, \vec{\sigma})}(2^l \cdot) \in \text{span}\{\psi_\mu^{(\vec{\sigma}, \vec{\sigma})} : \mu \in \nabla_{\sigma_\ell, \tilde{\sigma}_\ell}^{(\ell)}\} \quad (l \in \mathbb{N}_0, \lambda \in \nabla_{\sigma_\ell, \tilde{\sigma}_\ell}^{(\ell)}), \\ \psi_\lambda^{(\vec{0}, \vec{0})}(2^l \cdot) \in \text{span}\{\psi_\mu^{(\vec{0}, \vec{0})} : \mu \in \nabla^{(I)}\} \quad (l \in \mathbb{N}_0, \lambda \in \nabla^{(I)}). \end{cases}$$

As (\mathcal{W}_1) – (\mathcal{W}_6) , these conditions are satisfied by following the biorthogonal wavelet constructions on the interval from [Pri10, Dij09] ((\mathcal{W}_7) is not satisfied by the construction from [DKU99], but the following exposition can be adapted to apply to these wavelets as well).

Remark 5.1. In view of the boundary conditions that are imposed on the interfacets, see (\mathcal{D}_4) and (\mathcal{D}'_4) , it is actually sufficient to impose (\mathcal{W}_7) – (\mathcal{W}_{10}) for $(\sigma_\ell, \tilde{\sigma}_\ell), (\sigma_r, \tilde{\sigma}_r) \in \{(\lfloor t + \frac{1}{2} \rfloor, 0), (0, \lfloor \tilde{t} + \frac{1}{2} \rfloor)\}$.

We consider the setting of Proposition 4.4. W.l.o.g. we assume that $J = \{0\} \times \check{J}$, and $(0, 1) \times \check{J} \subset \hat{\Omega}_{k_1}^{(q-1)}$. We assume to have available a univariate extension operator

$$(5.2) \quad \check{G}_1 \in B(L_2(0, 1), L_2(-1, 1)) \text{ with } \begin{cases} \check{G}_1 \in B(H^t(0, 1), H^t(-1, 1)), \\ \check{G}_1^* \in B(H^{\tilde{t}}(-1, 1), H_{(\lfloor \tilde{t} + \frac{1}{2} \rfloor, 0)}^{\tilde{t}}(0, 1)). \end{cases}$$

Let η_1 and η_2 denote the extensions by zero of functions on $(0, 1)$ or on $(-1, 0)$ to functions on $(-1, 1)$, with R_1 and R_2 denoting their adjoints. We assume that \check{G}_1 and its ‘‘adjoint extension’’

$$\check{G}_2 := (\text{Id} - \eta_1 \check{G}_1^*) \eta_2$$

(cf. Remark 2.3) are local in the sense that

$$(5.3) \quad \begin{cases} \text{diam}(\text{supp } R_2 \check{G}_1 u) \lesssim \text{diam}(\text{supp } u) & (u \in L_2(0, 1)), \\ \text{diam}(\text{supp } R_1 \check{G}_2 u) \lesssim \text{diam}(\text{supp } u) & (u \in L_2(-1, 0)), \end{cases}$$

see Figure 4 for an illustration.




FIGURE 4. Univariate extensions and restrictions.

Examples are given by Hestenes extensions ([Hes41, DS99, KS06]), which are of the form

$$(5.4) \quad \check{G}_1 v(-x) = \sum_{l=0}^L \gamma_l (\zeta v)(\beta_l x) \quad (v \in L_2(\mathcal{I}), x \in \mathcal{I}),$$

(and, being an extension, $\check{G}_1 v(x) = v(x)$ for $x \in \mathcal{I}$), where $\gamma_l \in \mathbb{R}$, $\beta_l > 0$, and $\zeta : [0, \infty) \rightarrow [0, \infty)$ is some smooth cut-off function with $\zeta \equiv 1$ in a neighborhood of 0, and $\text{supp } \zeta \subset [0, \min_l(\beta_l, \beta_l^{-1})]$. Its adjoint reads as

$$\check{G}_1^* w(x) = w(x) + \zeta(x) \sum_{l=0}^L \frac{\gamma_l}{\beta_l} w\left(\frac{-x}{\beta_l}\right) \quad (w \in L_2(-1, 1), x \in \mathcal{I}).$$

A Hestenes extension satisfies (5.2) if and only if

$$\sum_{l=0}^L \gamma_l \beta_l^i = (-1)^i (\mathbb{N}_0 \ni i \leq \lfloor t - \frac{1}{2} \rfloor), \quad \sum_{l=0}^L \gamma_l \beta_l^{-(j+1)} = (-1)^{j+1} (\mathbb{N}_0 \ni j \leq \lfloor \tilde{t} - \frac{1}{2} \rfloor).$$

With a univariate extension \check{G}_1 as in (5.2) at hand, the obvious approach is to define $E_1^{(q)}$ according to Proposition 4.4 with $G_1 = \check{G}_1$. A problem with the choice $G_1 = \check{G}_1$ is that generally (5.3) does *not* imply the desirable property that $\text{diam}(\text{supp } G_1 u) \lesssim \text{diam}(\text{supp } u)$. Indeed, think of the application of a Hestenes extension to a u with a small support that is not located near the interface.

To solve this and the corresponding problem for the adjoint extension, in any case for u being any primal or dual wavelet, respectively, following [DS99] we will apply our construction using the modified, *scale-dependent* univariate extension operator

$$(5.5) \quad G_1 : u \mapsto \sum_{\lambda \in \nabla_{0,0}^{(\ell)}} \langle u, \tilde{\psi}_\lambda^{(\vec{0}, \vec{0})} \rangle_{L_2(\mathcal{I})} \check{G}_1 \psi_\lambda^{(\vec{0}, \vec{0})} + \sum_{\lambda \in \nabla^{(\ell)} \cup \nabla_{0,0}^{(r)}} \langle u, \tilde{\psi}_\lambda^{(\vec{0}, \vec{0})} \rangle_{L_2(\mathcal{I})} \eta_1 \psi_\lambda^{(\vec{0}, \vec{0})}.$$

Taking \check{G}_1 to be a Hestenes extension, under the condition of ρ being sufficiently small, its first advantage is that its application in (5.5) does not “see” the cut-off function ζ , which prevents potential quadrature problems.

Proposition 5.2. *Assuming ρ to be sufficiently small, the scale-dependent extension G_1 from (5.5) satisfies, for $\vec{\sigma} \in \{0, \dots, [t + \frac{1}{2}]\}^2$, $\vec{\sigma}_\ell \in \{0, \dots, [\tilde{t} + \frac{1}{2}]\}^2$*

$$(5.6) \quad G_1 \psi_\mu^{(\vec{\sigma}, \vec{\sigma})} = \begin{cases} \eta_1 \psi_\mu^{(\vec{\sigma}, \vec{\sigma})} & \text{when } \mu \in \nabla^{(I)} \cup \nabla_{\sigma_r, \tilde{\sigma}_r}^{(r)}, \\ \check{G}_1 \psi_\mu^{(\vec{\sigma}, \vec{\sigma})} & \text{when } \mu \in \nabla_{\sigma_\ell, \tilde{\sigma}_\ell}^{(\ell)}. \end{cases}$$

Assuming, additionally, \check{G}_1 to be a Hestenes extension with $\beta_l = 2^l$, the resulting adjoint extension $G_2 := (\text{Id} - \eta_1 G_1^*) \eta_2$ satisfies

$$(5.7) \quad G_2(\tilde{\psi}_\mu^{(\vec{\sigma}, \vec{\sigma})}(1 + \cdot)) = \begin{cases} \eta_2(\tilde{\psi}_\mu^{(\vec{\sigma}, \vec{\sigma})}(1 + \cdot)) & \text{when } \mu \in \nabla^{(I)} \cup \nabla_{\sigma_\ell, \tilde{\sigma}_\ell}^{(\ell)}, \\ \check{G}_2(\tilde{\psi}_\mu^{(\vec{\sigma}, \vec{\sigma})}(1 + \cdot)) & \text{when } \mu \in \nabla_{\sigma_r, \tilde{\sigma}_r}^{(r)}. \end{cases}$$

We have $G_1 \in B(L_2(0, 1), L_2(-1, 1))$, $G_1 \in B(H^t(0, 1), H^t(-1, 1))$, and $G_1^* \in B(H^{\tilde{t}}(-1, 1), H_{([\tilde{t} + \frac{1}{2}], 0)}^{\tilde{t}}(0, 1))$.

Finally, for $\mu \in \nabla_{\vec{\sigma}, \vec{\sigma}}$, it holds that

$$(5.8) \quad \begin{aligned} \text{diam}(\text{supp } \check{G}_1 \psi_\mu^{(\vec{\sigma}, \vec{\sigma})}) &\lesssim \text{diam}(\text{supp } \psi_\mu^{(\vec{\sigma}, \vec{\sigma})}), \\ \text{diam}(\text{supp } \check{G}_2 \tilde{\psi}_\mu^{(\vec{\sigma}, \vec{\sigma})}) &\lesssim \text{diam}(\text{supp } \tilde{\psi}_\mu^{(\vec{\sigma}, \vec{\sigma})}). \end{aligned}$$

Proof. By (\mathcal{W}_8) (ii), for $\mu \in \nabla^{(I)} \cup \nabla_{\sigma_r, \tilde{\sigma}_r}^{(r)}$, $\lambda \in \nabla_{0,0}^{(\ell)}$, one has $\langle \psi_\mu^{(\vec{\sigma}, \vec{\sigma})}, \tilde{\psi}_\lambda^{(\vec{0}, \vec{0})} \rangle_{L_2(\mathcal{I})} = 0$, and so $G_1 \psi_\mu^{(\vec{\sigma}, \vec{\sigma})} = \sum_{\lambda \in \nabla_{\vec{0}, \vec{0}}} \langle \psi_\mu^{(\vec{\sigma}, \vec{\sigma})}, \tilde{\psi}_\lambda^{(\vec{0}, \vec{0})} \rangle_{L_2(\mathcal{I})} \eta_1 \psi_\lambda^{(\vec{0}, \vec{0})} = \eta_1 \psi_\mu^{(\vec{\sigma}, \vec{\sigma})}$, the last equality from $\Psi^{(\vec{0}, \vec{0})}$ being a Riesz basis for $L_2(\mathcal{I})$, and η_1 being L_2 -bounded.

Similarly, for $\mu \in \nabla_{\sigma_\ell, \tilde{\sigma}_\ell}^{(\ell)}$, $\lambda \in \nabla^{(I)} \cup \nabla_{0,0}^{(r)}$, it holds that $\langle \psi_\mu^{(\vec{\sigma}, \vec{\sigma})}, \tilde{\psi}_\lambda^{(\vec{0}, \vec{0})} \rangle_{L_2(\mathcal{I})} = 0$, and so $G_1 \psi_\mu^{(\vec{\sigma}, \vec{\sigma})} = \sum_{\lambda \in \nabla_{(\vec{0}, \vec{0})}} \langle \psi_\mu^{(\vec{\sigma}, \vec{\sigma})}, \tilde{\psi}_\lambda^{(\vec{0}, \vec{0})} \rangle_{L_2(\mathcal{I})} \check{G}_1 \psi_\lambda^{(\vec{0}, \vec{0})} = \check{G}_1 \psi_\mu^{(\vec{\sigma}, \vec{\sigma})}$.

If \check{G}_1 is a Hestenes extension with $\beta_l = 2^l$, then for $v \in L_2(\mathcal{I})$,

$$(5.9) \quad \begin{aligned} G_1^* \eta_2(v(1 + \cdot)) &= \sum_{\lambda \in \nabla_{\vec{0}, \vec{0}}} \langle G_1^* \eta_2(v(1 + \cdot)), \psi_\lambda^{(\vec{0}, \vec{0})} \rangle_{L_2(\mathcal{I})} \tilde{\psi}_\lambda^{(\vec{0}, \vec{0})} \\ &= \sum_{\lambda \in \nabla_{\vec{0}, \vec{0}}} \langle v(1 - \cdot), (R_2 G_1 \psi_\lambda^{(\vec{0}, \vec{0})})(-\cdot) \rangle_{L_2(\mathcal{I})} \tilde{\psi}_\lambda^{(\vec{0}, \vec{0})} \\ &= \sum_{\lambda \in \nabla_{0,0}^{(\ell)}} \langle v(1 - \cdot), (R_2 \check{G}_1 \psi_\lambda^{(\vec{0}, \vec{0})})(-\cdot) \rangle_{L_2(\mathcal{I})} \tilde{\psi}_\lambda^{(\vec{0}, \vec{0})} \\ &= \sum_{\lambda \in \nabla_{0,0}^{(\ell)}} \left\langle v(1 - \cdot), \sum_{l=0}^L \gamma_l \psi_\lambda^{(\vec{0}, \vec{0})}(2^l \cdot) \right\rangle_{L_2(\mathcal{I})} \tilde{\psi}_\lambda^{(\vec{0}, \vec{0})}. \end{aligned}$$

For $v = \tilde{\psi}_\mu^{(\vec{\sigma}, \vec{\sigma})}$ and $\mu \in \nabla^{(I)} \cup \nabla_{\sigma_\ell, \tilde{\sigma}_\ell}^{(\ell)}$, (5.9) is zero by (\mathcal{W}_9) , (\mathcal{W}_{10}) , and (\mathcal{W}_8) (ii). For $v = \tilde{\psi}_\mu^{(\vec{\sigma}, \vec{\sigma})}$ and $\mu \in \nabla_{\sigma_r, \tilde{\sigma}_r}^{(r)}$, one has $\left\langle v(1 - \cdot), \sum_{l=0}^L \gamma_l (\zeta \psi_\lambda^{(\vec{0}, \vec{0})})(2^l \cdot) \right\rangle_{L_2(\mathcal{I})} = 0$ for $\lambda \in \nabla^{(I)} \cup \nabla_{0,0}^{(r)}$ by (\mathcal{W}_9) , (\mathcal{W}_{10}) , and (\mathcal{W}_8) (ii). So for those μ , one has $G_1^* \eta_2 \tilde{\psi}_\mu^{(\vec{\sigma}, \vec{\sigma})} = \check{G}_1^* \eta_2 \tilde{\psi}_\mu^{(\vec{\sigma}, \vec{\sigma})}$, which completes the proof of (5.7).

Since $\text{span}\{\psi_\mu^{(\vec{0},\vec{0})} : \mu \in \nabla^{(I)} \cup \nabla_{0,0}^{(r)}\} + \text{span}\{\psi_\mu^{(\vec{0},\vec{0})} : \mu \in \nabla_{0,0}^{(\ell)}\}$ defines a stable splitting of both $L_2(\mathcal{I})$ and $H^t(\mathcal{I})$ into two subspaces, the statements about the boundedness of G_1 follow from (5.6) with $(\vec{\sigma}, \vec{\sigma}) = (\vec{0}, \vec{0})$, (5.2), and (\mathcal{W}_8) (iii).

The mapping $P : u \mapsto \sum_{\mu \in \nabla^{(I)} \cup \nabla_{\sigma_\ell, \tilde{\sigma}_\ell}^{(\ell)}} \langle u, \psi_\mu^{(\vec{\sigma}, \vec{\sigma})}(1 + \cdot) \rangle_{L_2(-1,0)} \eta_2(\tilde{\psi}_\mu^{(\vec{\sigma}, \vec{\sigma})}(1 + \cdot))$ is in $B(H^{\tilde{t}}(-1, 1), H^{\tilde{t}}(-1, 1))$ by the assumption on $\tilde{\Psi}^{(\vec{\sigma}, \vec{\sigma})}$ and (\mathcal{W}_8) (ii). Since $\Psi^{(\vec{\sigma}, \vec{\sigma})}(1 + \cdot)$ is a Riesz basis for $L_2(-1, 0)$,

$$R_2(I - P)u = \sum_{\mu \in \nabla_{\sigma_r, \tilde{\sigma}_r}^{(r)}} \langle u, \psi_\mu^{(\vec{\sigma}, \vec{\sigma})}(1 + \cdot) \rangle_{L_2(-1,0)} \tilde{\psi}_\mu^{(\vec{\sigma}, \vec{\sigma})}(1 + \cdot).$$

We conclude that $(G_1^* - \check{G}_1^*)\eta_2 R_2(I - P) = 0$. Since G_1 and \check{G}_1 are extensions, we also have $G_1^* \eta_1 = \text{Id} = \check{G}_1 - 1^* \eta_1$, and so $G_1^*(I - P) = G_1^*(\eta_1 R_1 + \eta_2 R_2)(I - P) = \check{G}_1^*(I - P)$. Together with $G_1^* P = 0$, from (5.2) we conclude that $G_1^* \in B(H^{\tilde{t}}(-1, 1), H^{\tilde{t}}(\lfloor \tilde{t} + \frac{1}{2} \rfloor, 0)(0, 1))$.

The last statement is a direct consequence of (5.6) and (5.7). \square

Remark 5.3. Although implicitly claimed otherwise in [DS99, (4.3.12)], we note that (5.7), and so (5.8), *cannot* be expected for \check{G}_1 being a general Hestenes extension as given by (5.4), so without assuming that $\beta_l = 2^l$.

Moreover, (5.7), and so (5.8), are only guaranteed when, for $(\sigma_\ell, \tilde{\sigma}_\ell) = (0, \lfloor \tilde{t} + \frac{1}{2} \rfloor)$, for any $\lambda \in \nabla_{\vec{\sigma}, \vec{\sigma}}$ for which either $\psi_\lambda^{(\vec{\sigma}, \vec{\sigma})}$ or $\tilde{\psi}_\lambda^{(\vec{\sigma}, \vec{\sigma})}$ “depends on” the boundary conditions imposed at the left boundary, the primal wavelet $\psi_\lambda^{(\vec{\sigma}, \vec{\sigma})}$ is extended by the application of \check{G}_1 . The reason to emphasize this is that with common biorthogonal wavelet constructions on the interval, the number of dual wavelets that depend on the boundary conditions is larger than that of the primal ones. Note that even if dual wavelets may not enter the computations, their locality as given by (5.8) will be used to prove the forthcoming Theorem 5.6 about the approximation rates provided by the primal piecewise tensor product wavelets.

Some examples of relevant Hestenes extensions with $\beta_l = 2^l$ are:

- $L = 0$, $\gamma_0 = 1$ (reflection). Satisfies (5.2) for $t < \frac{3}{2}$, $\tilde{t} < \frac{1}{2}$,
- $L = 1$, $\gamma_0 = 3$, $\gamma_1 = -2$. Satisfies (5.2) for $t < \frac{5}{2}$, $\tilde{t} < \frac{1}{2}$,
- $L = 1$, $\gamma_0 = -3$, $\gamma_1 = 4$. Satisfies (5.2) for $t < \frac{3}{2}$, $\tilde{t} < \frac{3}{2}$,
- $L = 2$, $\gamma_0 = -5$, $\gamma_1 = 10$, $\gamma_2 = -4$. Satisfies (5.2) for $t < \frac{5}{2}$, $\tilde{t} < \frac{3}{2}$.

In order to identify individual wavelets from the collections constructed by the applications of the extension operators, we have to introduce some more notations. For $0 \leq q \leq N$, we set the index sets

$$\begin{aligned} \nabla_k^{(0)} &:= \nabla_{\sigma^{(k)}, \tilde{\sigma}^{(k)}} \times \{k\} \text{ and, for } q > 0, \\ \nabla_k^{(q)} &:= \begin{cases} \nabla_{k_1}^{(q-1)} \cup \nabla_{k_2}^{(q-1)} & \text{if } k = \bar{k}, \\ \nabla_{\bar{k}}^{(q-1)} & \text{if } k \in \{q, \dots, N\} \setminus \{\bar{k}\} \text{ and } \Omega_k^{(q)} = \Omega_{\bar{k}}^{(q-1)}, \end{cases} \end{aligned}$$

and, for $(\lambda, p) \in \nabla_k^{(q)}$, the primal and dual wavelets,

$$\psi_{\lambda,p}^{(0,k)} := \psi_\lambda^{(\sigma^{(p)}, \tilde{\sigma}^{(p)})}(\cdot - \alpha_p), \quad \tilde{\psi}_{\lambda,p}^{(0,k)} := \tilde{\psi}_\lambda^{(\sigma^{(p)}, \tilde{\sigma}^{(p)})}(\cdot - \alpha_p),$$

and, for $q > 0$,

$$\begin{aligned} \psi_{\lambda,p}^{(q,k)} &:= \begin{cases} \left\{ \begin{array}{l} E_1^{(q)} \psi_{\lambda,p}^{(q-1,k_1)} \\ \eta_2^{(q)} \psi_{\lambda,p}^{(q-1,k_2)} \end{array} \right. & (\lambda, p) \in \nabla_{k_1}^{(q-1)} \\ & (\lambda, p) \in \nabla_{k_2}^{(q-1)} \end{cases} \quad \text{if } k = \bar{k}, \\ & \psi_{\lambda,p}^{(q-1,\hat{k})} & \text{if } k \in \{q, \dots, N\} \setminus \{\bar{k}\} \text{ and } \Omega_k^{(q)} = \Omega_{\hat{k}}^{(q-1)}, \end{cases} \\ \tilde{\psi}_{\lambda,p}^{(q,k)} &:= \begin{cases} \left\{ \begin{array}{l} \eta_1^{(q)} \tilde{\psi}_{\lambda,p}^{(q-1,k_1)} \\ (\text{Id} - \eta_1^{(q)}(E_1^{(q)})^*) \eta_2^{(q)} \tilde{\psi}_{\lambda,p}^{(q-1,k_2)} \end{array} \right. & (\lambda, p) \in \nabla_{k_1}^{(q-1)} \\ & (\lambda, p) \in \nabla_{k_2}^{(q-1)} \end{cases} \quad \text{if } k = \bar{k}, \\ & \tilde{\psi}_{\lambda,p}^{(q-1,\hat{k})} & \text{if } k \in \{q, \dots, N\} \setminus \{\bar{k}\} \text{ and } \Omega_k^{(q)} = \Omega_{\hat{k}}^{(q-1)}, \end{cases} \end{aligned}$$

Then, as we have seen,

$$(\Psi_k^{(q)}, \tilde{\Psi}_k^{(q)}) := (\{\psi_{\lambda,p}^{(q,k)} : (\lambda, p) \in \nabla_k^{(q)}\}, \{\tilde{\psi}_{\lambda,p}^{(q,k)} : (\lambda, p) \in \nabla_k^{(q)}\})$$

is a pair of biorthogonal Riesz bases for $L_2(\hat{\Omega}_k^{(q)})$, and for $s \in [0, 1]$, $\Psi_k^{(q)}$ or $\tilde{\Psi}_k^{(q)}$ are, properly scaled, Riesz bases for $[L_2(\hat{\Omega}_k^{(q)}), \mathring{H}^t(\hat{\Omega}_k^{(q)})]_{s,2}$ and $[L_2(\hat{\Omega}_k^{(q)}), \mathring{H}^{\tilde{t}}(\hat{\Omega}_k^{(q)})]_{s,2}$, respectively.

Proposition 5.4. *With $E_1^{(q)}$ being defined using the scale-dependent extension operator as in Proposition 5.2, for $0 \leq q \leq k \leq N$, we have*

- (1) $\text{supp } \psi_{\lambda,p}^{(q,k)}$, $\text{supp } \tilde{\psi}_{\lambda,p}^{(q,k)}$ are contained in a hyperrectangle aligned with the Cartesian coordinates with sides in length of order $2^{-|\lambda|_1}, \dots, 2^{-|\lambda|_n}$,
- (2) for any $y \in \mathbb{R}^n$ and $\mathbf{j} \in \mathbb{N}_0^n$, the hyperrectangle $y + \prod_{i=1}^n [0, 2^{-j_i}]$ is intersected by the supports at most a uniformly bounded number of primal or dual wavelets $\psi_{\lambda,p}^{(q,k)}$, $\tilde{\psi}_{\lambda,p}^{(q,k)}$ with $|\lambda| = \mathbf{j}$,
- (3) let

$$V_{\mathbf{j}}(\hat{\Omega}_k^{(q)}) := \{u \in \mathring{H}^t(\hat{\Omega}_k^{(q)}) : u(\cdot + \alpha_{k'})|_{\square} \in \otimes_{i=1}^n V_{j_i}^{(\bar{0})} (\square_{k'} \subset \hat{\Omega}_k^{(q)})\}$$

$$Z_{\mathbf{j}}(\hat{\Omega}_k^{(q)}) := \text{span}\{\psi_{\lambda,p}^{(q,k)} : (\lambda, p) \in \nabla_k^{(q)}, |\lambda| \leq \mathbf{j}\},$$

and $\mathbf{e} := (1, \dots, 1)^\top \in \mathbb{R}^n$. Then for some constants $m_q, M_q \in \mathbb{N}_0$, for all $\mathbf{j} \in \{m_q, m_q + 1, \dots\}^n$,

$$V_{\mathbf{j}-m_q\mathbf{e}}(\hat{\Omega}_k^{(q)}) \subset Z_{\mathbf{j}}(\hat{\Omega}_k^{(q)}) \subset V_{\mathbf{j}+M_q\mathbf{e}}(\hat{\Omega}_k^{(q)}).$$

Proof. Parts (1) and (2) follow from the locality and the locally finiteness of the univariate primal and dual wavelets (\mathcal{W}_5) and (\mathcal{W}_6), and the locality of the extension and the adjoint extension given by (5.6) and (5.7).

By construction of the wavelet basis, the second inclusion in (3) follows from (5.6) and \check{G}_1 being a Hestenes extension with $\beta_l = 2^l$, (\mathcal{W}_{10}), (\mathcal{W}_9), and (\mathcal{W}_7). The constant M_q can be taken to be less than or equal to $2L$, or to L when the domain has no cracks.

The first inclusion in (3) holds true for $q = 0$ with $m_0 = 0$ by (\mathcal{W}_7). Suppose, for some m_{q-1} , it is true for $q-1$ and $q-1 \leq k \leq N$. For some constant $m_q \geq m_{q-1}$ that will be determined below, let $v \in V_{\mathbf{j}-m_q\mathbf{e}}(\hat{\Omega}_k^{(q)})$. Then $R_1^{(q)}v \in V_{\mathbf{j}-m_q\mathbf{e}}(\hat{\Omega}_{k_1}^{(q-1)}) \subset Z_{\mathbf{j}+(m_{q-1}-m_q)\mathbf{e}}(\hat{\Omega}_{k_1}^{(q-1)})$, and so

$$(5.10) \quad E_1^{(q)}R_1^{(q)}v \in Z_{\mathbf{j}+(m_{q-1}-m_q)\mathbf{e}}(\hat{\Omega}_k^{(q)}) \subset Z_{\mathbf{j}}(\hat{\Omega}_k^{(q)})$$

by definition of $\Psi_k^{(q)}$.

From (5.10), we have $E_1^{(q)} R_1^{(q)} v \in V_{\mathbf{j}+(m_{q-1}+M_q-m_q)\mathbf{e}}(\hat{\Omega}_{\bar{k}}^{(q)})$, and so $(I-E_1^{(q)} R_1^{(q)})v \in V_{\mathbf{j}+(m_{q-1}+M_q-m_q)\mathbf{e}}(\hat{\Omega}_{\bar{k}}^{(q)})$, and therefore

$$R_2^{(q)}(\text{Id} - E_1^{(q)} R_1^{(q)})v \in \{u \in L_2(\hat{\Omega}_{k_2}^{(q-1)}) : u(\cdot + \alpha_{k'})|_{\square} \in \otimes_{i=1}^n V_{\mathbf{j}_i+m_{q-1}+M_q-m_q}(\bar{\mathbf{0}}) (\square_{k'} \subset \hat{\Omega}_{k_2}^{(q-1)})\}$$

Since, as shown in the proof of Proposition 4.2, $(\text{Id} - E_1^{(q)} R_1^{(q)})v \in \mathfrak{S}(\eta_2^{(q)}|_{\hat{H}^t(\hat{\Omega}_{k_2}^{(q-1)})})$, and so $R_2^{(q)}(\text{Id} - E_1^{(q)} R_1^{(q)})v \in \hat{H}^t(\hat{\Omega}_{k_2}^{(q-1)})$, we infer that $R_2^{(q)}(\text{Id} - E_1^{(q)} R_1^{(q)})v \in V_{\mathbf{j}+(m_{q-1}+M_q-m_q)\mathbf{e}}(\hat{\Omega}_{k_2}^{(q-1)}) \subset Z_{\mathbf{j}+(2m_{q-1}+M_q-m_q)\mathbf{e}}(\hat{\Omega}_{k_2}^{(q-1)})$. By taking $m_q = 2m_{q-1} + M_q$, we conclude that $(\text{Id} - E_1^{(q)} R_1^{(q)})v = \eta_2^{(q)} R_2^{(q)}(\text{Id} - E_1^{(q)} R_1^{(q)})v \in Z_{\mathbf{j}}(\hat{\Omega}_{\bar{k}}^{(q)})$ by definition of $\Psi_{\bar{k}}^{(q)}$. Together with (5.10), this completes the proof. \square

Remark 5.5. The above proof shows that for $L = 0$ (reflection), $V_{\mathbf{j}}(\hat{\Omega}_{\bar{k}}^{(q)}) = Z_{\mathbf{j}}(\hat{\Omega}_{\bar{k}}^{(q)})$.

Now we are ready to study the question, raised at the beginning of this section, about the rate of approximation in $\hat{H}^m(\hat{\Omega})$ from the span of $\Psi := \Psi_N^{(N)}$.

Theorem 5.6. *Let the $E_1^{(q)}$ be defined using the scale-dependent extension operators as in Proposition 5.2. Then for any $\theta \in [0, d)$, and any $0 \leq q \leq k \leq N$, there exists a (nested) sequence $(\nabla_{k,M}^{(q)})_{M \in \mathbb{N}} \subset \nabla_k^{(q)}$ with $\#\nabla_{k,M}^{(q)} \approx M$, such that*

$$(5.11) \quad \inf_{v \in \text{span}\{\psi_{\lambda,p}^{(q,k)} : (\lambda,p) \in \nabla_{k,M}^{(q)}\}} \|u - v\|_{H^m(\hat{\Omega}_k^{(q)})} \lesssim M^{-(d-m)} \sqrt{\sum_{\square_{k'} \subset \hat{\Omega}_k^{(q)}} \|u\|_{\mathcal{H}_{m,\theta}^d(\square_{k'})}^2}$$

for any $u \in \hat{H}^m(\hat{\Omega}_k^{(q)})$ for which the right-hand side is finite (for $q = k = N$, i.e., for $\hat{\Omega}_k^{(q)} = \hat{\Omega}$, this is equivalent to saying that u satisfies (5.1) (with $\kappa = \text{Id}$)).

For $m = 0$, the factor $M^{-(d-m)}$ in (5.11) has to be read as $(\log M)^{(n-1)(\frac{1}{2}+d)} M^{-d}$.

Proof. We prove the statement with the additional property that the index sets $\nabla_{k,M}^{(q)}$ have the *multiple tree property* introduced in Theorem 3.2 for subsets of $\nabla_{(\sigma,\bar{\sigma})}^{(q)}$, and that in the current generalized setting reads as: For any $(\lambda, p) \in \nabla_{k,M}^{(q)}$ and any $\mathbf{j} \in \mathbb{N}_0^n$ with $\mathbf{j} \leq |\lambda|$, there exists a $(\lambda', p') \in \nabla_{k,M}^{(q)}$ with $|\lambda'| = \mathbf{j}$, and $\text{supp } \psi_{\lambda,p}^{(q,k)} \cap \text{supp } \psi_{\lambda',p'}^{(q,k)} \neq \emptyset$.

For $q = 0$, the so extended statement is equal to that of Theorem 3.2. Let us assume that the statement is valid for some $0 \leq q-1 \leq N-1$.

To prove the statement for q , it is sufficient to consider $k = \bar{k}$. Let ϱ be a smooth function on \mathbb{R}^n such that for some sufficiently small $\varepsilon_2 > \varepsilon_1 > 0$, $\varrho \equiv 1$ within distance ε_1 of the interface J between $\hat{\Omega}_{k_1}^{(q-1)}$ and $\hat{\Omega}_{k_2}^{(q-1)}$, and vanishes outside distance ε_2 of J . Writing any function v on $\hat{\Omega}_{\bar{k}}^{(q)}$ as $\varrho v + (1-\varrho)v$ induces a stable splitting of $\hat{H}^m(\hat{\Omega}_{\bar{k}}^{(q)}) \cap \prod_{\square_{k'} \subset \hat{\Omega}_{\bar{k}}^{(q)}} \mathcal{H}_{m,\theta}^d(\square_{k'})$ into two subspaces.

For functions u of type $(1-\varrho)v$, one has $u|_{\hat{\Omega}_{k_2}^{(q-1)}} \in \hat{H}^m(\hat{\Omega}_{k_2}^{(q-1)})$, and, assuming ϱ to be sufficiently small, $\langle u|_{\hat{\Omega}_{k_1}^{(q-1)}}, \tilde{\psi}_{\lambda,p}^{(q-1,k_1)} \rangle_{L_2(\hat{\Omega}_{k_1}^{(q-1)})} = 0$ for all $(\lambda, p) \in \nabla_{k_1}^{(q-1)}$ with $\eta_1^{(q)} \psi_{\lambda,p}^{(q-1,k_1)} \neq E_1^{(q)} \psi_{\lambda,p}^{(q-1,k_1)}$. We conclude that for such functions (5.11) is

valid when

$$\nabla_{\bar{k},M}^{(q)} \supseteq \nabla_{k_1,M}^{(q-1)} \cup \nabla_{k_2,M}^{(q-1)}.$$

In the remainder of this proof, we consider functions of type $u = \varrho v$, so with support inside some sufficiently small neighborhood of J . For $q-1 \leq k \leq N$, we set the biorthogonal projectors

$$P_{k,M}^{(q-1)} : v \mapsto \sum_{(\lambda,p) \in \nabla_{k,M}^{(q-1)}} \langle v, \tilde{\psi}_{\lambda,p}^{(q-1,k)} \rangle_{L_2(\hat{\Omega}_k^{(q-1)})} \psi_{\lambda,p}^{(q-1,k)}.$$

W.l.o.g. we assume $J = \{0\} \times \check{J}$ and define the (scale-independent) extension $\hat{E}_1^{(q)}$ as $E_1^{(q)}$ with G_1 reading as \hat{G}_1 , defined by $\hat{G}_1 v(-x) = \sum_{l=0}^L \gamma_l v(2^l x)$ and $\hat{G}_1 v(x) = v(x)$ ($x \in \mathcal{I}$). So \hat{G}_1 is the Hestenes extension \check{G}_1 without the smooth cut-off function which is not needed here because of the assumption on $\text{supp } u$.

It holds that $R_2^{(q)}(\text{Id} - \hat{E}_1^{(q)} R_1^{(q)})u \in \mathring{H}^m(\hat{\Omega}_{k_2}^{(q-1)})$ and $R_1^{(q)}u \in \mathring{H}^m(\hat{\Omega}_{k_1}^{(q-1)})$. Since $\hat{E}_1^{(q)}$ preserves the piecewise weighted Sobolev smoothness of a function supported near the interface, we have

$$(5.12) \quad \begin{aligned} & \sum_{\square_{k'} \subset \hat{\Omega}_{k_2}^{(q-1)}} \|R_2^{(q)}(\text{Id} - \hat{E}_1^{(q)} R_1^{(q)})u\|_{\mathcal{H}_{m,\theta}^d(\square_{k'})}^2 + \sum_{\square_{k'} \subset \hat{\Omega}_{k_1}^{(q-1)}} \|R_1^{(q)}u\|_{\mathcal{H}_{m,\theta}^d(\square_{k'})}^2 \\ & \lesssim \sum_{\square_{k'} \subset \hat{\Omega}_{\bar{k}}^{(q)}} \|u\|_{\mathcal{H}_{m,\theta}^d(\square_{k'})}^2. \end{aligned}$$

Setting $u_1 := P_{k_1,M}^{(q-1)} R_1^{(q)}u$, $u_2 := P_{k_2,M}^{(q-1)} R_2^{(q)}(\text{Id} - \hat{E}_1^{(q)} R_1^{(q)})u$, from $[\hat{E}_1^{(q)} \quad \eta_2^{(q)}] \in B(\mathring{H}^m(\hat{\Omega}_{k_1}^{(q-1)}) \times \mathring{H}^m(\hat{\Omega}_{k_2}^{(q-1)}), \mathring{H}^m(\hat{\Omega}_{\bar{k}}^{(q)}))$ (see Proposition 4.2), we conclude that

$$(5.13) \quad \begin{aligned} & \|u - (\hat{E}_1^{(q)} u_1 + \eta_2^{(q)} u_2)\|_{H^m(\hat{\Omega}_{\bar{k}}^{(q)})} \\ & = \left\| [\hat{E}_1^{(q)} \quad \eta_2^{(q)}] \left(\begin{bmatrix} R_1^{(q)} \\ R_2^{(q)}(\text{Id} - \hat{E}_1^{(q)} R_1^{(q)}) \end{bmatrix} u - \begin{bmatrix} u_1 \\ u_2 \end{bmatrix} \right) \right\|_{H^m(\hat{\Omega}_{\bar{k}}^{(q)})} \\ & \lesssim \sqrt{\|R_1^{(q)}u - u_1\|_{H^m(\hat{\Omega}_{k_1}^{(q-1)})}^2 + \|R_2^{(q)}(\text{Id} - \hat{E}_1^{(q)} R_1^{(q)})u - u_2\|_{H^m(\hat{\Omega}_{k_2}^{(q-1)})}^2} \\ & \lesssim M^{-(d-m)} \sqrt{\sum_{\square_{k'} \subset \hat{\Omega}_{\bar{k}}^{(q)}} \|u\|_{\mathcal{H}_{m,\theta}^d(\square_{k'})}^2}, \end{aligned}$$

the last inequality by the induction hypothesis and (5.12).

Next, we write

$$(5.14) \quad u - (E_1^{(q)} u_1 + \eta_2^{(q)} u_2) = u - (\hat{E}_1^{(q)} u_1 + \eta_2^{(q)} u_2) + (\hat{E}_1^{(q)} - E_1^{(q)})u_1.$$

By construction of G_1 from \check{G}_1 , we have that $(\text{Id} - \eta_2^{(q)} R_2^{(q)})(\hat{E}_1^{(q)} - E_1^{(q)})u_1 = 0$, and $R_2^{(q)}(\hat{E}_1^{(q)} - E_1^{(q)})u_1 \in \mathring{H}^m(\hat{\Omega}_{k_2}^{(q-1)})$, and so

$$(\hat{E}_1^{(q)} - E_1^{(q)})u_1 = \sum_{(\hat{\lambda}, \hat{p}) \in \nabla_{k_2}^{(q-1)}} \langle R_2^{(q)}(\hat{E}_1^{(q)} - E_1^{(q)})u_1, \tilde{\psi}_{\hat{\lambda}, \hat{p}}^{(q-1,k_2)} \rangle_{L_2(\hat{\Omega}_{k_2}^{(q-1)})} \eta_2^{(q)} \psi_{\hat{\lambda}, \hat{p}}^{(q-1,k_2)}.$$

We set

$$\hat{\nabla}_{k_2, M}^{(q-1)} := \{(\hat{\lambda}, \hat{p}) \in \nabla_{k_2}^{(q-1)} : \langle R_2^{(q)}(\hat{E}_1^{(q)} - E_1^{(q)})u_1, \tilde{\psi}_{\hat{\lambda}, \hat{p}}^{(q-1, k_2)} \rangle_{L_2(\hat{\Omega}_{k_2}^{(q-1)})} \neq 0, \\ \text{for some } u_1 \in \mathfrak{S}P_{k_1, M}^{(q-1)}\}.$$

Below we will show that, even after a possible enlargement to ensure the multiple tree property, it holds that $\#\hat{\nabla}_{k_2, M}^{(q-1)} \lesssim \#\nabla_{k_1, M}^{(q-1)}$. Defining $\nabla_{\bar{k}, M}^{(q)} := \nabla_{k_1, M}^{(q-1)} \cup \nabla_{k_2, M}^{(q-1)} \cup \hat{\nabla}_{k_2, M}^{(q-1)}$, the proof is completed.

If $(\hat{\lambda}, \hat{p}) \in \hat{\nabla}_{k_2, M}^{(q-1)}$, then $\langle R_2^{(q)}\hat{E}_1^{(q)}\psi_{\lambda, p}^{(q-1, k_1)}, \tilde{\psi}_{\hat{\lambda}, \hat{p}}^{(q-1, k_2)} \rangle_{L_2(\hat{\Omega}_{k_2}^{(q-1)})} \neq 0$ for some $(\lambda, p) \in \nabla_{k_1, M}^{(q-1)}$ with $R_2^{(q)}\hat{E}_1^{(q)}\psi_{\lambda, p}^{(q-1, k_1)} \in \mathring{H}^m(\hat{\Omega}_{k_2}^{(q-1)})$. Using $Z_{|\lambda|}(\hat{\Omega}_{k_1}^{(q-1)}) \subset V_{|\lambda|+M_{q-1}\mathbf{e}}(\hat{\Omega}_{k_1}^{(q-1)})$ and the assumptions on the extension, we have

$$R_2^{(q)}\hat{E}_1^{(q)}\psi_{\lambda, p}^{(q-1, k_1)} \in V_{|\lambda|+(M_{q-1}+L)\mathbf{e}}(\hat{\Omega}_{k_2}^{(q-1)}) \subset Z_{|\lambda|+(M_{q-1}+m_{q-1}+L)\mathbf{e}}(\hat{\Omega}_{k_2}^{(q-1)})$$

and so $|\hat{\lambda}| \leq |\lambda| + (M_{q-1} + m_{q-1} + L)\mathbf{e}$. Here we applied both inclusions from Proposition 5.4(3).

Thanks to the multiple tree property of $\nabla_{k_1, M}^{(q-1)}$, there exists a $(\lambda', p') \in \nabla_{k_1, M}^{(q-1)}$ with $|\lambda'|_i = \min(|\hat{\lambda}|_i, |\lambda|_i)$ ($1 \leq i \leq n$) and $\text{supp } \psi_{\lambda', p'}^{(q-1, k_1)} \cap \text{supp } \psi_{\lambda, p}^{(q-1, k_1)} \neq \emptyset$. Note that because of $|\hat{\lambda}| \leq |\lambda| + (M_{q-1} + m_{q-1} + L)\mathbf{e}$, we have $|\lambda'| \leq |\hat{\lambda}| \leq |\lambda'| + (M_{q-1} + m_{q-1} + L)\mathbf{e}$.

The ‘‘localness’’ of $\Psi_{k_1}^{(q-1)}$ as given by Proposition 5.4(1), the assumptions on the extension, and the ‘‘locally finiteness’’ of $\tilde{\Psi}_{k_2}^{(q-1)}$ as given by Proposition 5.4(2) show that the number of $(\hat{\lambda}, \hat{p}) \in \hat{\nabla}_{k_2, M}^{(q-1)}$ with $\langle R_2^{(q)}\hat{E}_1^{(q)}\psi_{\lambda, p}^{(q-1, k_1)}, \tilde{\psi}_{\hat{\lambda}, \hat{p}}^{(q-1, k_2)} \rangle_{L_2(\hat{\Omega}_{k_2}^{(q-1)})} \neq 0$ on the *same* level $|\hat{\lambda}| \leq |\lambda| + (M_{q-1} + m_{q-1} + L)\mathbf{e}$ is uniformly bounded. With this, we conclude that with the above mapping $(\hat{\lambda}, \hat{p}) \mapsto (\lambda', p')$, an at most uniformly bounded number of $(\hat{\lambda}, \hat{p}) \in \hat{\nabla}_{k_2, M}^{(q-1)}$ is mapped onto any $(\lambda', p') \in \nabla_{k_1, M}^{(q-1)}$, and so that $\#\hat{\nabla}_{k_2, M}^{(q-1)} \lesssim \#\nabla_{k_1, M}^{(q-1)}$.

Finally, to bound $\#\hat{\nabla}_{k_2, M}^{(q-1)}$, we only used that for $(\hat{\lambda}, \hat{p}) \in \hat{\nabla}_{k_2, M}^{(q-1)}$ there exists a $(\lambda, p) \in \nabla_{k_1, M}^{(q-1)}$ with $\text{supp } R_2^{(q)}\hat{E}_1^{(q)}\psi_{\lambda, p}^{(q-1, k_1)} \cap \text{supp } \tilde{\psi}_{\hat{\lambda}, \hat{p}}^{(q-1, k_1)} \neq \emptyset$ and $|\hat{\lambda}| \leq |\lambda| + (M_{q-1} + m_{q-1} + L)\mathbf{e}$. The same proof would have applied with the condition about the non-empty intersection of the supports reading as the condition that $\text{supp } R_2^{(q)}\hat{E}_1^{(q)}\psi_{\lambda, p}^{(q-1, k_1)}$ has non-empty intersection with some hyperrectangle, containing $\text{supp } \tilde{\psi}_{\hat{\lambda}, \hat{p}}^{(q-1, k_1)}$, that is aligned with the Cartesian coordinates with sides of lengths of order $2^{-|\hat{\lambda}|_1}, \dots, 2^{-|\hat{\lambda}|_n}$. In view of this, if $\hat{\nabla}_{k_2, M}^{(q-1)}$ does not already have the multiple tree property, then it can be enlarged to have this property while retaining $\#\hat{\nabla}_{k_2, M}^{(q-1)} \lesssim \#\nabla_{k_1, M}^{(q-1)}$. \square

6. REGULARITY

We study the issue whether we may expect (5.1) for u being the solution of an elliptic boundary value problem of order $2m = 2$.

6.1. Two-dimensional case. Let Ω be a *polygonal* domain. This means that its boundary is the union of a finite number of line segments, known as *edges*, with ends known as *corners*. It is not assumed that Ω is a Lipschitz domain, so it may

contain cracks. We denote with \mathcal{E} the set of edges, with \mathcal{C} the set of corners, and set for $\mathbf{c} \in \mathcal{C}$,

$$r_{\mathbf{c}}(\mathbf{x}) := \text{dist}(\mathbf{x}, \mathbf{c}).$$

Following [CDN10], for $m \in \mathbb{N}_0$, we define the (non-homogeneous) *weighted Sobolev space* $J_{\beta}^m(\Omega)$ as the set of $u \in L_2^{\text{loc}}(\Omega)$ that have a finite squared norm

$$\|v\|_{J_{\beta}^m(\Omega)}^2 := \sum_{k=0}^m \sum_{|\alpha|=k} \|\{\prod_{\mathbf{c} \in \mathcal{C}} r_{\mathbf{c}}^{\beta+m}\} \partial^{\alpha} v\|_{L_2(\Omega)}^2.$$

(in [CDN10] the generalization is considered of β being possibly dependent on \mathbf{c}).

Let A be a constant, real, symmetric and positive definite 2×2 matrix. Let $\mathcal{E}_D \subset \mathcal{E}$, and

$$V(\Omega) := \begin{cases} \{v \in H^1(\Omega) : v|_{\mathbf{e}} = 0 \ \forall \mathbf{e} \in \mathcal{E}_D\} & \text{when } \mathcal{E}_D \neq \emptyset, \\ \{v \in H^1(\Omega) : \int_{\Omega} v \, dx = 0\} & \text{otherwise.} \end{cases}$$

Given $g \in V(\Omega)'$, let $u \in V(\Omega)$ denote the solution of

$$(6.1) \quad \int_{\Omega} A \nabla u \cdot \nabla v \, dx = g(v) \quad (v \in V(\Omega)).$$

Theorem 6.1. *For $m \in \mathbb{N}_0$, there exists a $b^* \in (0, m+2]$ such that for any $b \in [0, b^*)$, the mapping $g \mapsto u \in B(J_{-b+1}^m(\Omega), J_{-b-1}^{m+2}(\Omega))$.*

The proof follows from [CDN10, formula (6.7)]. As stated in [CDN10, Example 6.7], for m sufficiently large, $b^* > \frac{1}{4}$.

We refer to [CDN10, Sect. 7] for generalizations of Theorem 6.1 to differential operators with variable coefficients and/or lower order terms.

Concerning the smoothness condition on the right-hand side g , note that for $b \leq m+1$,

$$H^m(\Omega) \hookrightarrow J_{-b+1}^m(\Omega).$$

Let us now consider the situation that $\Omega = \cup_{i=1}^K \Omega_i$ is an essentially disjoint subdivision into subdomains, where $\Omega_i = \kappa_i(\square)$ with κ_i being a regular parametrization. Let R_i denote the restriction of functions on Ω to Ω_i .

Proposition 6.2. *If $\kappa_i \in C^{m+2}(\overline{\square})$ and $b \leq m+1$, then*

$$\kappa_i^* R_i \in B(J_{-b-1}^{m+2}(\Omega), J_{-b-1}^{m+2}(\square)).$$

Proof. This follows from the smoothness of κ_i , and from the fact that $\kappa_i^* u|_{\Omega_i}$ localized near corners of \square that do not correspond to corners of Ω is a function in $H^{m+2} \hookrightarrow J_{-b-1}^{m+2}$, the latter by $-b-1+m+2 \geq 0$. \square

The following Proposition demonstrates (5.1).

Proposition 6.3. *For $d \in \mathbb{N}_0$, $\theta \geq \max(1, d-b/2)$, it holds that*

$$J_{-b-1}^{2d}(\square) \hookrightarrow \mathcal{H}_{\theta-1}^d(\mathcal{I}) \otimes \mathcal{H}_{\theta}^d(\mathcal{I}) \cap \mathcal{H}_{\theta}^d(\mathcal{I}) \otimes \mathcal{H}_{\theta-1}^d(\mathcal{I}) = \mathcal{H}_{1,\theta}^d(\square).$$

Proof. This follows from $\max(x^{\theta-1}y^{\theta}, x^{\theta}y^{\theta-1}) \leq r_0^{2\theta-1} \leq r_0^{2d-b-1}$ when $r_0 \in [0, 1]$. \square

6.2. Three-dimensional case. As in the previous section we follow [CDN10] closely. Let Ω be a *polyhedral* domain. This means that its boundary is the union of a finite number of polygons, known as the *faces*; the segments forming their boundaries are the *edges*, and the ends of the edges are the *corners*. It is not assumed that Ω is a Lipschitz domain, so it may contain crack surfaces. We denote with \mathcal{F} , \mathcal{E} , and \mathcal{C} the set of faces, edges, and corners, respectively, and set for $\mathbf{e} \in \mathcal{E}$ and $\mathbf{c} \in \mathcal{C}$,

$$r_{\mathbf{e}}(\mathbf{x}) := \text{dist}(\mathbf{x}, \mathbf{e}), \quad r_{\mathbf{c}}(\mathbf{x}) := \text{dist}(\mathbf{x}, \mathbf{c}), \quad r_{\mathcal{C}}(\mathbf{x}) := \min_{\mathbf{c} \in \mathcal{C}} r_{\mathbf{c}}(\mathbf{x}), \quad r_{\mathcal{E}}(\mathbf{x}) := \min_{\mathbf{e} \in \mathcal{E}} r_{\mathbf{e}}(\mathbf{x}).$$

There exists an $\varepsilon > 0$ small enough such that if we set

$$\begin{aligned} \Omega_{\mathbf{e}} &:= \{\mathbf{x} \in \Omega : r_{\mathbf{e}}(\mathbf{x}) < \varepsilon, r_{\bar{\mathbf{e}}}(\mathbf{x}) > r_{\mathbf{e}}(\mathbf{x}) \ (\mathbf{e} \neq \bar{\mathbf{e}} \in \mathcal{E}), \text{ and } r_{\mathcal{C}}(\mathbf{x}) > \frac{\varepsilon}{2}\} \\ \Omega_{\mathbf{c}} &:= \{\mathbf{x} \in \Omega : r_{\mathbf{c}}(\mathbf{x}) < \varepsilon \text{ and } r_{\mathcal{E}}(\mathbf{x}) > \frac{\varepsilon}{2} r_{\mathbf{c}}(\mathbf{x})\} \\ \Omega_{\mathbf{ce}} &:= \{\mathbf{x} \in \Omega : r_{\mathbf{c}}(\mathbf{x}) < \varepsilon \text{ and } r_{\mathbf{e}}(\mathbf{x}) < \varepsilon r_{\mathbf{c}}(\mathbf{x})\} \\ \Omega_I &:= \{\mathbf{x} \in \Omega : r_{\mathcal{E}}(\mathbf{x}) > \frac{\varepsilon}{2}\} \end{aligned}$$

we have the following properties

$$\overline{\Omega_{\mathbf{e}}} \cap \overline{\Omega_{\mathbf{e}'}} = \emptyset, \quad \overline{\Omega_{\mathbf{ce}}} \cap \overline{\Omega_{\mathbf{ce}'}} = \{\mathbf{c}\} \quad (\mathbf{e} \neq \mathbf{e}' \in \mathcal{E}, \mathbf{c} \in \mathcal{C}),$$

$$\overline{B(\mathbf{c}; \varepsilon)} \cap \overline{B(\mathbf{c}'; \varepsilon)} = \emptyset \quad (\mathbf{c} \neq \mathbf{c}' \in \mathcal{C}), \quad \Omega = \Omega_I \cup_{\{\mathbf{c} \in \mathcal{C}\}} \Omega_{\mathbf{c}} \cup_{\{\mathbf{e} \in \mathcal{E}\}} \Omega_{\mathbf{e}} \cup_{\{\mathbf{c} \in \mathcal{C}, \mathbf{e} \in \mathcal{E}\}} \Omega_{\mathbf{ce}}.$$

In a neighborhood of any edge $\mathbf{e} \in \mathcal{E}$, we will take partial derivatives in an orthogonal coordinate system with one of the coordinate directions being parallel to e . For a multi-index α in that coordinate system, $|\alpha_{\perp}|$ will denote the sum of the coordinates in the directions perpendicular to e , and $|\alpha_{\parallel}| := |\alpha| - |\alpha_{\perp}|$.

For $m \in \mathbb{N}_0$, $\beta > -m$, and $\mathcal{E}_0 \subset \mathcal{E}$, we define the *anisotropic weighted Sobolev space*

(6.2)

$$\begin{aligned} N_{\beta}^m(\Omega, \mathcal{C}, \mathcal{E}_0) &:= \left\{ u \in L_2^{\text{loc}}(\Omega) : \forall \alpha, |\alpha| \leq m, \partial^{\alpha} u \in L_2(\Omega_I), \right. \\ &\quad r_{\mathbf{c}}(\mathbf{x})^{\beta+|\alpha|} \partial^{\alpha} u \in L_2(\Omega_{\mathbf{c}}) \quad \forall \mathbf{c} \in \mathcal{C}, \\ &\quad r_{\mathbf{e}}(\mathbf{x})^{\beta+|\alpha_{\perp}|} \partial^{\alpha} u \in L_2(\Omega_{\mathbf{e}}) \quad \forall \mathbf{e} \in \mathcal{E}_0, \\ &\quad r_{\mathbf{c}}(\mathbf{x})^{\beta+|\alpha|} (r_{\mathbf{e}}(\mathbf{x})/r_{\mathbf{c}}(\mathbf{x}))^{\beta+|\alpha_{\perp}|} \partial^{\alpha} u \in L_2(\Omega_{\mathbf{ce}}) \quad \forall \mathbf{c} \in \mathcal{C}, \mathbf{e} \in \mathcal{E}_0 \\ &\quad r_{\mathbf{e}}(\mathbf{x})^{\max(\beta+|\alpha_{\perp}|, 0)} \partial^{\alpha} u \in L_2(\Omega_{\mathbf{e}}) \quad \forall \mathbf{e} \in \mathcal{E} \setminus \mathcal{E}_0, \\ &\quad \left. r_{\mathbf{c}}(\mathbf{x})^{\beta+|\alpha|} (r_{\mathbf{e}}(\mathbf{x})/r_{\mathbf{c}}(\mathbf{x}))^{\max(\beta+|\alpha_{\perp}|, 0)} \partial^{\alpha} u \in L_2(\Omega_{\mathbf{ce}}) \quad \forall \mathbf{c} \in \mathcal{C}, \mathbf{e} \in \mathcal{E} \setminus \mathcal{E}_0 \right\}, \end{aligned}$$

with squared norm being the sum over $|\alpha| \leq m$ of the squared L_2 -norms over Ω_I , $\Omega_{\mathbf{c}}$, $\Omega_{\mathbf{e}}$, $\Omega_{\mathbf{ce}}$, and $\mathbf{c} \in \mathcal{C}$, $\mathbf{e} \in \mathcal{E}$, respectively. (As in the two-dimensional case, this definition can be generalized to β being possibly dependent on \mathbf{c} and \mathbf{e}).

The definition of $N_{\beta}^m(\Omega, \mathcal{C}, \mathcal{E}_0)$ is a special case of a definition of $N_{\beta}^m(\Omega, \mathcal{C}_0, \mathcal{E}_0)$ from [CDN10] for general $\mathcal{C}_0 \subseteq \mathcal{C}$. In particular, the definition of the (fully) non-homogeneous anisotropic weighted Sobolev space $N_{\beta}^m(\Omega) := N_{\beta}^m(\Omega, \emptyset, \emptyset)$ is obtained from (6.2) by taking $\mathcal{E}_0 = \emptyset$, and by replacing $r_{\mathbf{c}}(\mathbf{x})^{\beta+|\alpha|}$ by $r_{\mathbf{c}}(\mathbf{x})^{\max(\beta+|\alpha|, 0)}$ on all three occurrences. Obviously,

$$(6.3) \quad N_{\beta}^m(\Omega, \mathcal{C}, \mathcal{E}_0) \hookrightarrow N_{\beta}^m(\Omega).$$

Let A be a constant, real, symmetric and positive definite 3×3 matrix. Let $\mathcal{F}_D \subset \mathcal{F}$, and

$$V(\Omega) := \begin{cases} \{v \in H^1(\Omega) : v|_{\mathbf{f}} = 0 \ \forall \mathbf{f} \in \mathcal{F}_D\} & \text{when } \mathcal{F}_D \neq \emptyset, \\ \{v \in H^1(\Omega) : \int_{\Omega} v \, dx = 0\} & \text{otherwise.} \end{cases}$$

Given $g \in V(\Omega)'$, let $u \in V(\Omega)$ denote the solution of

$$(6.4) \quad \int_{\Omega} A \nabla u \cdot \nabla v \, dx = g(v) \quad (v \in V(\Omega)).$$

Theorem 6.4. *Let \mathcal{E}_0 be the set of all $\mathbf{e} \in \mathcal{E}$ that are an edge of an $\mathbf{f} \in \mathcal{F}_D$. There exists a $b^* \in (0, 1]$ such that for $m \in \mathbb{N}$, $m > 1$, and for any $b \in [0, b^*)$, the mapping $g \mapsto u \in B(N_{1-b}^m(\Omega, \mathcal{C}, \mathcal{E}_0), N_{-1-b}^m(\Omega, \mathcal{C}, \mathcal{E}_0))$.*

Indeed, with the isotropic weighted Sobolev spaces $J_{\beta}^m(\Omega)$ as defined in [CDN10, Def. 5.9] (where we consider the value of β to be independent of the edges and corners), [MR03, Th. 7.1] shows that $g \mapsto u \in B(J_{1-b}^0(\Omega), J_{1-b}^2(\Omega))$, and thus that $g \mapsto u \in B(J_{1-b}^0(\Omega), J_{1-b}^1(\Omega))$. Using that $N_{1-b}^m(\Omega, \mathcal{C}, \mathcal{E}_0) \hookrightarrow J_{1-b}^0(\Omega)$, we conclude the statement of the theorem from the *anisotropic regularity shift theorem* [CDN10, (5.25)(a)]. Here we used that the Assumptions 5.5 and 5.13 from [CDN10] for $\mathbf{e} \in \mathcal{E}_0$ or $\mathbf{e} \in \mathcal{E} \setminus \mathcal{E}_0$, respectively, are satisfied by an application of [MR03, Th. 7.2].

Concerning the smoothness condition on the right-hand side g , note that for $b \leq 1$,

$$H^m(\Omega) \hookrightarrow N_{1-b}^m(\Omega, \mathcal{C}, \mathcal{E}_0).$$

The fact, as proven in Thm. 6.4, that for sufficiently smooth right-hand side, the tangential derivatives of sufficiently high order along the edges of Ω of the solution of (6.4) are in the (unweighted) $L_2(\Omega)$ space, is essential for our goal of proving approximation rates with piecewise tensor product approximation as for one-dimensional problems.

Let us consider the situation that $\Omega = \cup_{i=1}^K \Omega_i$ is an essentially disjoint conforming subdivision into hexahedra that are images of \square under trilinear diffeomorphisms κ_i , with $\inf_{\mathbf{x} \in \square} |D\kappa_i(\mathbf{x})| > 0$.

Aiming at deriving a three dimensional analogue of Proposition 6.2, care has to be taken that for $\kappa_i^* R_i u$ to be in $N_{-1-b}^m(\square)$, its tangential derivatives along an edge up to order m have to be in $L_2(\square)$. Therefore, we have to ensure that if an edge of \square is mapped onto the boundary of Ω , then lines parallel to this edge are (smoothly) mapped onto lines parallel to the boundary of Ω .

Proposition 6.5. *Let for any i , κ_i be such that it maps an edge \mathbf{e} of \square to an edge of Ω , then it maps all three edges that are parallel to \mathbf{e} to edges that are parallel to $\kappa_i(\mathbf{e})$. Then*

$$\kappa_i^* R_i \in B(N_{-1-b}^m(\Omega), N_{-1-b}^m(\square)).$$

Proof. What has to be shown is that if an edge \mathbf{e} of \square is mapped to the boundary of Ω , then the tangential derivatives along $\mathbf{e} \circ \kappa_i$ up to order m are a smooth functions of the tangential derivatives of u along $\kappa_i(\mathbf{e})$ up to order m .

W.l.o.g., let \mathbf{e} be one of the edges $\mathbf{e}^{(1)}, \dots, \mathbf{e}^{(4)}$ that are parallel to the first unit vector. The vector $\partial_1 \kappa_i(\mathbf{x})$ is a bilinear function of x_2 and x_3 , and so in particular constant on each of the $\mathbf{e}^{(j)}$. These constant vectors are the differences of the

endpoints of $\kappa_i(\mathbf{e}^{(j)})$, and so, by assumption, multiples of $\partial_1 \kappa_i|_{\mathbf{e}}$. We conclude that $\partial_1 \kappa_i(\mathbf{x})$ is a multiple of a bilinear *scalar* function and $\partial_1 \kappa_i|_{\mathbf{e}}$. \square

Next we will show that the condition on the parametrizations imposed in Proposition 6.5 can always be satisfied by making some refinement of the initial conforming subdivision into hexahedra: Let us cut each hexahedron in the partition along 6 planes parallel to the 6 faces of the hexahedron on distance $\zeta > 0$, see Figure 5. When ζ is small enough, then the planes parallel to opposite faces of the




FIGURE 5. Hexahedron cut into 3^3 subhexahedra.

hexahedron do not intersect inside the hexahedron, and we obtain a subdivision of the hexahedron in 3^3 hexahedra. Eight of these hexahedra share a corner with the original hexahedron and so have three edges on edges of this hexahedron, and so possibly three edges on edges of Ω . These hexahedra are parallelepipeds and so satisfy the condition from Proposition 6.5. Twelve other hexahedra have one edge on an edge of the original hexahedron, and so possibly on an edge of Ω . For each of these hexahedra, the edges opposite to this specific edge are parallel to this edge and so satisfy the condition from Proposition 6.5. The remaining seven hexahedra have no edges on edges of the original hexahedron, and thus no edge on an edge of Ω . Six of them have a face on a face of the original hexahedron, whereas the boundary of the remaining “interior” hexahedron has empty intersection with the boundary of the original hexahedron.

Above subdivision of a hexahedron induces a subdivision of each of its faces into 3^2 quadrilaterals; 4 parallelograms at the corners, 4 trapezoids at the edges, and one interior quadrilateral. Conversely, such a subdivision of 3 non-opposite faces of the hexahedron, where the interior quadrilaterals are sufficiently large, determines uniquely the subdivision of the hexahedron into 3^3 subhexahedra by making cuts along planes parallel to the faces. So if we start with a subdivision of one hexahedron and use the resulting subdivision of its faces to induce subdivisions of its neighbors, then by choosing ζ small enough we obtain a refinement of the original conforming decomposition into hexahedra to a conforming decomposition into hexahedra that satisfy the conditions needed for Proposition 6.5.

What is left to show is whether the hexahedra in the refined subdivision are images of \square under trilinear diffeomorphisms κ_i , with $\inf_{\mathbf{x} \in \square} |D\kappa_i(\mathbf{x})| > 0$. When the aforementioned parameter ζ tends to zero, the interior hexahedron converges

to the hexahedron in the original decomposition, which was assumed to have this property. So for ζ small enough, the interior hexahedra have this property.

The other hexahedra in the refined subdivision have at least two parallel faces, and so are instances of a prismatoid. Let us consider such a hexahedron with its parallel faces, being convex quadrilaterals, on the planes $z = 0$ and $z = 1$. Let $q_1, q_2 : (0, 1)^2 \rightarrow \mathbb{R}^2$ be bilinear parametrizations of the bottom and top face with $\inf_{(x,y) \in (0,1)^2} |Dq_i(x,y)| > 0$, and such that the images of each corner of $(0, 1)^2$ under q_1 and q_2 are connected by an edge in the hexahedron. Then a trilinear parametrization $\square \rightarrow \mathbb{R}^3$ is given by

$$\kappa(x, y, z) = (1 - z)q_1(x, y) + zq_2(x, y)$$

and so $\inf_{(x,y,z) \in \square} |D\kappa(x, y, z)| = \inf_{(x,y,z) \in \square} |(1 - z)Dq_1(x, y) + zDq_2(x, y)| > 0$.

The following Proposition demonstrates (5.1).

Proposition 6.6. *For $d \in \mathbb{N}_0$, $\theta \geq \max(1, d - \frac{b}{3})$ where $b > 0$, it holds that*

$$N_{-1-b}^{3d}(\square) \hookrightarrow \mathcal{H}_{1,\theta}^d(\square).$$

Proof. It is sufficient to show continuity of the embedding of the spaces restricted to $\Omega_{\mathbf{c}}$, $\Omega_{\mathbf{e}}$, and $\Omega_{\mathbf{ce}}$ intersected with $(0, \frac{1}{2})^3$, where $\mathbf{c} = (0, 0, 0)$ and $\mathbf{e} = \mathbf{e}_1$.

For $\|\alpha\|_\infty \leq d$, the conditions on θ show that on $\Omega_{\mathbf{c}} \cap (0, \frac{1}{2})^3$,

$$\max(x^{\theta-1}y^\theta z^\theta, x^\theta y^{\theta-1}z^\theta, x^\theta y^\theta z^{\theta-1}) \leq r_{\mathbf{c}}(\mathbf{x})^{3\theta-1} \leq r_{\mathbf{c}}(\mathbf{x})^{\max(-1-b+|\alpha|, 0)},$$

and on $\Omega_{\mathbf{e}} \cap (0, \frac{1}{2})^3$,

$$\max(y^\theta z^\theta, y^{\theta-1}z^\theta, y^\theta z^{\theta-1}) \leq r_{\mathbf{e}}(\mathbf{x})^{2\theta-1} \leq r_{\mathbf{e}}(\mathbf{x})^{\max(-1-b+\alpha_2+\alpha_3, 0)}.$$

On $\Omega_{\mathbf{ce}} \cap (0, \frac{1}{2})^3$, we have

$$\max(x^{\theta-1}y^\theta z^\theta, x^\theta y^{\theta-1}z^\theta, x^\theta y^\theta z^{\theta-1}) \leq r_{\mathbf{c}}(\mathbf{x})^\theta r_{\mathbf{e}}(\mathbf{x})^{2\theta-1}.$$

To show that this right-hand side can be bounded on

$$r_{\mathbf{c}}(\mathbf{x})^{\max(-1-b+|\alpha|, 0) - \max(-1-b+\alpha_2+\alpha_3, 0)} r_{\mathbf{e}}(\mathbf{x})^{\max(-1-b+\alpha_2+\alpha_3, 0)}$$

we distinguish between 3 cases: For $-1 - b + |\alpha| \leq 0$, this results from $\theta \geq 0$ and $2\theta - 1 \geq 0$. For $-1 - b + |\alpha| \geq 0 \geq -1 - b + \alpha_2 + \alpha_3$, we have $r_{\mathbf{c}}(\mathbf{x})^\theta r_{\mathbf{e}}(\mathbf{x})^{2\theta-1} \leq r_{\mathbf{c}}(\mathbf{x})^{3\theta-1} \leq r_{\mathbf{c}}(\mathbf{x})^{-1-b+|\alpha|}$ by $\theta \geq d - \frac{b}{3}$. For $-1 - b + \alpha_2 + \alpha_3 \geq 0$, $r_{\mathbf{c}}(\mathbf{x})^\theta r_{\mathbf{e}}(\mathbf{x})^{2\theta-1} \leq r_{\mathbf{c}}(\mathbf{x})^{\theta+\frac{b}{3}} r_{\mathbf{e}}(\mathbf{x})^{2\theta-1-\frac{b}{3}} \leq r_{\mathbf{c}}(\mathbf{x})^{\alpha_1} r_{\mathbf{e}}(\mathbf{x})^{-1-b+\alpha_2+\alpha_3}$ by $\theta \geq d - \frac{b}{3}$. \square

7. NUMERICAL RESULTS

As the univariate building block of the piecewise tensor product wavelet construction, we apply the C^1 , piecewise quartic (so $d = 5$) (multi-) wavelets, with (discontinuous) piecewise quartic duals as constructed in [CS11]. The primal wavelets satisfy Dirichlet boundary conditions of order 1 at both boundaries 0 and 1, i.e., $\vec{\sigma} = (\sigma_\ell, \sigma_r) = (1, 1)$, whereas at the dual side no boundary conditions can be imposed, i.e., $\vec{\sigma} = (\tilde{\sigma}_\ell, \tilde{\sigma}_r) = (0, 0)$.

For the present work, we generalized this construction to obtain also wavelet collections that satisfy no boundary conditions (at primal side) at either or both boundaries, i.e., $\vec{\sigma} \in \{0, 1\}^2 \setminus \{(1, 1)\}$. Actually, we also slightly modified the biorthogonal collections $(\Psi_{(1,1),(0,0)}, \tilde{\Psi}_{(1,1),(0,0)})$ from [CS11] with the aim to minimize, for $\vec{\sigma} \in \{0, 1\}^2 \setminus \{(1, 1)\}$, the number of $\lambda \in \nabla_{\vec{\sigma}, (0,0)}$ for which either

$\psi_\lambda^{(\vec{\sigma},(0,0))} \notin \Psi_{(1,1),(0,0)}$ or $\tilde{\psi}_\lambda^{(\vec{\sigma},(0,0))} \notin \tilde{\Psi}_{(1,1),(0,0)}$. Indeed, recall from Remark 5.3 that the extension operator has to be applied to all primal wavelets with such indices λ (at either left or right boundary). We obtained the result that the number of such λ on each level at left or right boundary is equal to 2. One of them corresponds to a primal wavelet that does not vanish at the boundary and therefore has to be extended to obtain a continuous extension, whereas the primal wavelet corresponding to the other only has to be extended to guarantee locality of the resulting dual wavelets by an application of Proposition 5.2.

As extension operator, we apply the simple reflection suited for $1 \leq t < \frac{3}{2}$, $0 < \tilde{t} < \frac{1}{2}$.

As domains, we consider the 2-dimensional slit domain $\Omega = (0, 2)^2 \setminus \{1\} \times [1, 2]$, being the union of 4 squares from $\mathbb{Z}^2 + (0, 1)^2$, and the 3-dimensional L-shaped domain $\Omega = (0, 2) \times (0, 2) \times (0, 1) \setminus (1, 2) \times (1, 2) \times (0, 1)$, being the union of 3 cubes from $\mathbb{Z}^3 + (0, 1)^3$. Aiming at constructing Riesz bases for $[H^t(\Omega), H_0^t(\Omega)]_{s,2}$ ($s \in [0, 1]$), in particular for $H_0^1(\Omega)$, we impose homogeneous Dirichlet boundary conditions of order 1 at $\partial\Omega$.

In the slit domain case, we consider tensor product wavelet bases on the squares $(0, 1)^2$, $\{(1, 0)\} + (0, 1)^2$ with no boundary conditions on the left edge, and $\{(0, 1)\} + (0, 1)^2$ and $\{(1, 1)\} + (0, 1)^2$ with no boundary conditions on their bottom edge, with homogeneous Dirichlet boundary conditions of order 1 on all other edges. By applying the scale-dependent extension, first from $\{(1, 0)\} + (0, 1)^2$ to $(0, 2) \times (0, 1)$, and then from both top domains $\{(0, 1)\} + (0, 1)^2$ and $\{(1, 1)\} + (0, 1)^2$ over their bottom edges to Ω , we end up with the piecewise tensor product basis.

In the L-shaped domain case, we consider tensor product wavelet bases on the cubes $(0, 1)^3$, and $\{(1, 0, 0)\} + (0, 1)^3$ and $\{(0, 1, 0)\} + (0, 1)^3$ with no boundary conditions on their interface with $(0, 1)^3$, with homogeneous Dirichlet boundary conditions of order 1 on all other faces. By applying the scale-dependent extension from $\{(1, 0, 0)\} + (0, 1)^3$ to $(0, 2) \times (0, 1)^2$, and then from $\{(0, 1, 0)\} + (0, 1)^3$ to Ω , the piecewise tensor product basis is obtained.

Using these piecewise tensor product bases, we solved the Poisson problem of finding $u \in H_0^1(\Omega)$ such that

$$\int_{\Omega} \nabla u \cdot \nabla v = f(v) \quad (v \in H_0^1(\Omega))$$

by applying the *adaptive wavelet-Galerkin method* ([CDD01, Ste09]). This method is known to produce a sequence of approximations from the span of the basis that converge in $H^1(\Omega)$ -norm with the best possible rate. Assuming a sufficiently smooth right-hand side, Theorem 5.6 together with the regularity results from §6.1 or §6.2 show that this rate is $d - m = 5 - 1 = 4$ (indeed an even higher rate can generally not be expected).

Furthermore, if the bi-infinite stiffness matrix of the PDE w.r.t. the basis is sufficiently close to a sparse matrix, in the sense that it is *s*-compressible* for some $s^* > 4$, then this adaptive method has optimal computational complexity. The univariate wavelet basis from [CS11] was designed such that any second order PDE on $(0, 1)^n$ with homogeneous Dirichlet boundary conditions gives rise, w.r.t. the tensor product basis, to a bi-infinite stiffness matrix which is *truly sparse*. By losing the Dirichlet boundary conditions on one side of each interface between subdomains, and the application of reflections, this sparsity, however, is partly lost in the sense

that columns corresponding to wavelets that are not zero at an interface contain infinitely many non-zero entries. The sizes of these entries, however, decay that fast as function of the difference in levels of the wavelets involved that nevertheless the stiffness matrix is s^* -compressible with $s^* = \infty$, meaning that indeed the adaptive method has optimal computational complexity.

To investigate how the application of the extensions, and the incorporation of wavelets with other boundary conditions, affects the conditioning of the bi-infinite stiffness matrix, we computed numerically the condition number of the stiffness matrix (“preconditioned” by its diagonal) w.r.t. the piecewise tensor product wavelet basis on the slit domain, as well as that corresponding to the Poisson problem on $(0, 1)^2$ with the tensor product wavelet basis (with homogeneous Dirichlet boundary conditions). The results are given in Table 1.

J	0	1	2	3	4	5	6
square	14	37	60	86	110	131	149
slit domain	378	564	755	1077	1345	1437	1782

TABLE 1. Condition numbers of the diagonally preconditioned stiffness matrix on $(0, 1)^2$ with tensor product basis, and that on the slit domain $(0, 2)^2 \setminus \{1\} \times [1, 2]$ with piecewise tensor product basis. The bi-infinite stiffness matrices are restricted to the square block corresponding to row and column indices λ with $\|\lambda\|_\infty \leq J$. The cardinality of this set of row- or column-indices is (approximately) equal to $(3 \cdot 2^{J+1})^2$ or $4 \cdot (3 \cdot 2^{J+1})^2$, respectively.

We consider the Poisson problem on the slit domain for right-hand side $f = 1$. Its solution is illustrated in Figure 6. In Figure 7 we give the support lengths, in




FIGURE 6. The solution of the Poisson problem on the slit domain $(0, 2)^2 \setminus \{1\} \times [1, 2]$ with right-hand side $f = 1$.

piecewise tensor product wavelet coordinates, of the approximate solutions obtained by the adaptive wavelet-Galerkin scheme vs. the (relative) ℓ_2 -norm of their residual in the bi-infinite matrix vector system, the latter being equivalent to the $H^1(\Omega)$ -norm of the error. The optimal rate -4 indicated by the slope of the hypotenuse of the triangle is accurately approached for the problems sizes near the end of the computation. The centers of supports of the piecewise tensor product wavelets that were selected by the adaptive wavelet-Galerkin scheme are indicated Figure 8.




FIGURE 7. Support length vs. relative residual of the approximations produced by the adaptive wavelet-Galerkin scheme for the Poisson problem on the slit domain $(0, 2)^2 \setminus \{1\} \times [1, 2]$ with right-hand side $f = 1$ and the piecewise tensor product basis.




FIGURE 8. Centers of the supports of the piecewise tensor product wavelets that were selected by the adaptive wavelet-Galerkin scheme. The right picture is a zoom in of the left one.

Next we give numerical results for the Poisson problem on the L-shaped domain $\Omega = (0, 2) \times (0, 2) \times (0, 1) \setminus (1, 2) \times (1, 2) \times (0, 1)$ with right-hand side $f(x, y, z) = x(x - 2)y(y - 2)z$. In Figure 9 we give the support lengths, in piecewise tensor product wavelet coordinates, of the approximate solutions obtained by the adaptive wavelet-Galerkin scheme vs. the (relative) ℓ_2 -norm of their residual in the bi-infinite matrix vector system, the latter being equivalent to the $H^1(\Omega)$ -norm of the error. With the maximum problem size $\approx 2 \cdot 10^5$ that we could reach on our PC, the rate is still not very close to the asymptotic rate -4 . We expect that this is caused by the combination of a strong singularity in the solution, a three-dimensional domain, and the application of (piecewise) tensor product approximation. Indeed, although the asymptotic rate is independent of the spatial dimension, it is generally observed that with tensor product approximation in higher dimensions it takes an increasingly longer time before this rate becomes visible. Note that a rate -4 in

the $H^1(\Omega)$ -norm with an isotropic method would require approximation of order 13, if already attainable at all in view of regularity constraints.




FIGURE 9. Support length vs. relative residual of the approximations produced by the adaptive wavelet-Galerkin scheme for the Poisson problem on the slit domain L-shaped domain $\Omega = (0, 2) \times (0, 2) \times (0, 1) \setminus (1, 2) \times (1, 2) \times (0, 1)$ with right-hand side $f(x, y, z) = x(x - 2)y(y - 2)z$ and the piecewise tensor product basis.

The centers of supports of the piecewise tensor product wavelets that were selected by the adaptive wavelet-Galerkin scheme are indicated in Figure 10.




FIGURE 10. Centers of the supports of the piecewise tensor product wavelets that were selected by the adaptive wavelet-Galerkin scheme.

REFERENCES

- [Ape99] T. Apel. *Anisotropic finite elements: local estimates and applications*. Advances in Numerical Mathematics. B. G. Teubner, Stuttgart, 1999.

- [CDD01] A. Cohen, W. Dahmen, and R. DeVore. Adaptive wavelet methods for elliptic operator equations – Convergence rates. *Math. Comp.*, 70:27–75, 2001.
- [CDN10] M. Costabel, M. Dauge, and S. Nicaise. Analytic regularity for linear elliptic systems in polygons and polyhedra. Technical report, 2010. arXiv:1002.1772v1 [math.AP].
- [CF83] Z. Ciesielski and T. Figiel. Spline bases in classical function spaces on compact C^∞ manifolds. I and II. *Studia Math.*, 76(2):1–58 and 95–136, 1983.
- [CS11] N.G. Chegini and R.P. Stevenson. The adaptive tensor product wavelet scheme: Sparse matrices and the application to singularly perturbed problems. *IMA J. Numer. Anal.*, 2011. doi: 10.1093/imanum/drr013.
- [Dij09] T.J. Dijkema. *Adaptive tensor product wavelet methods for solving PDEs*. PhD thesis, Utrecht University, 2009.
- [DKU99] W. Dahmen, A. Kunoth, and K. Urban. Biorthogonal spline-wavelets on the interval - Stability and moment conditions. *Appl. Comp. Harm. Anal.*, 6:132–196, 1999.
- [DS99] W. Dahmen and R. Schneider. Wavelets on manifolds I: Construction and domain decomposition. *SIAM J. Math. Anal.*, 31:184–230, 1999.
- [DS10] M. Dauge and R.P. Stevenson. Sparse tensor product wavelet approximation of singular functions. *SIAM J. Math. Anal.*, 42(5):2203–2228, 2010.
- [Hes41] M.R. Hestenes. Extension of the range of differentiable functions. *Duke Math. J.*, 8(1):183–192, 1941.
- [KS06] A. Kunoth and J. Sahner. Wavelets on manifolds: An optimized construction. *Math. Comp.*, 75:1319–1349, 2006.
- [MR03] V. G. Maz'ya and J. Roßmann. Weighted L_p estimates of solutions to boundary value problems for second order elliptic systems in polyhedral domains. *ZAMM Z. Angew. Math. Mech.*, 83(7):435–467, 2003.
- [Nit05] P.-A. Nitsche. Sparse approximation of singularity functions. *Constr. Approx.*, 21(1):63–81, 2005.
- [Nit06] P.-A. Nitsche. Best N -term approximation spaces for tensor product wavelet bases. *Constr. Approx.*, 24(1):49–70, 2006.
- [Pri10] M. Primbs. New stable biorthogonal spline-wavelets on the interval. *Results Math.*, 57(1-2):121–162, 2010.
- [Ste09] R.P. Stevenson. Adaptive wavelet methods for solving operator equations: An overview. In R.A. DeVore and A. Kunoth, editors, *Multiscale, Nonlinear and Adaptive Approximation: Dedicated to Wolfgang Dahmen on the Occasion of his 60th Birthday*, pages 543–598. Springer, Berlin, 2009.
- [SU09] W. Sickel and T. Ullrich. Tensor products of Sobolev-Besov spaces and applications to approximation from the hyperbolic cross. *J. Approx. Theory*, 161:748–786, 2009.

KORTEWEG-DE VRIES INSTITUTE FOR MATHEMATICS, UNIVERSITY OF AMSTERDAM, P.O. BOX 94248, 1090 GE AMSTERDAM, THE NETHERLANDS

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCES, PHILIPPS-UNIVERSITY MARBURG, HANS-MEERWEIN STR., LAHNBERGE, 35032 MARBURG, GERMANY

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCES, PHILIPPS-UNIVERSITY MARBURG, HANS-MEERWEIN STR., LAHNBERGE, 35032 MARBURG, GERMANY

KORTEWEG-DE VRIES INSTITUTE FOR MATHEMATICS, UNIVERSITY OF AMSTERDAM, P.O. BOX 94248, 1090 GE AMSTERDAM, THE NETHERLANDS

E-mail address: n.godarzvandchegini@uva.nl, dahlke@mathematik.uni-marburg.de, friedrich@mathematik.uni-marburg.de, r.p.stevenson@uva.nl

Preprint Series DFG-SPP 1324

<http://www.dfg-spp1324.de>

Reports

- [1] R. Ramlau, G. Teschke, and M. Zhariy. A Compressive Landweber Iteration for Solving Ill-Posed Inverse Problems. Preprint 1, DFG-SPP 1324, September 2008.
- [2] G. Plonka. The Easy Path Wavelet Transform: A New Adaptive Wavelet Transform for Sparse Representation of Two-dimensional Data. Preprint 2, DFG-SPP 1324, September 2008.
- [3] E. Novak and H. Woźniakowski. Optimal Order of Convergence and (In-) Tractability of Multivariate Approximation of Smooth Functions. Preprint 3, DFG-SPP 1324, October 2008.
- [4] M. Espig, L. Grasedyck, and W. Hackbusch. Black Box Low Tensor Rank Approximation Using Fibre-Crosses. Preprint 4, DFG-SPP 1324, October 2008.
- [5] T. Bonesky, S. Dahlke, P. Maass, and T. Raasch. Adaptive Wavelet Methods and Sparsity Reconstruction for Inverse Heat Conduction Problems. Preprint 5, DFG-SPP 1324, January 2009.
- [6] E. Novak and H. Woźniakowski. Approximation of Infinitely Differentiable Multivariate Functions Is Intractable. Preprint 6, DFG-SPP 1324, January 2009.
- [7] J. Ma and G. Plonka. A Review of Curvelets and Recent Applications. Preprint 7, DFG-SPP 1324, February 2009.
- [8] L. Denis, D. A. Lorenz, and D. Tiede. Greedy Solution of Ill-Posed Problems: Error Bounds and Exact Inversion. Preprint 8, DFG-SPP 1324, April 2009.
- [9] U. Friedrich. A Two Parameter Generalization of Lions' Nonoverlapping Domain Decomposition Method for Linear Elliptic PDEs. Preprint 9, DFG-SPP 1324, April 2009.
- [10] K. Bredies and D. A. Lorenz. Minimization of Non-smooth, Non-convex Functionals by Iterative Thresholding. Preprint 10, DFG-SPP 1324, April 2009.
- [11] K. Bredies and D. A. Lorenz. Regularization with Non-convex Separable Constraints. Preprint 11, DFG-SPP 1324, April 2009.

- [12] M. Döhler, S. Kunis, and D. Potts. Nonequispaced Hyperbolic Cross Fast Fourier Transform. Preprint 12, DFG-SPP 1324, April 2009.
- [13] C. Bender. Dual Pricing of Multi-Exercise Options under Volume Constraints. Preprint 13, DFG-SPP 1324, April 2009.
- [14] T. Müller-Gronbach and K. Ritter. Variable Subspace Sampling and Multi-level Algorithms. Preprint 14, DFG-SPP 1324, May 2009.
- [15] G. Plonka, S. Tenorth, and A. Iske. Optimally Sparse Image Representation by the Easy Path Wavelet Transform. Preprint 15, DFG-SPP 1324, May 2009.
- [16] S. Dahlke, E. Novak, and W. Sickel. Optimal Approximation of Elliptic Problems by Linear and Nonlinear Mappings IV: Errors in L_2 and Other Norms. Preprint 16, DFG-SPP 1324, June 2009.
- [17] B. Jin, T. Khan, P. Maass, and M. Pidcock. Function Spaces and Optimal Currents in Impedance Tomography. Preprint 17, DFG-SPP 1324, June 2009.
- [18] G. Plonka and J. Ma. Curvelet-Wavelet Regularized Split Bregman Iteration for Compressed Sensing. Preprint 18, DFG-SPP 1324, June 2009.
- [19] G. Teschke and C. Borries. Accelerated Projected Steepest Descent Method for Nonlinear Inverse Problems with Sparsity Constraints. Preprint 19, DFG-SPP 1324, July 2009.
- [20] L. Grasedyck. Hierarchical Singular Value Decomposition of Tensors. Preprint 20, DFG-SPP 1324, July 2009.
- [21] D. Rudolf. Error Bounds for Computing the Expectation by Markov Chain Monte Carlo. Preprint 21, DFG-SPP 1324, July 2009.
- [22] M. Hansen and W. Sickel. Best m-term Approximation and Lizorkin-Triebel Spaces. Preprint 22, DFG-SPP 1324, August 2009.
- [23] F.J. Hickernell, T. Müller-Gronbach, B. Niu, and K. Ritter. Multi-level Monte Carlo Algorithms for Infinite-dimensional Integration on \mathbb{R}^N . Preprint 23, DFG-SPP 1324, August 2009.
- [24] S. Dereich and F. Heidenreich. A Multilevel Monte Carlo Algorithm for Lévy Driven Stochastic Differential Equations. Preprint 24, DFG-SPP 1324, August 2009.
- [25] S. Dahlke, M. Fornasier, and T. Raasch. Multilevel Preconditioning for Adaptive Sparse Optimization. Preprint 25, DFG-SPP 1324, August 2009.

- [26] S. Dereich. Multilevel Monte Carlo Algorithms for Lévy-driven SDEs with Gaussian Correction. Preprint 26, DFG-SPP 1324, August 2009.
- [27] G. Plonka, S. Tenorth, and D. Roşca. A New Hybrid Method for Image Approximation using the Easy Path Wavelet Transform. Preprint 27, DFG-SPP 1324, October 2009.
- [28] O. Koch and C. Lubich. Dynamical Low-rank Approximation of Tensors. Preprint 28, DFG-SPP 1324, November 2009.
- [29] E. Faou, V. Gradinaru, and C. Lubich. Computing Semi-classical Quantum Dynamics with Hagedorn Wavepackets. Preprint 29, DFG-SPP 1324, November 2009.
- [30] D. Conte and C. Lubich. An Error Analysis of the Multi-configuration Time-dependent Hartree Method of Quantum Dynamics. Preprint 30, DFG-SPP 1324, November 2009.
- [31] C. E. Powell and E. Ullmann. Preconditioning Stochastic Galerkin Saddle Point Problems. Preprint 31, DFG-SPP 1324, November 2009.
- [32] O. G. Ernst and E. Ullmann. Stochastic Galerkin Matrices. Preprint 32, DFG-SPP 1324, November 2009.
- [33] F. Lindner and R. L. Schilling. Weak Order for the Discretization of the Stochastic Heat Equation Driven by Impulsive Noise. Preprint 33, DFG-SPP 1324, November 2009.
- [34] L. Kämmerer and S. Kunis. On the Stability of the Hyperbolic Cross Discrete Fourier Transform. Preprint 34, DFG-SPP 1324, December 2009.
- [35] P. Cerejeiras, M. Ferreira, U. Kähler, and G. Teschke. Inversion of the noisy Radon transform on $SO(3)$ by Gabor frames and sparse recovery principles. Preprint 35, DFG-SPP 1324, January 2010.
- [36] T. Jahnke and T. Udrescu. Solving Chemical Master Equations by Adaptive Wavelet Compression. Preprint 36, DFG-SPP 1324, January 2010.
- [37] P. Kittipoom, G. Kutyniok, and W.-Q. Lim. Irregular Shearlet Frames: Geometry and Approximation Properties. Preprint 37, DFG-SPP 1324, February 2010.
- [38] G. Kutyniok and W.-Q. Lim. Compactly Supported Shearlets are Optimally Sparse. Preprint 38, DFG-SPP 1324, February 2010.

- [39] M. Hansen and W. Sickel. Best m -Term Approximation and Tensor Products of Sobolev and Besov Spaces – the Case of Non-compact Embeddings. Preprint 39, DFG-SPP 1324, March 2010.
- [40] B. Niu, F.J. Hickernell, T. Müller-Gronbach, and K. Ritter. Deterministic Multi-level Algorithms for Infinite-dimensional Integration on $\mathbb{R}^{\mathbb{N}}$. Preprint 40, DFG-SPP 1324, March 2010.
- [41] P. Kittipoom, G. Kutyniok, and W.-Q. Lim. Construction of Compactly Supported Shearlet Frames. Preprint 41, DFG-SPP 1324, March 2010.
- [42] C. Bender and J. Steiner. Error Criteria for Numerical Solutions of Backward SDEs. Preprint 42, DFG-SPP 1324, April 2010.
- [43] L. Grasedyck. Polynomial Approximation in Hierarchical Tucker Format by Vector-Tensorization. Preprint 43, DFG-SPP 1324, April 2010.
- [44] M. Hansen und W. Sickel. Best m -Term Approximation and Sobolev-Besov Spaces of Dominating Mixed Smoothness - the Case of Compact Embeddings. Preprint 44, DFG-SPP 1324, April 2010.
- [45] P. Binev, W. Dahmen, and P. Lamby. Fast High-Dimensional Approximation with Sparse Occupancy Trees. Preprint 45, DFG-SPP 1324, May 2010.
- [46] J. Ballani and L. Grasedyck. A Projection Method to Solve Linear Systems in Tensor Format. Preprint 46, DFG-SPP 1324, May 2010.
- [47] P. Binev, A. Cohen, W. Dahmen, R. DeVore, G. Petrova, and P. Wojtaszczyk. Convergence Rates for Greedy Algorithms in Reduced Basis Methods. Preprint 47, DFG-SPP 1324, May 2010.
- [48] S. Kestler and K. Urban. Adaptive Wavelet Methods on Unbounded Domains. Preprint 48, DFG-SPP 1324, June 2010.
- [49] H. Yserentant. The Mixed Regularity of Electronic Wave Functions Multiplied by Explicit Correlation Factors. Preprint 49, DFG-SPP 1324, June 2010.
- [50] H. Yserentant. On the Complexity of the Electronic Schrödinger Equation. Preprint 50, DFG-SPP 1324, June 2010.
- [51] M. Guillemard and A. Iske. Curvature Analysis of Frequency Modulated Manifolds in Dimensionality Reduction. Preprint 51, DFG-SPP 1324, June 2010.
- [52] E. Herrholz and G. Teschke. Compressive Sensing Principles and Iterative Sparse Recovery for Inverse and Ill-Posed Problems. Preprint 52, DFG-SPP 1324, July 2010.

- [53] L. Kämmerer, S. Kunis, and D. Potts. Interpolation Lattices for Hyperbolic Cross Trigonometric Polynomials. Preprint 53, DFG-SPP 1324, July 2010.
- [54] G. Kutyniok and W.-Q Lim. Shearlets on Bounded Domains. Preprint 54, DFG-SPP 1324, July 2010.
- [55] A. Zeiser. Wavelet Approximation in Weighted Sobolev Spaces of Mixed Order with Applications to the Electronic Schrödinger Equation. Preprint 55, DFG-SPP 1324, July 2010.
- [56] G. Kutyniok, J. Lemvig, and W.-Q Lim. Compactly Supported Shearlets. Preprint 56, DFG-SPP 1324, July 2010.
- [57] A. Zeiser. On the Optimality of the Inexact Inverse Iteration Coupled with Adaptive Finite Element Methods. Preprint 57, DFG-SPP 1324, July 2010.
- [58] S. Jokar. Sparse Recovery and Kronecker Products. Preprint 58, DFG-SPP 1324, August 2010.
- [59] T. Aboiyar, E. H. Georgoulis, and A. Iske. Adaptive ADER Methods Using Kernel-Based Polyharmonic Spline WENO Reconstruction. Preprint 59, DFG-SPP 1324, August 2010.
- [60] O. G. Ernst, A. Mugler, H.-J. Starkloff, and E. Ullmann. On the Convergence of Generalized Polynomial Chaos Expansions. Preprint 60, DFG-SPP 1324, August 2010.
- [61] S. Holtz, T. Rohwedder, and R. Schneider. On Manifolds of Tensors of Fixed TT-Rank. Preprint 61, DFG-SPP 1324, September 2010.
- [62] J. Ballani, L. Grasedyck, and M. Kluge. Black Box Approximation of Tensors in Hierarchical Tucker Format. Preprint 62, DFG-SPP 1324, October 2010.
- [63] M. Hansen. On Tensor Products of Quasi-Banach Spaces. Preprint 63, DFG-SPP 1324, October 2010.
- [64] S. Dahlke, G. Steidl, and G. Teschke. Shearlet Coorbit Spaces: Compactly Supported Analyzing Shearlets, Traces and Embeddings. Preprint 64, DFG-SPP 1324, October 2010.
- [65] W. Hackbusch. Tensorisation of Vectors and their Efficient Convolution. Preprint 65, DFG-SPP 1324, November 2010.
- [66] P. A. Cioica, S. Dahlke, S. Kinzel, F. Lindner, T. Raasch, K. Ritter, and R. L. Schilling. Spatial Besov Regularity for Stochastic Partial Differential Equations on Lipschitz Domains. Preprint 66, DFG-SPP 1324, November 2010.

- [67] E. Novak and H. Woźniakowski. On the Power of Function Values for the Approximation Problem in Various Settings. Preprint 67, DFG-SPP 1324, November 2010.
- [68] A. Hinrichs, E. Novak, and H. Woźniakowski. The Curse of Dimensionality for Monotone and Convex Functions of Many Variables. Preprint 68, DFG-SPP 1324, November 2010.
- [69] G. Kutyniok and W.-Q. Lim. Image Separation Using Shearlets. Preprint 69, DFG-SPP 1324, November 2010.
- [70] B. Jin and P. Maass. An Analysis of Electrical Impedance Tomography with Applications to Tikhonov Regularization. Preprint 70, DFG-SPP 1324, December 2010.
- [71] S. Holtz, T. Rohwedder, and R. Schneider. The Alternating Linear Scheme for Tensor Optimisation in the TT Format. Preprint 71, DFG-SPP 1324, December 2010.
- [72] T. Müller-Gronbach and K. Ritter. A Local Refinement Strategy for Constructive Quantization of Scalar SDEs. Preprint 72, DFG-SPP 1324, December 2010.
- [73] T. Rohwedder and R. Schneider. An Analysis for the DIIS Acceleration Method used in Quantum Chemistry Calculations. Preprint 73, DFG-SPP 1324, December 2010.
- [74] C. Bender and J. Steiner. Least-Squares Monte Carlo for Backward SDEs. Preprint 74, DFG-SPP 1324, December 2010.
- [75] C. Bender. Primal and Dual Pricing of Multiple Exercise Options in Continuous Time. Preprint 75, DFG-SPP 1324, December 2010.
- [76] H. Harbrecht, M. Peters, and R. Schneider. On the Low-rank Approximation by the Pivoted Cholesky Decomposition. Preprint 76, DFG-SPP 1324, December 2010.
- [77] P. A. Cioica, S. Dahlke, N. Döhring, S. Kinzel, F. Lindner, T. Raasch, K. Ritter, and R. L. Schilling. Adaptive Wavelet Methods for Elliptic Stochastic Partial Differential Equations. Preprint 77, DFG-SPP 1324, January 2011.
- [78] G. Plonka, S. Tenorth, and A. Iske. Optimal Representation of Piecewise Hölder Smooth Bivariate Functions by the Easy Path Wavelet Transform. Preprint 78, DFG-SPP 1324, January 2011.

- [79] A. Mugler and H.-J. Starkloff. On Elliptic Partial Differential Equations with Random Coefficients. Preprint 79, DFG-SPP 1324, January 2011.
- [80] T. Müller-Gronbach, K. Ritter, and L. Yaroslavtseva. A Derandomization of the Euler Scheme for Scalar Stochastic Differential Equations. Preprint 80, DFG-SPP 1324, January 2011.
- [81] W. Dahmen, C. Huang, C. Schwab, and G. Welper. Adaptive Petrov-Galerkin methods for first order transport equations. Preprint 81, DFG-SPP 1324, January 2011.
- [82] K. Grella and C. Schwab. Sparse Tensor Spherical Harmonics Approximation in Radiative Transfer. Preprint 82, DFG-SPP 1324, January 2011.
- [83] D.A. Lorenz, S. Schiffler, and D. Trede. Beyond Convergence Rates: Exact Inversion With Tikhonov Regularization With Sparsity Constraints. Preprint 83, DFG-SPP 1324, January 2011.
- [84] S. Dereich, M. Scheutzow, and R. Schottstedt. Constructive quantization: Approximation by empirical measures. Preprint 84, DFG-SPP 1324, January 2011.
- [85] S. Dahlke and W. Sickel. On Besov Regularity of Solutions to Nonlinear Elliptic Partial Differential Equations. Preprint 85, DFG-SPP 1324, January 2011.
- [86] S. Dahlke, U. Friedrich, P. Maass, T. Raasch, and R.A. Ressel. An adaptive wavelet method for parameter identification problems in parabolic partial differential equations. Preprint 86, DFG-SPP 1324, January 2011.
- [87] A. Cohen, W. Dahmen, and G. Welper. Adaptivity and Variational Stabilization for Convection-Diffusion Equations. Preprint 87, DFG-SPP 1324, January 2011.
- [88] T. Jahnke. On Reduced Models for the Chemical Master Equation. Preprint 88, DFG-SPP 1324, January 2011.
- [89] P. Binev, W. Dahmen, R. DeVore, P. Lamby, D. Savu, and R. Sharpley. Compressed Sensing and Electron Microscopy. Preprint 89, DFG-SPP 1324, March 2011.
- [90] P. Binev, F. Blanco-Silva, D. Blom, W. Dahmen, P. Lamby, R. Sharpley, and T. Vogt. High Quality Image Formation by Nonlocal Means Applied to High-Angle Annular Dark Field Scanning Transmission Electron Microscopy (HAADF-STEM). Preprint 90, DFG-SPP 1324, March 2011.
- [91] R. A. Ressel. A Parameter Identification Problem for a Nonlinear Parabolic Differential Equation. Preprint 91, DFG-SPP 1324, May 2011.

- [92] G. Kutyniok. Data Separation by Sparse Representations. Preprint 92, DFG-SPP 1324, May 2011.
- [93] M. A. Davenport, M. F. Duarte, Y. C. Eldar, and G. Kutyniok. Introduction to Compressed Sensing. Preprint 93, DFG-SPP 1324, May 2011.
- [94] H.-C. Kreuzler and H. Yserentant. The Mixed Regularity of Electronic Wave Functions in Fractional Order and Weighted Sobolev Spaces. Preprint 94, DFG-SPP 1324, June 2011.
- [95] E. Ullmann, H. C. Elman, and O. G. Ernst. Efficient Iterative Solvers for Stochastic Galerkin Discretizations of Log-Transformed Random Diffusion Problems. Preprint 95, DFG-SPP 1324, June 2011.
- [96] S. Kunis and I. Melzer. On the Butterfly Sparse Fourier Transform. Preprint 96, DFG-SPP 1324, June 2011.
- [97] T. Rohwedder. The Continuous Coupled Cluster Formulation for the Electronic Schrödinger Equation. Preprint 97, DFG-SPP 1324, June 2011.
- [98] T. Rohwedder and R. Schneider. Error Estimates for the Coupled Cluster Method. Preprint 98, DFG-SPP 1324, June 2011.
- [99] P. A. Cioica and S. Dahlke. Spatial Besov Regularity for Semilinear Stochastic Partial Differential Equations on Bounded Lipschitz Domains. Preprint 99, DFG-SPP 1324, July 2011.
- [100] L. Grasedyck and W. Hackbusch. An Introduction to Hierarchical (H-) Rank and TT-Rank of Tensors with Examples. Preprint 100, DFG-SPP 1324, August 2011.
- [101] N. Chegini, S. Dahlke, U. Friedrich, and R. Stevenson. Piecewise Tensor Product Wavelet Bases by Extensions and Approximation Rates. Preprint 101, DFG-SPP 1324, September 2011.