

DFG-Schwerpunktprogramm 1324

„Extraktion quantifizierbarer Information aus komplexen Systemen“

Discontinuous information in the worst case and randomized settings

A. Hinrichs, E. Novak, H. Woźniakowski

Preprint 106



Edited by

AG Numerik/Optimierung
Fachbereich 12 - Mathematik und Informatik
Philipps-Universität Marburg
Hans-Meerwein-Str.
35032 Marburg

DFG-Schwerpunktprogramm 1324

„Extraktion quantifizierbarer Information aus komplexen Systemen“

Discontinuous information in the worst case and randomized settings

A. Hinrichs, E. Novak, H. Woźniakowski

Preprint 106



The consecutive numbering of the publications is determined by their chronological order.

The aim of this preprint series is to make new research rapidly available for scientific discussion. Therefore, the responsibility for the contents is solely due to the authors. The publications will be distributed by the authors.

Discontinuous information in the worst case and randomized settings

Aicke Hinrichs, Erich Novak, Henryk Woźniakowski

September 16, 2011

Dedicated to Hans Triebel on the occasion of his 75th birthday

Abstract

We believe that discontinuous linear information is never more powerful than continuous linear information for approximating continuous operators. We prove such a result in the worst case setting. In the randomized setting we consider compact linear operators defined between Hilbert spaces. In this case, the use of discontinuous linear information in the randomized setting cannot be much more powerful than continuous linear information in the worst case setting. These results can be applied when function evaluations are used even if function values are defined only almost everywhere.

1 Introduction

We study the approximation of an operator S defined between normed spaces F and G . The operator S does not have to be linear or continuous. We approximate $S(f)$ by algorithms that use information consisting of finitely many continuous or discontinuous linear functionals $L_i : F \rightarrow \mathbb{R}$. The error of such algorithms is defined either in the worst case or randomized setting.

For continuous S , it is hard to imagine that one can learn about $S(f)$ by using discontinuous information. On the other hand, it is well known that the Monte Carlo algorithm works nicely for multivariate integration defined for L_2 -functions. This algorithm uses linear functionals given by function evaluations which are indeed discontinuous or even not always well defined. Hence, discontinuous information is actually used in computational practice and seems to be useful, at least in the randomized setting.

This is the subject of this paper. We want to verify the power of *discontinuous* linear information and compare it to the power of *continuous* linear approximation. We study the worst case and randomized settings. This is done by comparing the n th minimal (worst case and randomized) errors which we can achieve by using n discontinuous or continuous linear functionals.

In the worst case setting, we prove that as long as S is a continuous operator (not necessarily linear) then the n th minimal errors are exactly the same for the class $\tilde{\Lambda}^{\text{all}}$ of all discontinuous or continuous linear functionals and the class Λ^{all} of all continuous linear functionals, see Theorem 1. This means that the use of discontinuous linear functionals does not help. The situation is quite different if S is discontinuous. We present an easy example of a discontinuous linear functional S for which the n th minimal errors for the class $\tilde{\Lambda}^{\text{all}}$ are zero for all $n \geq 1$, whereas the n th minimal errors for the class Λ^{all} are infinity for all $n \geq 1$.

In the randomized setting, we mostly consider compact linear operators S defined between Hilbert spaces F and G . In this case, we know from [14] that the power of continuous linear functionals in the randomized setting is roughly the same as in the worst case setting, see Lemma 2. Here, the word “roughly” means that the n th minimal error in the randomized setting is at least as large as a half of the $(4n - 1)$ st minimal error in the worst case setting, and obviously it is at most as large as the n th minimal error in the worst case setting. By combining with the result from the worst case setting, we conclude that the power of discontinuous linear functionals in the randomized setting is roughly the same as the power of continuous linear functionals in the worst case setting. On the other hand, if we drop the assumption that S is a compact linear operator between Hilbert spaces then we can construct a problem S which is *not* solvable in the worst case setting and solvable and relatively easy in the randomized setting. Here, not solvable means that the n th minimal errors in the worst case setting do *not* converge to zero, and relatively easy means the n th minimal errors in the randomized setting are of order $n^{-1/2}$.

For many applications the class F consists of functions and we can only use function evaluations for the approximation of S . The class of such evaluations is called *standard* and denoted by Λ^{std} . These evaluations are always linear but not always continuous. That is, we always have $\Lambda^{\text{std}} \subseteq \tilde{\Lambda}^{\text{all}}$, and, depending on the space F , we sometimes have $\Lambda^{\text{std}} \subseteq \Lambda^{\text{all}}$. In either case, our results apply. In particular, if all function evaluations are discontinuous then they may be useless in the worst case setting since the minimal worst case error of any algorithm that uses n function values is as good as a constant algorithm that uses *no* function values, see Remark 2.

For some applications the space F consists of equivalence classes of functions that are equal almost everywhere. This is the case for $F_1 = L_2(D)$ for some $D \subseteq \mathbb{R}^d$. Then function evaluations are not even well defined. We extend our analysis also to such function

evaluations and show that again the same results as before hold.

2 Worst Case Setting

For arbitrary normed spaces F and G , consider an arbitrary operator $S : F \rightarrow G$ that does not have to be linear or continuous. We approximate $S(f)$ for f from the unit ball of F by algorithms that use finitely many linear functionals from Λ^{all} or from $\tilde{\Lambda}^{\text{all}}$, respectively. More precisely, we consider algorithms $A_n : F \rightarrow G$ given by

$$A_n(f) = \varphi_n(L_1(f), L_2(f), \dots, L_n(f)), \quad (1)$$

where n is a nonnegative integer, $\varphi_n : \mathbb{R}^n \rightarrow G$ is an arbitrary mapping, and $L_j \in \Lambda$, where $\Lambda \in \{\Lambda^{\text{all}}, \tilde{\Lambda}^{\text{all}}\}$. Hence, for $\Lambda = \Lambda^{\text{all}}$ we only use *continuous* linear functionals, whereas for $\Lambda = \tilde{\Lambda}^{\text{all}}$ we may also use *discontinuous* linear functionals.

The choice of L_j can be *nonadaptive* or *adaptive*. It is nonadaptive if the functionals L_j are the same for all $f \in F$, and it is adaptive if L_j depends on the already computed values $L_1(f), L_2(f), \dots, L_{j-1}(f)$. That is,

$$N(f) = (L_1(f), L_2(f), \dots, L_n(f))$$

is the information used by the algorithm A_n and $L_j = L_j(\cdot; L_1(f), L_2(f), \dots, L_{j-1}(f)) \in \Lambda$. If the choice of all L_j 's is independent of $f \in F$ then N is *nonadaptive* information, otherwise if at least one L_j varies with $f \in F$ then N is *adaptive* information. For $n = 0$, the mapping A_n is a constant element of the space G . More details can be found in e.g., [13, 16, 18]. We define the error of such algorithms by taking the worst case setting, i.e.,

$$e(A_n) = \sup_{\|f\|_F < 1} \|S(f) - A_n(f)\|_G.$$

Observe that it is enough that the operator S is defined on the open unit ball in F , not necessarily on the whole space F . We take the open unit ball instead of the more standard case of the closed unit ball of F in the definition of the worst case error since this includes also operators with singularities on the boundary of the unit ball. For linear continuous S , or more generally for S uniformly continuous on the closed unit ball of F , this does not make a difference.

We define the n th minimal errors of approximation of S in the worst case setting as follows.

Definition 1. For $n = 0$ and $n \in \mathbb{N} := \{1, 2, \dots\}$, let

$$e_n^{\text{all-wor}}(S) = \inf_{A_n \text{ with } L_j \in \Lambda^{\text{all}}} e(A_n)$$

and

$$\tilde{e}_n^{\text{all-wor}}(S) = \inf_{A_n \text{ with } L_j \in \tilde{\Lambda}^{\text{all}}} e(A_n).$$

For $n = 0$, we obtain

$$e_0^{\text{all-wor}}(S) = \tilde{e}_0^{\text{all-wor}}(S) = \inf_{g \in G} \sup_{\|f\|_F < 1} \|S(f) - g\|_G.$$

It is easy to see that the best algorithm is $A_0 = 0$ if we assume that $S(f) = -S(-f)$ for all $\|f\|_F < 1$. Then

$$e_0^{\text{all-wor}}(S) = \tilde{e}_0^{\text{all-wor}}(S) = \sup_{\|f\|_F < 1} \|S(f)\|_G.$$

The error $e_0^{\text{all-wor}}(S)$ is the initial error that can be achieved without computing any linear functional on the elements $f \in F$. Clearly,

$$\tilde{e}_n^{\text{all-wor}}(S) \leq e_n^{\text{all-wor}}(S) \quad \text{for all } n \in \mathbb{N}.$$

The sequences $\{\tilde{e}_n^{\text{all-wor}}(S)\}$ and $\{e_n^{\text{all-wor}}(S)\}$ are both non-increasing but not necessarily convergent to zero.

We will use the following fact from functional analysis, see, e.g., [1, Ch. 3].

Lemma 1. Assume that F is a normed space and $L \in \tilde{\Lambda}^{\text{all}}$ is discontinuous. Then for all real α the set $\{f \in F \mid L(f) = \alpha\}$ is dense in F .

We are ready to prove that discontinuous linear functionals do not help for the approximation of continuous operators in the worst case setting.

Theorem 1. Let F and G be normed spaces and let $S : F \rightarrow G$ be continuous. Then

$$e_n^{\text{all-wor}}(S) = \tilde{e}_n^{\text{all-wor}}(S).$$

Proof. We may assume that $\dim(F) = \infty$ since otherwise $\tilde{\Lambda}^{\text{all}} = \Lambda^{\text{all}}$ and there is nothing to prove. Consider arbitrary adaptive information $N = (L_1, L_2, \dots, L_n)$ with

$$L_j = L_j(\cdot; y_1, y_2, \dots, y_{j-1}) \in \tilde{\Lambda}^{\text{all}} \quad \text{and} \quad y_i = L_i(f; y_1, y_2, \dots, y_{i-1}) \quad \text{for } f \in F.$$

It is well known that the infimum of the worst case errors of algorithms A_n that use information N is given by the radius of information N ,

$$r(N) = \sup_{y \in N(F)} \text{rad}(\{S(f) \mid N(f) = y, \|f\|_F < 1\}), \quad (2)$$

where $\text{rad}(A) = \inf_{g \in G} \sup_{a \in A} \|g - a\|$ denotes the radius of a set $A \subseteq G$, see [18]. Without loss of generality we may assume that the linear functionals L_1, L_2, \dots, L_n are linearly independent since otherwise the choice of a linear functional, say, L_j that is linearly dependent on L_1, L_2, \dots, L_{j-1} does not increase our knowledge about the element f . This implies that we may assume that $N(F) = \mathbb{R}^n$.

For $y \in N(F) = \mathbb{R}^n$ and $j = 1, 2, \dots, n$, define

$$B_k = B_k(y) = \{f \in F \mid L_j(f) = y_j \text{ for } j = 1, 2, \dots, k\},$$

Then the B_k are affine subspaces of F .

For each affine subspace B of F , we associate the uniquely determined linear subspace \tilde{B} such that $B = f + \tilde{B}$ for any $f \in B$. It is easily seen that a linear functional on F is continuous on B if and only if it is continuous on \tilde{B} . In particular,

$$\tilde{B}_k = \tilde{B}_k(y) = \ker L_1 \cap \ker L_2 \cap \dots \cap \ker L_k,$$

and continuity of L_{k+1} on B_k is equivalent to continuity of L_{k+1} on \tilde{B}_k .

We may now further assume for $k = 1, \dots, n-1$ that the functional L_{k+1} satisfies the following condition:

either L_{k+1} is continuous on F or L_{k+1} is discontinuous on B_k .

Indeed, assume that L_{k+1} is continuous on B_k . Then it is also continuous on \tilde{B}_k . Let L be a continuous linear extension of L_{k+1} on \tilde{B}_k to the whole space F . Then $L_{k+1} - L$ is a linear functional which is 0 on \tilde{B}_k , so that $\ker(L_{k+1} - L) \supset \tilde{B}_k$. This implies that $L_{k+1} - L$ is in the span of L_1, L_2, \dots, L_k , and for some numbers a_j we have $L(f) = L_{k+1}(f) + \sum_{j=1}^k a_j L_j(f)$ for all f . Hence knowing $L_j(f)$ for $j = 1, 2, \dots, k$, we know $L(f)$ iff we know $L_{k+1}(f)$. This means that we can replace the functional L_{k+1} in the information N with the continuous functional L without essentially changing the information and without changing its radius.

We now define the information $N^* = (L_1^*, L_2^*, \dots, L_n^*)$ with adaptively chosen

$$L_j^* = L_j^*(\cdot; y_1^*, y_2^*, \dots, y_{j-1}^*) \in \Lambda^{\text{all}} \quad \text{and} \quad y_i^* = L_i^*(f; y_1^*, y_2^*, \dots, y_{i-1}^*) \text{ for } f \in F$$

such that $r(N^*) \leq r(N)$. Since N is arbitrary and N^* consists of continuous linear functionals, this will prove the theorem.

The functionals L_j^* are defined inductively. We define $L_1^* = 0$ if L_1 is discontinuous on F , and otherwise we take $L_1^* = L_1$. Observe that in the case $L_1^* = 0$ the next functional L_2^* cannot be chosen adaptively since $L_1^*(f) = 0$ for all $f \in F$. Therefore, in general, $L_2^* = L_2^*(\cdot, 0)$ will be different from $L_2 = L_2(\cdot; y_1)$, with $y_1 = L_1(f)$, even if L_2 is continuous. Assume now inductively that for all $j = 1, 2, \dots, k < n$, we have already defined

$$L_j^* = L_j^*(\cdot; y_1^*, y_2^*, \dots, y_{j-1}^*) \in \Lambda^{\text{all}} \quad \text{with} \quad y_j^* = L_j^*(f; y_1^*, y_2^*, \dots, y_{j-1}^*) \quad \text{for all } f \in F.$$

Let $L_{k+1} = L_{k+1}(\cdot; y_1^*, y_2^*, \dots, y_k^*) \in \tilde{\Lambda}^{\text{all}}$ be the next linear functional for the original information N . Let $B_k = B_k(y^*)$ be defined as above for $y = y^*$. Define also

$$A_k = A_k(y^*) = \{f \in F \mid L_j^*(f) = y_j^* \text{ for } j = 1, 2, \dots, k\}.$$

If L_{k+1} is continuous, we set $L_{k+1}^* = L_{k+1}$, if L_{k+1} is discontinuous on B_k , we define $L_{k+1}^* = 0$. In the latter case only $y_{k+1}^* = 0$ needs to be further considered. As shown above, this completes the definition of $N^* = (L_1^*, L_2^*, \dots, L_n^*)$ that consists of n continuous linear functionals L_j^* .

We now show inductively that $B_j \subseteq A_j$ and B_j is dense in A_j for all $j = 1, 2, \dots, n$.

Indeed, for $j = 1$ we have $B_1 = A_1$ if L_1 is continuous on F , and

$$B_1 = \{f \in F \mid L_1(f) = 0\} \subseteq A_1 = F$$

if L_1 is discontinuous on F . From Lemma 1 we know that B_1 is dense in A_1 .

Assume now that $B_j \subseteq A_j$ and B_j is dense in A_j for $j = 1, 2, \dots, k < n$ with $k \geq 1$. Consider first the case when L_{k+1} is continuous. Then we have $L_{k+1}^* = L_{k+1}$ and

$$\begin{aligned} A_{k+1} &= \{f \in A_k \mid L_{k+1}^*(f) = y_{k+1}^*\} \\ B_{k+1} &= \{f \in B_k \mid L_{k+1}(f) = y_{k+1}^*\} = \{f \in B_k \mid L_{k+1}^*(f) = y_{k+1}^*\}. \end{aligned}$$

Hence

$$A_{k+1} = A_k \cap C_k \quad \text{and} \quad B_{k+1} = B_k \cap C_k \quad \text{with} \quad C_k = \{f \in F \mid L_{k+1}^*(f) = y_{k+1}^*\}.$$

Clearly, $B_k \subseteq A_k$ implies that $B_{k+1} \subseteq A_{k+1}$. We need to show that if B_k is dense in A_k then $B_k \cap C_k$ is dense in $A_k \cap C_k$. This is obvious if $L_{k+1}^* = 0$. Assume then that $L_{k+1}^* \neq 0$. Take $f \in A_k \cap C_k$. Then for any positive ε there exists $f_\varepsilon \in B_k$ such that $\|f - f_\varepsilon\| \leq \varepsilon$. For $g_{k+1} \in F$ with $L_j(g_{k+1}) = 0$ for $j = 1, 2, \dots, k$ and $L_{k+1}^*(g_{k+1}) = 1$, define

$$\tilde{f}_\varepsilon = f_\varepsilon + (y_{k+1}^* - L_{k+1}^*(f_\varepsilon)) g_{k+1}.$$

Then $\tilde{f}_\varepsilon \in B_k \cap C_k$ and since $L_{k+1}^*(f) = y_{k+1}^*$ we have

$$\|f - \tilde{f}_\varepsilon\| \leq \|f - f_\varepsilon\| + |L_{k+1}^*(f) - L_{k+1}^*(f_\varepsilon)| \|g_{k+1}\| \leq \|f - f_\varepsilon\| (1 + \|L_{k+1}^*\| \|g_{k+1}\|).$$

Hence $B_k \cap C_k$ is dense in $A_k \cap C_k$, as needed.

Consider now the case when L_{k+1} is discontinuous on B_k . Then $L_{k+1}^* = 0$ and $y_{k+1}^* = 0$. We now have

$$A_{k+1} = A_k \quad \text{and} \quad B_{k+1} = B_k \cap C_k \quad \text{with} \quad C_k = \{f \in F \mid L_{k+1}(f) = 0\}.$$

Clearly $B_{k+1} \subseteq B_k \subseteq A_k = A_{k+1}$. Since L_{k+1} is discontinuous on B_k it is also discontinuous on F . Then Lemma 1 says that C_k is a linear subspace which is dense in F . We want to show that $B_k \cap C_k$ is dense in $A_k = A_{k+1}$. Similarly as before, we take $f \in A_k$. For any positive ε we can find $f_\varepsilon \in B_k$ such that $\|f - f_\varepsilon\| \leq \varepsilon$. If $L_{k+1}(f_\varepsilon) = 0$ then $f_\varepsilon \in B_k \cap C_k$ and we are done. Assume then that $L_{k+1}(f_\varepsilon) \neq 0$. We now choose

$$g_{k+1} \in \tilde{B}_k = \ker L_1 \cap \ker L_2 \cdots \cap \ker L_k \quad \text{with} \quad L_{k+1}(g_{k+1}) = 1.$$

Since L_{k+1} is discontinuous on B_k , it is also discontinuous on \tilde{B}_k , so the element g_{k+1} can be of an arbitrary small norm. We choose a nonzero g_{k+1} such that $\|g_{k+1}\| \leq \varepsilon/|L_{k+1}(f_\varepsilon)|$. Then

$$\tilde{f}_\varepsilon = f_\varepsilon - L_{k+1}(f_\varepsilon) g_{k+1}$$

belongs to $B_k \cap C_k$ and

$$\|f - \tilde{f}_\varepsilon\| \leq \|f - f_\varepsilon\| + |L_{k+1}(f_\varepsilon)| \|g_{k+1}\| \leq 2\varepsilon.$$

Hence, $B_k \cap C_k$ is dense in A_k , as needed. This completes the proof that $B_n = N^{-1}(y^*)$ is dense in $A_n = (N^*)^{-1}(y^*)$.

Due to continuity of S , the set $B(y^*) := \{S(f) \in G \mid N(f) = y^*, \|f\|_F < 1\}$ is dense in $A(y^*) := \{S(f) \in G \mid N^*(f) = y^*, \|f\|_F < 1\}$ and therefore $\text{rad}(A(y^*)) = \text{rad}(B(y^*))$. This holds for all $y^* \in N^*(F)$ and therefore $r(N^*) \leq r(N)$, as needed. \square

The assumption on continuity of S in Theorem 1 is needed. Indeed, assume that $\dim(F) = \infty$. Then there are discontinuous linear functionals $L : F \rightarrow \mathbb{R}$. Define $S = L$. Note that now $e_0^{\text{all-wor}}(S) = \tilde{e}_0^{\text{all-wor}}(S) = \infty$.

Clearly, the worst case error $A_1(f) = L(f) = S(f)$ is zero. Therefore $\tilde{e}_n^{\text{all-wor}}(S) = 0$ for all $n \geq 1$. On the other hand, we know that adaption does not help for linear functionals as proved by Bakhvalov. Furthermore, if we use n linear continuous nonadaptive functionals L_j then Smolyak's theorem tells us that the best φ in (1) is linear, i.e., there are some real

numbers a_j for which the algorithm $A_n(f) = \sum_{j=1}^n a_j L_j(f)$ minimizes the worst case error among all algorithms that use $N = (L_1, L_2, \dots, L_n)$. The results of Bakhvalov and Smolyak can be found in [13, 16, 17, 18]. However, $S(f) - A_n(f)$ is still a discontinuous linear functional and therefore its worst case error is infinite. Since this holds for all continuous N , we have $e_n^{\text{all-wor}}(S) = \infty$. Hence,

$$\tilde{e}_n^{\text{all-wor}}(S) = 0 < e_n^{\text{all-wor}}(S) = \infty \quad \text{for all } n \in \mathbb{N}.$$

Although Theorem 1 deals with adaptive information, it is known that adaptive information does *not* help for many problems. This holds for linear operators S defined over Hilbert spaces F or if S is a linear functional, whereas for linear operators defined over arbitrary normed spaces adaption may help at most by a factor of two. The reader may find a survey of such results in Chapter 4 of [16].

Remark 1. Seeing the proof of Theorem 1, we may think that the use of discontinuous linear functionals is useless for approximating continuous S . More precisely, assume that we use nonadaptive $N = (L_1, L_2, \dots, L_n)$ for which *all* L_j 's are discontinuous linear functionals. What is the radius of N ? Is it the same as the radius of zero information? This is not true. We now show that we can achieve the radius of nonadaptive information consisting of $n - 1$ *continuous* linear functionals. Indeed, let $N^* = (L_2^*, L_3^*, \dots, L_n^*)$ be nonadaptive continuous information, $L_j^* \in \Lambda^{\text{all}}$. Take $N = (L_1, L_1 + L_2^*, \dots, L_1 + L_n^*)$ with a discontinuous linear functional L_1 . Then all $L_1 + L_j^*$'s are discontinuous. However, if we compute $y_1 = L_1(f)$ and $y_j = L_1(f) + L_j^*(f)$ then we also know $L_j^*(f) = y_j - y_1$ for $j = 2, 3, \dots, n$. Hence, we know $N^*(f)$ and therefore $r(N) \leq r(N^*)$, as claimed.

It is interesting to see what happens if we apply the proof of Theorem 1 to N . Since L_1 is discontinuous we obtain $L_1^* = 0$. However, $L_1 + L_2^*$ on $B_1 = \{f \in F \mid L_1(f) = y_1\}$ is $y_1 + L_2^*$ and therefore it is *continuous*. Then we replace L_2 by L_2^* . Similarly, all L_j with $j \geq 2$ will be replaced by L_j^* . The proof of Theorem 1 shows that $r(N) = r(N^*)$.

Remark 2. Assume now that F is a space of functions $f : D \rightarrow \mathbb{R}$. We consider the class Λ^{std} of all function evaluations given by the linear functionals $L_x(f) = f(x)$ for $f \in F$. Let $e_n^{\text{std-wor}}(S)$ denote the n th minimal worst case errors of algorithm A_n with $L_j \in \Lambda^{\text{std}}$, i.e., algorithms that use at most n function evaluations.

Assume that L_x is *discontinuous* on

$$F \cap \ker L_{x_1} \cap \ker L_{x_2} \cap \dots \cap \ker L_{x_n}$$

for all $x \in D \setminus \{x_1, x_2, \dots, x_n\}$. For instance $F = L_2(D) \cap C(D)$ equipped with the $L_2(D)$ norm is such an example.

Then it is easy to check that the proof of Theorem 1 yields $L_j^* = 0$ for all j . That is,

$$e_n^{\text{std-wor}}(S) = e_0^{\text{std-wor}}(S) = e_0^{\text{all-wor}}(S).$$

Hence, in this case the use of function evaluations is completely useless.

However, similarly as in Remark 1, one can construct examples where all function evaluations are *discontinuous* on F but may be *continuous* on $F \cap \ker L_{x_1} \cap \ker L_{x_2} \cap \dots \cap \ker L_{x_n}$, and still they are useful. Indeed, consider

$$F = C([0, 1]) \cap \left\{ f \mid \int_0^1 f(x) dx = \frac{f(0) + f(1)}{2} \right\}$$

equipped with the L_2 norm. Then the integration problem $S(f) = \int_0^1 f(x) dx$ is continuous and all function evaluations are discontinuous on F . Nevertheless, $L_1(f) = f(1) = S(f)/2$ on $F \cap \ker L_0$ is continuous, and we can compute $S(f)$ exactly using two function values of f .

Remark 3. For a continuous linear S and a Banach space F , Theorem 1 can be proved modulo a factor of $\frac{1}{2}$ by using the known relations between the Gelfand numbers $c_n(S)$ and the minimal errors $e_n^{\text{all-wor}}(S)$, see [16] for a survey of related results. In particular, we use that

$$c_n(S) \leq e_n^{\text{all-wor}}(S) \leq 2c_n(S),$$

for any linear and continuous operator S , see [18, Section 5.4 of Chapter 4].

It is known, see [2, Prop. 2.7.5], that the Gelfand numbers $c_n(S)$ are local in the sense that

$$c_n(S) = \sup_M c_n(S|_M)$$

where the supremum is taken over all finite dimensional subspaces M and $S|_M$ is the restriction of S to M .

Altogether, we obtain the following inequality:

$$\begin{aligned} e_n^{\text{all-wor}}(S) &\leq 2c_n(S) = 2 \sup_M c_n(S|_M) \leq 2 \sup_M e_n^{\text{all-wor}}(S|_M) = 2 \sup_M \tilde{e}_n^{\text{all-wor}}(S|_M) \\ &\leq 2\tilde{e}_n^{\text{all-wor}}(S), \end{aligned}$$

i.e.

$$e_n^{\text{all-wor}}(S) \leq 2 \cdot \tilde{e}_n^{\text{all-wor}}(S).$$

We return to general continuous not necessarily linear operators S . We conjecture that the worst case error $e_n^{\text{all-wor}}(S)$ is itself a local quantity at least for compact operators S , i.e.,

$$e_n^{\text{all-wor}}(S) = \sup_M e_n^{\text{all-wor}}(S|_M), \quad (3)$$

where again the supremum is taken over all finite dimensional subspaces M of F , and $S|_M$ is the restriction of S to M . This would give another and a much shorter proof of Theorem 1. Indeed, (3) implies

$$e_n^{\text{all-wor}}(S) = \sup_M e_n^{\text{all-wor}}(S|_M) = \sup_M \tilde{e}_n^{\text{all-wor}}(S|_M) \leq \tilde{e}_n^{\text{all-wor}}(S),$$

as claimed.

Although we do not know if (3) holds, we easily conclude from the local property of the Gelfand numbers that at least the weak local property holds for a continuous linear S and a Banach space F , namely

$$e_n^{\text{all-wor}}(S) \leq c_n(S) = \sup_M c_N(S|_M) \leq 2 \sup_M e_n^{\text{all-wor}}(S|_M).$$

It is also known that the approximation numbers $a_n(S)$ are local as long as S is a compact operator or G is a dual space, see [2, Prop. 2.7.1 and 2.7.3]. That translates into the fact that for linear nonadaptive algorithms in the worst case setting, the use of discontinuous functionals does not help. A weak form (with a factor 5) is true for arbitrary G and S , see [2, Prop. 2.7.4].

Remark 4. Heinrich [3] proves relations between linear n -widths and approximation numbers and shows that they coincide for compact and absolutely convex subsets of a normed space. There is also an example showing that, in general, for relatively compact absolutely convex sets equality does not hold. The spirit of these results is similar, but there is a difference as can be seen from Proposition 1.3 of that paper: The aim is to compare general or continuous linear information applied to $g = S(f) \in G$, while we compare general or continuous linear information applied to $f \in F$.

3 Randomized Setting

We now deal with randomized algorithms. We consider, as in [16, Theorem 4.42], only measurable algorithms. Hence we use the following definitions.

A randomized algorithm A is a pair consisting of a probability space (Ω, Σ, μ) and a family $(N_\omega, \varphi_\omega)_{\omega \in \Omega}$ of mappings such that the following holds:

1. For each fixed $\omega \in \Omega$, the mapping $A_\omega = \varphi_\omega \circ N_\omega$ is a deterministic algorithm defined as before, based on adaptive information N_ω consisting of linear functionals from a class Λ .
2. Let $n(f, \omega)$ be the cardinality of the information N_ω for $f \in F$. We assume that the function n is measurable.
3. The mapping $(f, \omega) \mapsto \varphi_\omega(N_\omega(f)) \in G$ is measurable.

Let A be a randomized algorithm. Then the cardinality of A is defined as

$$n(A) = \sup_{\|f\|_F < 1} \int_{\Omega} n(f, \omega) \, d\mu(\omega),$$

whereas the error of A in the randomized setting is

$$e^{\text{ran}}(A) = \sup_{\|f\|_F < 1} \left(\int_{\Omega} \|S(f) - \varphi_\omega(N_\omega(f))\|^2 \, d\mu(\omega) \right)^{1/2}.$$

For $n \in \mathbb{N}$, we define the n th minimal error of S in the randomized setting by

$$e_n^{\text{all-ran}}(S) = \inf\{e^{\text{ran}}(A) : n(A) \leq n\},$$

if A uses linear functionals from $\Lambda = \Lambda^{\text{all}}$. Similarly we define $\tilde{e}_n^{\text{all-ran}}(S)$ with $\Lambda = \tilde{\Lambda}^{\text{all}}$.

Obviously, we can interpret deterministic algorithms as randomized with a singleton Ω . That is why the n th minimal errors in the randomized setting cannot be larger than the n th minima errors in the worst case setting,

$$\tilde{e}_n^{\text{all-ran}}(S) \leq \tilde{e}_n^{\text{all-wor}}(S) \quad \text{and} \quad e_n^{\text{all-ran}}(S) \leq e_n^{\text{all-wor}}(S).$$

Basically, there exists only one proof technique to obtain lower bounds for randomized algorithms, and this technique goes back to Bakhvalov, see Section 4.3.3 of [16]. The main point is to observe that the errors in the randomized setting cannot be smaller than the errors in the average case setting for an arbitrary probability measure on F .

If we restrict ourselves to measurable randomized algorithms, then

$$e_n^{\text{all-ran}}(S) \geq \frac{\sqrt{2}}{2} e^{\text{avg}}(2n, \varrho), \tag{4}$$

where $e^{\text{avg}}(2n, \varrho)$ denotes the $2n$ -th minimal average case error of deterministic algorithms that use at most $2n$ linear functionals from Λ^{all} and ϱ is an arbitrary (Borel) probability measure on F , for more details see Lemma 4.37 and Remark 4.41 of [16]. The next result is identical with Theorem 4.42 from [16], see also [14]. It is stated here with a proof since we need a small modification of this result.

Lemma 2. Assume that $S : F \rightarrow G$ is a compact linear operator between Hilbert spaces F and G . Then

$$\frac{1}{2} e_{4n-1}^{\text{all-wor}}(S) \leq e_n^{\text{all-ran}}(S).$$

Proof. We know that $S(e_i) = \sigma_i \tilde{e}_i$ with orthonormal $\{e_i\}$ in F and $\{\tilde{e}_i\}$ in G . Here, $\{\sigma_i\}$ is a sequence of non-increasing singular values σ_i of S and $\lim_i \sigma_i = 0$. We also know that $e_n^{\text{all-wor}}(S) = \sigma_{n+1}$. For $m > n$, consider the normed $(m-1)$ -dimensional Lebesgue measure ϱ_m on the unit sphere $E_m = \{\sum_{i=1}^m \alpha_i e_i : \alpha_i \in \mathbb{R}, \sum_{i=1}^m \alpha_i^2 = 1\}$. Then

$$A_n^* \left(\sum_{i=1}^{\infty} \alpha_i e_i \right) = \sum_{i=1}^n \sigma_i \alpha_i \tilde{e}_i$$

is the optimal algorithm using continuous linear information of cardinality n . This is true for the worst case setting, with error σ_{n+1} , as well as for the average case setting with respect to ϱ_m . Hence

$$e^{\text{avg}}(n, \varrho_m)^2 = \int_{E_m} \sum_{i=n+1}^m \sigma_i^2 \alpha_i^2 d\varrho_m(\alpha).$$

Since $\int_{E_m} \alpha_i^2 d\varrho_m(\alpha) = 1/m$ we obtain

$$e^{\text{avg}}(n, \varrho_m)^2 = \frac{1}{m} \sum_{i=n+1}^m \sigma_i^2.$$

If we put $m = 2n$ then we obtain

$$e^{\text{avg}}(n, \varrho_{2n}) \geq \frac{1}{2} \sqrt{2} \sigma_{2n}.$$

Together with (4), we obtain

$$e_n^{\text{all-ran}}(S) \geq \frac{1}{2} \sqrt{2} e^{\text{avg}}(2n, \varrho_{4n}) \geq \frac{1}{2} \sigma_{4n} = \frac{1}{2} e_{4n-1}^{\text{all-wor}}(S).$$

□

We stress that in the proof we only use finite dimensional subspaces of F and linear functionals on such subspaces. Of course, for finite dimensional spaces, we have $\tilde{\Lambda}^{\text{all}} = \Lambda^{\text{all}}$ and therefore we obtain the following result.

Theorem 2. Assume that $S : F \rightarrow G$ is a compact operator between Hilbert spaces. Then

$$\frac{1}{2} e_{4n-1}^{\text{all-wor}}(S) \leq \tilde{e}_n^{\text{all-ran}}(S) \leq e_n^{\text{all-wor}}(S).$$

In this sense, randomization as well as allowing discontinuous linear functionals from $\widetilde{\Lambda}^{\text{all}}$ does not (essentially) help for the approximation of compact operators between Hilbert spaces.

Remark 5. Assume that $S : F \rightarrow G$ is linear and F and G are normed spaces. We do not know whether

$$\lim_{n \rightarrow \infty} e_n^{\text{all-ran}}(S) = 0$$

implies that S is compact. It is shown in [10] that the embedding $I : \ell_1 \rightarrow \ell_\infty$ is a universal non-compact operator in the sense that it factors through any non-compact linear bounded operator $S : F \rightarrow G$ between Banach spaces F and G :

$$\begin{array}{ccc} F & \xrightarrow{S} & G \\ \uparrow V & & \downarrow U \\ \ell_1 & \xrightarrow{I} & \ell_\infty. \end{array}$$

Here, $I = USV$ for some linear bounded operators U and V . It follows that

$$e_n^{\text{all-ran}}(I) = e_n^{\text{all-ran}}(USV) \leq \|U\| e_n^{\text{all-ran}}(S) \|V\|.$$

Thus it is sufficient to decide whether

$$\lim_{n \rightarrow \infty} e_n^{\text{all-ran}}(I) > 0 \quad \text{or} \quad \lim_{n \rightarrow \infty} e_n^{\text{all-ran}}(I) = 0.$$

Remark 6. Also in the randomized setting it would be very interesting to know whether the error $e_n^{\text{all-ran}}(S)$ is a local quantity, i.e. whether

$$e_n^{\text{all-ran}}(S) = \sup_M e_n^{\text{all-ran}}(S|_M),$$

or at least

$$e_n^{\text{all-ran}}(S) \leq c \sup_M e_n^{\text{all-ran}}(S|_M)$$

for some constant c independent of n . This would lead to an analogue of Theorem 1 in the randomized setting.

Remark 7. It is interesting to mention that there is a continuous *nonlinear* operator $S : F \rightarrow G$ for normed spaces F and G which is solvable in the randomized setting but not in the worst case setting, i.e.,

$$\lim_n \widetilde{e}_n^{\text{all-wor}}(S) = \lim_n e_n^{\text{all-wor}}(S) > 0 \quad \text{and} \quad \lim_n e_n^{\text{all-ran}}(S) = 0.$$

Obviously, the first equality above follows from Theorem 1. Indeed, let

$$S : F := \ell_1 \rightarrow G := \mathbb{R}, \quad S(x) = \|x\|_2^2 \quad \text{for all } x \in F.$$

Clearly, S is a continuous nonlinear functional. Note that the constant algorithm $\frac{1}{2}$ has the worst case error $\frac{1}{2}$ since we have $S(x) \in [0, 1]$ for x from the unit ball of ℓ_1 . Let

$$c := \lim_{n \rightarrow \infty} e_n^{\text{all-wor}}(S).$$

Then $c \leq \frac{1}{2}$. We now show that $c > 0$. We will use the known result of Kashin about the Gelfand width $c_n(B_1^m, \ell_2^m)$ for the unit ball B_1^m of $x \in \mathbb{R}^m$ with $\|x\|_1 \leq 1$, and with the error measured in the ℓ_2 norm. Namely, there are two positive numbers $c_{1,2}$ and $C_{1,2}$ such that for all $n < m$ we have

$$c_{1,2} \min \left(1, \frac{\ln(m/n) + 1}{n} \right)^{1/2} \leq c_n(B_1^m, \ell_2^m) \leq C_{1,2} \min \left(1, \frac{\ln(m/n) + 1}{n} \right)^{1/2}.$$

This means that $\lim_{m \rightarrow \infty} c_n(B_1^m, \ell_2) \geq c_{1,2}$. Using the definition of the Gelfand width, we conclude that

$$\lim_{n \rightarrow \infty} \inf_{L_1, L_2, \dots, L_n \in \tilde{\Lambda}^{\text{all}}} \sup_{x \in \ell_1, L_j(x)=0, j=1,2,\dots,n, \|x\|_1 \leq 1} \|x\|_2 \geq c_{1,2}.$$

Take now arbitrary adaptive $N = (L_1, L_2, \dots, L_n)$ with linear functionals L_i . Then there exists x in the unit ball of ℓ_1 such that $L_j(x) = 0$ for $j = 1, 2, \dots, n$ and $\|x\|_2 \geq c_{1,2}/2$. Let $\alpha \in [0, 1]$. Then αx belongs to the unit ball of ℓ_1 and $N(\alpha x) = 0$. For an arbitrary algorithm $A = \varphi_n(N(\cdot))$ we have $A_n(\alpha x) = \varphi_n(0)$ and $S(\alpha x) - \varphi_n(0) = \alpha \|x\|_2^2 - \varphi_n(0)$. Therefore

$$e(A_n) \geq \max_{\alpha \in [0,1]} |\alpha \|x\|_2^2 - \varphi_n(0)| \geq \frac{1}{2} \|x\|_2^2 \geq \frac{1}{4} c_{1,2}.$$

Hence $e_n^{\text{all-wor}}(S) \geq \frac{1}{4} c_{1,2}$ for all n . This proves that $c > 0$, as claimed.

In the randomized setting, consider the (random) linear functional

$$L(x) = \sum_{k=1}^{\infty} (\pm) x_k$$

with random and independent signs of probability $\frac{1}{2}$. The variance of the random variable $L(x)$ is $\sigma^2(L(x)) = \|x\|_2^2$ and it can be easily estimated with independent copies L_i of L . It is well known that the ‘‘empirical variance’’

$$A_n(x) = \frac{1}{n-1} \sum_{i=1}^n \left(L_i - \sum_{k=1}^n L_k \right)^2$$

with independent copies L_i of L has expectation $S(x) = \|x\|_2^2$ and variance

$$\sigma^2(A_n(x)) \leq \frac{1}{n} \cdot \|x\|_4^4 \leq \frac{1}{n} \cdot \|x\|_1^4.$$

Therefore

$$e^{\text{ran}}(A_n) = \sup_{\|x\|_{\ell_1} < 1} \sigma(A_n(x)) \leq \frac{1}{\sqrt{n}},$$

and $\lim_n e_n^{\text{all-ran}}(S) = 0$, as claimed.

4 Function Values

So far we did not assume that F is a space of functions and we only compared continuous linear information with arbitrary linear information. Now we assume that F is a normed space of functions $f : D \rightarrow \mathbb{R}$ for some nonempty set D .

Let $L_x(f) = f(x)$ for all $f \in F$ and $x \in D$. Since $f(x)$ is well defined for all $f \in F$ and $x \in D$, the functionals L_x 's are linear but *not* necessarily continuous. This class of information is called *standard* and denoted by Λ^{std} . Obviously $\Lambda^{\text{std}} \subseteq \tilde{\Lambda}^{\text{all}}$, however, $\Lambda^{\text{std}} \subseteq \Lambda^{\text{all}}$ only if all L_x 's are continuous. Still Theorems 1 and 2 apply in this case.

We now consider a more general case when F is a space of equivalence classes of functions $f : D \rightarrow \mathbb{R}$. A major example is $F = L_2(D)$. Then L_x is not even well defined for $f \in F$. On the other hand, we know that for $F = L_2(D)$ the functional $L_x(\tilde{f}) = \tilde{f}(x)$ is well defined for each $\tilde{f} \in f$ and all x from D . Here $\tilde{f} \in f$ means that the well defined function \tilde{f} is in the equivalence class f . This type of information is successfully used for multivariate integration by the standard Monte Carlo algorithm M_n in the randomized setting. Here for $D = [0, 1]^d$, we have $M_n(f) = n^{-1} \sum_{j=1}^n f(x_j)$ for independent and uniformly distributed points x_j . Then

$$M_n(\tilde{f}_1) = M_n(\tilde{f}_2) \quad a.s.$$

if $\tilde{f}_1, \tilde{f}_2 \in f$ and one usually uses only algorithms with this property. We would like to extend the analysis presented in the previous section also for the case when the elements of F are equivalence classes of functions.

We argue as follows. As in Section 3, we assume that $S : F \rightarrow G$ is a compact linear operator between the Hilbert spaces F and G . We know that then

$$S(e_i) = \sigma_i e'_i$$

with a non-increasing sequence of singular values σ_i of S , $\lim_i \sigma_i = 0$, with orthonormal $\{e_i\}$ in F and orthonormal $\{e'_i\}$ in G . We also know that $e_n^{\text{all-wor}}(S) = \sigma_{n+1}$. Then

$$A_n^* \left(\sum_{i=1}^{\infty} \alpha_i e_i \right) = \sum_{i=1}^n \sigma_i \alpha_i e'_i$$

is an optimal algorithm using continuous linear information of cardinality n . This is also true if we replace F by the $n+k$ dimensional space $V_{n+k} = \text{span}(e_1, e_2, \dots, e_{n+k})$. Hence

$$\tilde{e}_n^{\text{all-wor}}(V_{n+k}) = e_n^{\text{all-wor}}(V_{n+k}) = \sigma_{n+1} = e_n^{\text{all-wor}}(S) \quad \text{for all } k \geq 1.$$

Suppose that the functions \tilde{e}_i are elements of the equivalence classes e_i , for all $i = 1, 2, \dots, n+k$. Then we have functions in $\tilde{V}_{n+k} = \text{span}(\tilde{e}_1, \tilde{e}_2, \dots, \tilde{e}_{n+k})$ that are well defined everywhere. With this assumption we only make the oracle more powerful, i.e., the lower bound is even stronger. In this sense we can think of Λ^{std} as a subset of $\tilde{\Lambda}^{\text{all}}$. By $e_n^{\text{std-wor}}(S)$ we denote the n minimal worst case errors of algorithms that use at most n function values for approximating S over \tilde{V}_{n+k} . We obtain the following corollary.

Corollary 1. Assume that $S : F \rightarrow G$ is a compact linear operator between Hilbert spaces F and G . Then

$$e_n^{\text{std-wor}}(S) \geq e_n^{\text{all-wor}}(S) \quad \text{for all } n \in \mathbb{N}.$$

The same can be said for randomized algorithms. In this case we take $k \geq 3n$, update the definition of $e_n^{\text{std-ran}}(S)$ and obtain the following corollary.

Corollary 2. Assume that $S : F \rightarrow G$ is a compact linear operator between Hilbert spaces F and G . Then

$$e_n^{\text{std-ran}}(S) \geq \frac{1}{2} e_{4n-1}^{\text{all-wor}}(S).$$

Remark 8. Note that we did *not* compare $e_n^{\text{std-ran}}(S)$ with $e_n^{\text{std-wor}}(S)$. We stress that randomization may help a lot for the class Λ^{std} . This holds if function evaluations are continuous and also if function evaluations are not continuous. Examples for both cases can be found in Chapter 17 of [17].

A major example is the embedding of a function space into another (larger) function space. The literature is very rich, see, e.g., [4, 5, 6, 7, 8, 11, 12, 15, 16, 19, 20]. One can study the classes Λ^{all} , $\tilde{\Lambda}^{\text{all}}$ as well as Λ^{std} in the worst case setting and in the randomized setting. In the randomized setting we do *not* know whether Λ^{all} and $\tilde{\Lambda}^{\text{all}}$ always lead to the same results since in Theorem 2 we assume that both F and G are Hilbert spaces. It is open what happens for general normed spaces F and G and if an analogue of Theorem 1 for the worst case setting also holds in the randomized setting.

We also add that the lower bounds in the randomized setting of Heinrich and Mathé for specific spaces F and G are valid not only for Λ^{all} but also for $\tilde{\Lambda}^{\text{all}}$, see [6, 7, 8, 11]. The reason is similar as above in the proof of Theorem 2. Namely, finite dimensional subspaces of F can be used for the lower bounds and here all linear functionals are continuous.

Added in proof: Stefan Heinrich [9] recently proved that (3) is true for compact operators, but not in general.

Acknowledgement: We thank Stefan Heinrich for valuable remarks.

References

- [1] B. Bollobás, *Linear Analysis*, Cambridge University Press, Cambridge, 1990.
- [2] B. Carl and I. Stephani, *Entropy, Compactness and the Approximation of Operators*, Cambridge University Press, Cambridge, 1990.
- [3] S. Heinrich, On the relation between linear n -widths and approximation numbers, *J. Approx. Th.* **58** (1989), 315–333.
- [4] S. Heinrich, Lower bounds for the complexity of Monte Carlo function approximation, *J. Complexity* **8** (1992), 277–300.
- [5] S. Heinrich, Randomized approximation of Sobolev embeddings, in: Monte Carlo and Quasi-Monte Carlo Methods 2006 (A. Keller, S. Heinrich, H. Niederreiter, eds.), Springer, Berlin, 2008, 445–459.
- [6] S. Heinrich, Randomized approximation of Sobolev embeddings II, *J. Complexity* **25** (2009), 455–472.
- [7] S. Heinrich, Randomized approximation of Sobolev embeddings III, *J. Complexity* **25** (2009), 473–507.
- [8] S. Heinrich, Stochastic approximation of functions and applications, in: Monte Carlo and Quasi-Monte Carlo Methods 2010 (L. Plaskota, H. Woźniakowski, eds.), Springer, Berlin, to appear.
- [9] S. Heinrich, Ultrastability of n -th minimal errors, Preprint.
- [10] W. B. Johnson, A universal non-compact operator, *Coll. Mathematicum* **23** (1971), 267–268.

- [11] P. Mathé, Random approximation of Sobolev embeddings, *J. Complexity* **7** (1991), 261–281.
- [12] P. Mathé, A minimax principle for the optimal error of Monte Carlo methods, *Constr. Approx.* **9** (1993), 23–39.
- [13] E. Novak, *Deterministic and Stochastic Error Bounds in Numerical Analysis*, LNIM **1349**, Springer-Verlag, Berlin, 1988.
- [14] E. Novak, Optimal linear randomized methods for linear operators in Hilbert spaces, *J. Complexity* **8** (1992), 22–36.
- [15] E. Novak, H. Triebel, Function spaces in Lipschitz domains and optimal rates of convergence for sampling, *Constr. Approx.* **23** (2006), 325–350.
- [16] E. Novak and H. Woźniakowski, *Tractability of Multivariate Problems*, Volume I, Linear Information, European Math. Soc., Zürich, 2008.
- [17] E. Novak and H. Woźniakowski, *Tractability of Multivariate Problems*, Volume II: Standard Information for Functionals, European Math. Soc. Publ. House, Zürich, 2010.
- [18] J. F. Traub, G. W. Wasilkowski and H. Woźniakowski, *Information-Based Complexity*, Academic Press, New York, 1988.
- [19] H. Triebel, *Bases in Function Spaces, Sampling, Discrepancy, Numerical Integration*, European Math. Soc., Zürich, 2010.
- [20] J. Vybíral, Sampling numbers and function spaces, *J. Complexity* **23** (2007), 773–792.

Preprint Series DFG-SPP 1324

<http://www.dfg-spp1324.de>

Reports

- [1] R. Ramlau, G. Teschke, and M. Zhariy. A Compressive Landweber Iteration for Solving Ill-Posed Inverse Problems. Preprint 1, DFG-SPP 1324, September 2008.
- [2] G. Plonka. The Easy Path Wavelet Transform: A New Adaptive Wavelet Transform for Sparse Representation of Two-dimensional Data. Preprint 2, DFG-SPP 1324, September 2008.
- [3] E. Novak and H. Woźniakowski. Optimal Order of Convergence and (In-) Tractability of Multivariate Approximation of Smooth Functions. Preprint 3, DFG-SPP 1324, October 2008.
- [4] M. Espig, L. Grasedyck, and W. Hackbusch. Black Box Low Tensor Rank Approximation Using Fibre-Crosses. Preprint 4, DFG-SPP 1324, October 2008.
- [5] T. Bonesky, S. Dahlke, P. Maass, and T. Raasch. Adaptive Wavelet Methods and Sparsity Reconstruction for Inverse Heat Conduction Problems. Preprint 5, DFG-SPP 1324, January 2009.
- [6] E. Novak and H. Woźniakowski. Approximation of Infinitely Differentiable Multivariate Functions Is Intractable. Preprint 6, DFG-SPP 1324, January 2009.
- [7] J. Ma and G. Plonka. A Review of Curvelets and Recent Applications. Preprint 7, DFG-SPP 1324, February 2009.
- [8] L. Denis, D. A. Lorenz, and D. Trede. Greedy Solution of Ill-Posed Problems: Error Bounds and Exact Inversion. Preprint 8, DFG-SPP 1324, April 2009.
- [9] U. Friedrich. A Two Parameter Generalization of Lions' Nonoverlapping Domain Decomposition Method for Linear Elliptic PDEs. Preprint 9, DFG-SPP 1324, April 2009.
- [10] K. Bredies and D. A. Lorenz. Minimization of Non-smooth, Non-convex Functionals by Iterative Thresholding. Preprint 10, DFG-SPP 1324, April 2009.
- [11] K. Bredies and D. A. Lorenz. Regularization with Non-convex Separable Constraints. Preprint 11, DFG-SPP 1324, April 2009.

- [12] M. Döhler, S. Kunis, and D. Potts. Nonequispaced Hyperbolic Cross Fast Fourier Transform. Preprint 12, DFG-SPP 1324, April 2009.
- [13] C. Bender. Dual Pricing of Multi-Exercise Options under Volume Constraints. Preprint 13, DFG-SPP 1324, April 2009.
- [14] T. Müller-Gronbach and K. Ritter. Variable Subspace Sampling and Multi-level Algorithms. Preprint 14, DFG-SPP 1324, May 2009.
- [15] G. Plonka, S. Tenorth, and A. Iske. Optimally Sparse Image Representation by the Easy Path Wavelet Transform. Preprint 15, DFG-SPP 1324, May 2009.
- [16] S. Dahlke, E. Novak, and W. Sickel. Optimal Approximation of Elliptic Problems by Linear and Nonlinear Mappings IV: Errors in L_2 and Other Norms. Preprint 16, DFG-SPP 1324, June 2009.
- [17] B. Jin, T. Khan, P. Maass, and M. Pidcock. Function Spaces and Optimal Currents in Impedance Tomography. Preprint 17, DFG-SPP 1324, June 2009.
- [18] G. Plonka and J. Ma. Curvelet-Wavelet Regularized Split Bregman Iteration for Compressed Sensing. Preprint 18, DFG-SPP 1324, June 2009.
- [19] G. Teschke and C. Borries. Accelerated Projected Steepest Descent Method for Nonlinear Inverse Problems with Sparsity Constraints. Preprint 19, DFG-SPP 1324, July 2009.
- [20] L. Grasedyck. Hierarchical Singular Value Decomposition of Tensors. Preprint 20, DFG-SPP 1324, July 2009.
- [21] D. Rudolf. Error Bounds for Computing the Expectation by Markov Chain Monte Carlo. Preprint 21, DFG-SPP 1324, July 2009.
- [22] M. Hansen and W. Sickel. Best m -term Approximation and Lizorkin-Triebel Spaces. Preprint 22, DFG-SPP 1324, August 2009.
- [23] F.J. Hickernell, T. Müller-Gronbach, B. Niu, and K. Ritter. Multi-level Monte Carlo Algorithms for Infinite-dimensional Integration on \mathbb{R}^N . Preprint 23, DFG-SPP 1324, August 2009.
- [24] S. Dereich and F. Heidenreich. A Multilevel Monte Carlo Algorithm for Lévy Driven Stochastic Differential Equations. Preprint 24, DFG-SPP 1324, August 2009.
- [25] S. Dahlke, M. Fornasier, and T. Raasch. Multilevel Preconditioning for Adaptive Sparse Optimization. Preprint 25, DFG-SPP 1324, August 2009.

- [26] S. Dereich. Multilevel Monte Carlo Algorithms for Lévy-driven SDEs with Gaussian Correction. Preprint 26, DFG-SPP 1324, August 2009.
- [27] G. Plonka, S. Tenorth, and D. Roşca. A New Hybrid Method for Image Approximation using the Easy Path Wavelet Transform. Preprint 27, DFG-SPP 1324, October 2009.
- [28] O. Koch and C. Lubich. Dynamical Low-rank Approximation of Tensors. Preprint 28, DFG-SPP 1324, November 2009.
- [29] E. Faou, V. Gradinaru, and C. Lubich. Computing Semi-classical Quantum Dynamics with Hagedorn Wavepackets. Preprint 29, DFG-SPP 1324, November 2009.
- [30] D. Conte and C. Lubich. An Error Analysis of the Multi-configuration Time-dependent Hartree Method of Quantum Dynamics. Preprint 30, DFG-SPP 1324, November 2009.
- [31] C. E. Powell and E. Ullmann. Preconditioning Stochastic Galerkin Saddle Point Problems. Preprint 31, DFG-SPP 1324, November 2009.
- [32] O. G. Ernst and E. Ullmann. Stochastic Galerkin Matrices. Preprint 32, DFG-SPP 1324, November 2009.
- [33] F. Lindner and R. L. Schilling. Weak Order for the Discretization of the Stochastic Heat Equation Driven by Impulsive Noise. Preprint 33, DFG-SPP 1324, November 2009.
- [34] L. Kämmerer and S. Kunis. On the Stability of the Hyperbolic Cross Discrete Fourier Transform. Preprint 34, DFG-SPP 1324, December 2009.
- [35] P. Cerejeiras, M. Ferreira, U. Kähler, and G. Teschke. Inversion of the noisy Radon transform on $SO(3)$ by Gabor frames and sparse recovery principles. Preprint 35, DFG-SPP 1324, January 2010.
- [36] T. Jahnke and T. Udrescu. Solving Chemical Master Equations by Adaptive Wavelet Compression. Preprint 36, DFG-SPP 1324, January 2010.
- [37] P. Kittipoom, G. Kutyniok, and W.-Q. Lim. Irregular Shearlet Frames: Geometry and Approximation Properties. Preprint 37, DFG-SPP 1324, February 2010.
- [38] G. Kutyniok and W.-Q. Lim. Compactly Supported Shearlets are Optimally Sparse. Preprint 38, DFG-SPP 1324, February 2010.

- [39] M. Hansen and W. Sickel. Best m -Term Approximation and Tensor Products of Sobolev and Besov Spaces – the Case of Non-compact Embeddings. Preprint 39, DFG-SPP 1324, March 2010.
- [40] B. Niu, F.J. Hickernell, T. Müller-Gronbach, and K. Ritter. Deterministic Multi-level Algorithms for Infinite-dimensional Integration on $\mathbb{R}^{\mathbb{N}}$. Preprint 40, DFG-SPP 1324, March 2010.
- [41] P. Kittipoom, G. Kutyniok, and W.-Q. Lim. Construction of Compactly Supported Shearlet Frames. Preprint 41, DFG-SPP 1324, March 2010.
- [42] C. Bender and J. Steiner. Error Criteria for Numerical Solutions of Backward SDEs. Preprint 42, DFG-SPP 1324, April 2010.
- [43] L. Grasedyck. Polynomial Approximation in Hierarchical Tucker Format by Vector-Tensorization. Preprint 43, DFG-SPP 1324, April 2010.
- [44] M. Hansen und W. Sickel. Best m -Term Approximation and Sobolev-Besov Spaces of Dominating Mixed Smoothness - the Case of Compact Embeddings. Preprint 44, DFG-SPP 1324, April 2010.
- [45] P. Binev, W. Dahmen, and P. Lamby. Fast High-Dimensional Approximation with Sparse Occupancy Trees. Preprint 45, DFG-SPP 1324, May 2010.
- [46] J. Ballani and L. Grasedyck. A Projection Method to Solve Linear Systems in Tensor Format. Preprint 46, DFG-SPP 1324, May 2010.
- [47] P. Binev, A. Cohen, W. Dahmen, R. DeVore, G. Petrova, and P. Wojtaszczyk. Convergence Rates for Greedy Algorithms in Reduced Basis Methods. Preprint 47, DFG-SPP 1324, May 2010.
- [48] S. Kestler and K. Urban. Adaptive Wavelet Methods on Unbounded Domains. Preprint 48, DFG-SPP 1324, June 2010.
- [49] H. Yserentant. The Mixed Regularity of Electronic Wave Functions Multiplied by Explicit Correlation Factors. Preprint 49, DFG-SPP 1324, June 2010.
- [50] H. Yserentant. On the Complexity of the Electronic Schrödinger Equation. Preprint 50, DFG-SPP 1324, June 2010.
- [51] M. Guillemard and A. Iske. Curvature Analysis of Frequency Modulated Manifolds in Dimensionality Reduction. Preprint 51, DFG-SPP 1324, June 2010.
- [52] E. Herrholz and G. Teschke. Compressive Sensing Principles and Iterative Sparse Recovery for Inverse and Ill-Posed Problems. Preprint 52, DFG-SPP 1324, July 2010.

- [53] L. Kämmerer, S. Kunis, and D. Potts. Interpolation Lattices for Hyperbolic Cross Trigonometric Polynomials. Preprint 53, DFG-SPP 1324, July 2010.
- [54] G. Kutyniok and W.-Q Lim. Shearlets on Bounded Domains. Preprint 54, DFG-SPP 1324, July 2010.
- [55] A. Zeiser. Wavelet Approximation in Weighted Sobolev Spaces of Mixed Order with Applications to the Electronic Schrödinger Equation. Preprint 55, DFG-SPP 1324, July 2010.
- [56] G. Kutyniok, J. Lemvig, and W.-Q Lim. Compactly Supported Shearlets. Preprint 56, DFG-SPP 1324, July 2010.
- [57] A. Zeiser. On the Optimality of the Inexact Inverse Iteration Coupled with Adaptive Finite Element Methods. Preprint 57, DFG-SPP 1324, July 2010.
- [58] S. Jokar. Sparse Recovery and Kronecker Products. Preprint 58, DFG-SPP 1324, August 2010.
- [59] T. Aboiyar, E. H. Georgoulis, and A. Iske. Adaptive ADER Methods Using Kernel-Based Polyharmonic Spline WENO Reconstruction. Preprint 59, DFG-SPP 1324, August 2010.
- [60] O. G. Ernst, A. Mugler, H.-J. Starkloff, and E. Ullmann. On the Convergence of Generalized Polynomial Chaos Expansions. Preprint 60, DFG-SPP 1324, August 2010.
- [61] S. Holtz, T. Rohwedder, and R. Schneider. On Manifolds of Tensors of Fixed TT-Rank. Preprint 61, DFG-SPP 1324, September 2010.
- [62] J. Ballani, L. Grasedyck, and M. Kluge. Black Box Approximation of Tensors in Hierarchical Tucker Format. Preprint 62, DFG-SPP 1324, October 2010.
- [63] M. Hansen. On Tensor Products of Quasi-Banach Spaces. Preprint 63, DFG-SPP 1324, October 2010.
- [64] S. Dahlke, G. Steidl, and G. Teschke. Shearlet Coorbit Spaces: Compactly Supported Analyzing Shearlets, Traces and Embeddings. Preprint 64, DFG-SPP 1324, October 2010.
- [65] W. Hackbusch. Tensorisation of Vectors and their Efficient Convolution. Preprint 65, DFG-SPP 1324, November 2010.
- [66] P. A. Cioica, S. Dahlke, S. Kinzel, F. Lindner, T. Raasch, K. Ritter, and R. L. Schilling. Spatial Besov Regularity for Stochastic Partial Differential Equations on Lipschitz Domains. Preprint 66, DFG-SPP 1324, November 2010.

- [67] E. Novak and H. Woźniakowski. On the Power of Function Values for the Approximation Problem in Various Settings. Preprint 67, DFG-SPP 1324, November 2010.
- [68] A. Hinrichs, E. Novak, and H. Woźniakowski. The Curse of Dimensionality for Monotone and Convex Functions of Many Variables. Preprint 68, DFG-SPP 1324, November 2010.
- [69] G. Kutyniok and W.-Q. Lim. Image Separation Using Shearlets. Preprint 69, DFG-SPP 1324, November 2010.
- [70] B. Jin and P. Maass. An Analysis of Electrical Impedance Tomography with Applications to Tikhonov Regularization. Preprint 70, DFG-SPP 1324, December 2010.
- [71] S. Holtz, T. Rohwedder, and R. Schneider. The Alternating Linear Scheme for Tensor Optimisation in the TT Format. Preprint 71, DFG-SPP 1324, December 2010.
- [72] T. Müller-Gronbach and K. Ritter. A Local Refinement Strategy for Constructive Quantization of Scalar SDEs. Preprint 72, DFG-SPP 1324, December 2010.
- [73] T. Rohwedder and R. Schneider. An Analysis for the DIIS Acceleration Method used in Quantum Chemistry Calculations. Preprint 73, DFG-SPP 1324, December 2010.
- [74] C. Bender and J. Steiner. Least-Squares Monte Carlo for Backward SDEs. Preprint 74, DFG-SPP 1324, December 2010.
- [75] C. Bender. Primal and Dual Pricing of Multiple Exercise Options in Continuous Time. Preprint 75, DFG-SPP 1324, December 2010.
- [76] H. Harbrecht, M. Peters, and R. Schneider. On the Low-rank Approximation by the Pivoted Cholesky Decomposition. Preprint 76, DFG-SPP 1324, December 2010.
- [77] P. A. Cioica, S. Dahlke, N. Döhring, S. Kinzel, F. Lindner, T. Raasch, K. Ritter, and R. L. Schilling. Adaptive Wavelet Methods for Elliptic Stochastic Partial Differential Equations. Preprint 77, DFG-SPP 1324, January 2011.
- [78] G. Plonka, S. Tenorth, and A. Iske. Optimal Representation of Piecewise Hölder Smooth Bivariate Functions by the Easy Path Wavelet Transform. Preprint 78, DFG-SPP 1324, January 2011.

- [79] A. Mugler and H.-J. Starkloff. On Elliptic Partial Differential Equations with Random Coefficients. Preprint 79, DFG-SPP 1324, January 2011.
- [80] T. Müller-Gronbach, K. Ritter, and L. Yaroslavtseva. A Derandomization of the Euler Scheme for Scalar Stochastic Differential Equations. Preprint 80, DFG-SPP 1324, January 2011.
- [81] W. Dahmen, C. Huang, C. Schwab, and G. Welper. Adaptive Petrov-Galerkin methods for first order transport equations. Preprint 81, DFG-SPP 1324, January 2011.
- [82] K. Grella and C. Schwab. Sparse Tensor Spherical Harmonics Approximation in Radiative Transfer. Preprint 82, DFG-SPP 1324, January 2011.
- [83] D.A. Lorenz, S. Schiffler, and D. Trede. Beyond Convergence Rates: Exact Inversion With Tikhonov Regularization With Sparsity Constraints. Preprint 83, DFG-SPP 1324, January 2011.
- [84] S. Dereich, M. Scheutzow, and R. Schottstedt. Constructive quantization: Approximation by empirical measures. Preprint 84, DFG-SPP 1324, January 2011.
- [85] S. Dahlke and W. Sickel. On Besov Regularity of Solutions to Nonlinear Elliptic Partial Differential Equations. Preprint 85, DFG-SPP 1324, January 2011.
- [86] S. Dahlke, U. Friedrich, P. Maass, T. Raasch, and R.A. Ressel. An adaptive wavelet method for parameter identification problems in parabolic partial differential equations. Preprint 86, DFG-SPP 1324, January 2011.
- [87] A. Cohen, W. Dahmen, and G. Welper. Adaptivity and Variational Stabilization for Convection-Diffusion Equations. Preprint 87, DFG-SPP 1324, January 2011.
- [88] T. Jahnke. On Reduced Models for the Chemical Master Equation. Preprint 88, DFG-SPP 1324, January 2011.
- [89] P. Binev, W. Dahmen, R. DeVore, P. Lamby, D. Savu, and R. Sharpley. Compressed Sensing and Electron Microscopy. Preprint 89, DFG-SPP 1324, March 2011.
- [90] P. Binev, F. Blanco-Silva, D. Blom, W. Dahmen, P. Lamby, R. Sharpley, and T. Vogt. High Quality Image Formation by Nonlocal Means Applied to High-Angle Annular Dark Field Scanning Transmission Electron Microscopy (HAADF-STEM). Preprint 90, DFG-SPP 1324, March 2011.
- [91] R. A. Ressel. A Parameter Identification Problem for a Nonlinear Parabolic Differential Equation. Preprint 91, DFG-SPP 1324, May 2011.

- [92] G. Kutyniok. Data Separation by Sparse Representations. Preprint 92, DFG-SPP 1324, May 2011.
- [93] M. A. Davenport, M. F. Duarte, Y. C. Eldar, and G. Kutyniok. Introduction to Compressed Sensing. Preprint 93, DFG-SPP 1324, May 2011.
- [94] H.-C. Kreuzler and H. Yserentant. The Mixed Regularity of Electronic Wave Functions in Fractional Order and Weighted Sobolev Spaces. Preprint 94, DFG-SPP 1324, June 2011.
- [95] E. Ullmann, H. C. Elman, and O. G. Ernst. Efficient Iterative Solvers for Stochastic Galerkin Discretizations of Log-Transformed Random Diffusion Problems. Preprint 95, DFG-SPP 1324, June 2011.
- [96] S. Kunis and I. Melzer. On the Butterfly Sparse Fourier Transform. Preprint 96, DFG-SPP 1324, June 2011.
- [97] T. Rohwedder. The Continuous Coupled Cluster Formulation for the Electronic Schrödinger Equation. Preprint 97, DFG-SPP 1324, June 2011.
- [98] T. Rohwedder and R. Schneider. Error Estimates for the Coupled Cluster Method. Preprint 98, DFG-SPP 1324, June 2011.
- [99] P. A. Cioica and S. Dahlke. Spatial Besov Regularity for Semilinear Stochastic Partial Differential Equations on Bounded Lipschitz Domains. Preprint 99, DFG-SPP 1324, July 2011.
- [100] L. Grasedyck and W. Hackbusch. An Introduction to Hierarchical (H-) Rank and TT-Rank of Tensors with Examples. Preprint 100, DFG-SPP 1324, August 2011.
- [101] N. Chegini, S. Dahlke, U. Friedrich, and R. Stevenson. Piecewise Tensor Product Wavelet Bases by Extensions and Approximation Rates. Preprint 101, DFG-SPP 1324, September 2011.
- [102] S. Dahlke, P. Oswald, and T. Raasch. A Note on Quarkonial Systems and Multi-level Partition of Unity Methods. Preprint 102, DFG-SPP 1324, September 2011.
- [103] A. Uschmajew. Local Convergence of the Alternating Least Squares Algorithm For Canonical Tensor Approximation. Preprint 103, DFG-SPP 1324, September 2011.
- [104] S. Kvaal. Multiconfigurational time-dependent Hartree method for describing particle loss due to absorbing boundary conditions. Preprint 104, DFG-SPP 1324, September 2011.

- [105] M. Guillemard and A. Iske. On Groupoid C^* -Algebras, Persistent Homology and Time-Frequency Analysis. Preprint 105, DFG-SPP 1324, September 2011.
- [106] A. Hinrichs, E. Novak, and H. Woźniakowski. Discontinuous information in the worst case and randomized settings. Preprint 106, DFG-SPP 1324, September 2011.