

DFG-Schwerpunktprogramm 1324

„Extraktion quantifizierbarer Information aus komplexen Systemen“

On the convergence of the stochastic Galerkin method for random elliptic partial differential equations

A. Mugler, H.-J. Starkloff

Preprint 123



Edited by

AG Numerik/Optimierung
Fachbereich 12 - Mathematik und Informatik
Philipps-Universität Marburg
Hans-Meerwein-Str.
35032 Marburg


DFG-Schwerpunktprogramm 1324

„Extraktion quantifizierbarer Information aus komplexen Systemen“

On the convergence of the stochastic Galerkin method for random elliptic partial differential equations

A. Mugler, H.-J. Starkloff

Preprint 123



The consecutive numbering of the publications is determined by their chronological order.

The aim of this preprint series is to make new research rapidly available for scientific discussion. Therefore, the responsibility for the contents is solely due to the authors. The publications will be distributed by the authors.

On the convergence of the stochastic Galerkin method for random elliptic partial differential equations*

Antje Mugler[†] and Hans-Jörg Starkloff[‡]

June 12, 2012

Abstract

In this article we consider elliptic partial differential equations with random coefficients and/or random forcing terms. In the current treatment of such problems by stochastic Galerkin methods it is standard to assume that the random diffusion coefficient is bounded by positive deterministic constants. In contrast, we make the significantly weaker assumption that the non-negative random coefficients can be bounded strictly away from zero and infinity by random variables only. We show that in this case the standard stochastic Galerkin approach does not necessarily produce a sequence of approximate solutions that converges in the natural norm to the exact solution. By using weighted test function spaces we develop an alternative stochastic Galerkin approach and prove that the associated sequence of approximate solutions converges to the exact solution in the natural norm.

Key words. equations with random data, stochastic Galerkin method, generalized polynomial chaos, spectral methods

1 Introduction

The stochastic Galerkin method developed by Ghanem and Spanos in [21] and refined by many researchers (see e. g. the books of LeMaître and Knio [26] and Xiu [38], the Acta Numerica article of Schwab and Gittelson [33] and the references therein) is a powerful tool for the numerical solution of different types of random equations. In many applications like e. g. subsurface flows in porous media input parameters cannot be modeled as deterministic functions but depend also on a random component. In this setting, it is desirable to quantify the uncertainty of the modeled system. To this end a variant of the stochastic Galerkin method, based on Wiener-Hermite or generalized polynomial chaos expansions, is mainly used for computing the approximate solution of the underlying random equation. The concept of polynomial chaos was introduced by N. Wiener [37] and the concept of generalized polynomial chaos expansions was introduced by Xiu and Karniadakis in [40, 36]. Some fundamental properties of such expansions have been investigated in [13, 14].

In this work we focus on elliptic partial differential equations of the form

$$-\nabla \cdot (\kappa \nabla u) = f$$

*This work was supported by the Deutsche Forschungsgemeinschaft Priority Programme 1324.

[†]Address: Mathematisches Institut, Brandenburg University of Technology, 03013 Cottbus, Germany.

e-mail: mugler@math.tu-cottbus.de

[‡]Address: Institut für Stochastik, TU Bergakademie Freiberg, 09596 Freiberg, Germany.

e-mail: hjstark@math.tu-freiberg.de

with random coefficient κ and random force term f together with appropriate boundary conditions. If the random coefficient κ can be bounded strictly away from zero and infinity by deterministic constants, i. e., there are constants $\underline{\kappa}, \bar{\kappa} > 0$, such that

$$0 < \underline{\kappa} \leq \kappa(x, \omega) \leq \bar{\kappa} < \infty \quad \text{a. e. and a. s.},$$

then using the Lax-Milgram theorem it can be shown, that there exists a unique solution of a corresponding stochastic variational problem (see e. g. Babuška et al. [11, 2, 3, 4] or Schwab et al. [15, 5, 6, 34, 10]). In addition, it can be shown that the standard stochastic Galerkin approach described in these works yields approximate solutions which converge to the exact solution in the natural norm.

However, in some applications where the coefficient describes a physical property which has to be positive (e. g. the permeability of a medium) the random coefficient is modeled as a lognormal random field (see e. g. [20, 19, 16, 44, 32]) which cannot be bounded by deterministic constants. In contrast, such random fields are strictly bounded away from zero and infinity by *random variables*. In this case we cannot use the Lax-Milgram theorem directly to prove the existence of a unique weak solution. In recent works Galvis and Sarkis [17] and Gittelsohn [22] prove the existence result using alternative techniques. However, the question of convergence of corresponding standard stochastic Galerkin approximations to the exact solution is not addressed.

In this work we generalize the results of Galvis and Sarkis and Gittelsohn, respectively, and consider more general random coefficients κ , for which there are random variables $\kappa_{min}, \kappa_{max} > 0$ a. s. satisfying

$$0 < \kappa_{min}(\omega) \leq \kappa(x, \omega) \leq \kappa_{max}(\omega) < \infty \quad \text{a. e. and a. s.}$$

Thus, we allow coefficients with distributions different from a lognormal one and do not use special properties of the normal distribution in the proofs. We investigate the existence of a unique weak solution for elliptic problems with random coefficients of this type as well as the convergence of the standard stochastic Galerkin approximations. Furthermore we construct an example where the standard stochastic Galerkin approximations do not converge in the natural norm to the exact solution. In order to overcome this convergence problem we introduce an alternative stochastic Galerkin approach which uses a modified finite-dimensional test function space corresponding to a stochastic Petrov-Galerkin approach. We prove the convergence of the associated stochastic Petrov-Galerkin approximations to the exact solution in the natural norm.

The paper is organized as follows. In Section 2, we describe our model problem and present two different stochastic variational formulations with standard unweighted and weighted (bi)linear forms, respectively. We discuss the existence of a unique solution for each case. In Section 3, we apply the stochastic Galerkin method to both formulations. We arrive at two discrete stochastic variational problems which can be interpreted as two different stochastic Galerkin approaches for the variational formulation with unweighted (bi)linear forms. One approach corresponds to the standard stochastic Galerkin ansatz, and the other one corresponds to a stochastic Petrov-Galerkin ansatz where the test function space is weighted by a suitable random variable. For the stochastic Petrov-Galerkin method we establish also convergence results w. r. t. the natural norm. Examples, demonstrating that the standard stochastic Galerkin method is not stable and their stochastic Galerkin solutions may not converge in the natural norm, are given in Section 4. Here, the non-convergence is explicitly proved and illustrated numerically. In contrast, we show that our new stochastic Petrov-Galerkin approach yields a convergent sequence of approximate solutions.

2 Stochastic Variational Problem Formulations

Let $D \subset \mathbb{R}^d$, $d \in \mathbb{N}$, be a bounded Lipschitz domain and $(\Omega, \mathfrak{A}, \mathbf{P})$ a probability space consisting of a sample space Ω , a σ -algebra \mathfrak{A} of subsets of Ω and a probability measure \mathbf{P} defined on the σ -algebra \mathfrak{A} . We consider the following boundary value problem

$$\begin{aligned} -\nabla \cdot (\kappa(x, \omega) \nabla u(x, \omega)) &= f(x, \omega) & x \in D, \omega \in \Omega, \\ u(x, \omega) &= 0 & x \in \partial D, \omega \in \Omega, \end{aligned} \quad (2.1)$$

for a given random coefficient κ and a random forcing term f . In contrast to many articles (compare e. g. with Babuška et al. [2, 3, 4] or Schwab et al. [5, 6, 34]) we do not assume that the coefficient κ can be uniformly bounded by deterministic constants. We only assume that each realization of the coefficient κ can be strictly bounded away from zero and infinity.

Assumption 2.1 *The coefficient $\kappa : D \times \Omega \rightarrow \mathbb{R}$ is a strongly measurable random variable with values in $L^\infty(D)$ and there exist real-valued random variables κ_{\min} and κ_{\max} on $(\Omega, \mathfrak{A}, \mathbf{P})$ such that*

$$0 < \kappa_{\min}(\omega) \leq \kappa(x, \omega) \leq \kappa_{\max}(\omega) < \infty \quad a. e. \text{ and } a. s. \quad (2.2)$$

W. l. o. g. we assume that κ_{\min} and κ_{\max} are measurable with respect to the σ -algebra $\sigma(\kappa)$ which is generated by the random variable κ .

Remark 2.2 *A random variable η with values in a Banach space \mathcal{B} is called strongly measurable iff there exists a sequence of countably-valued random variables in \mathcal{B} converging almost surely in \mathcal{B} to η (see e. g. [24]). We assume strongly measurable coefficients since the space $L^\infty(D)$ is a non-separable Banach space which causes some problems for probabilistic investigations. In most applications coefficients have additional regularity, for example, coefficients are assumed to be continuous or to belong to another separable subspace of $L^\infty(D)$.*

Assumption 2.1 is satisfied for random coefficients with constant bounds as well as lognormal random fields under weak conditions but other $L^\infty(D)$ -valued random variables can be treated as well. Assumption 2.1 is not very restrictive: For any non-negative strongly measurable random variable κ in $L^\infty(D)$ where $1/\kappa$ belongs almost surely to $L^\infty(D)$, there exist random bounds $0 < \kappa_{\min} < \kappa_{\max}$ as in (2.2). Though Assumption 2.1 is fairly general the pathwise weak problem, i. e. a variational formulation associated with the spatial variables, is a well-posed problem. To see this, we define the corresponding pathwise bilinear form $b(\cdot, \cdot; \omega) : H^1(D) \times H^1(D) \rightarrow \mathbb{R}$ by

$$b(u, v; \omega) = \int_D \kappa(x, \omega) \nabla u(x) \cdot \nabla v(x) dx$$

for $\omega \in \Omega$ and we denote by $\langle g, v \rangle_{H^{-1}, \dot{H}^1}$ the duality pairing between $g \in H^{-1}(D)$ and $v \in \dot{H}^1(D)$. Now, assuming that f is a random variable with values in $H^{-1}(D)$, we consider a pathwise weak formulation of the boundary-value problem (2.1).

Problem 2.3 (Pathwise Weak Formulation) *For a given random variable f with values in $H^{-1}(D)$ find a random variable \tilde{u} with values in $\dot{H}^1(D)$, such that*

$$b(\tilde{u}(\omega), v; \omega) = \langle f(\omega), v \rangle_{H^{-1}, \dot{H}^1} \quad \text{for all } v \in \dot{H}^1(D) \quad (2.3)$$

holds almost surely.

Since almost every realization of the coefficient κ is bounded by Assumption 2.1 and f is a random variable with values in $H^{-1}(D)$, the Lax-Milgram theorem (see e.g. [8] Theorem 2.7.7) tells us that there exists a mapping $\tilde{u} : \Omega \rightarrow \mathring{H}^1(D)$, $\omega \mapsto \tilde{u}(\omega)$ which satisfies the variational equation (2.3) almost surely. Furthermore, the Lax-Milgram theorem provides the estimate

$$\|\tilde{u}(\omega)\|_{H^1(D)} \leq C \frac{\|f(\omega)\|_{H^{-1}(D)}}{\kappa_{\min}(\omega)} \quad \text{a. s.} \quad (2.4)$$

where $C > 0$ is a suitable constant independent of $\omega \in \Omega$. Moreover, due to Assumption 2.1 this mapping \tilde{u} can be chosen as a random variable which is measurable w. r. t. the σ -algebra $\sigma(\kappa, f)$ generated by the random variables κ and f . Thus, the mapping \tilde{u} is the unique solution of the pathwise weak formulation, i. e. of Problem 2.3. For further details we refer to [14, 30].

In analogy to purely deterministic boundary-value problems we wish to formulate a variational problem in both spatial and random variables associated with (2.1). This requires a suitable Hilbert space of random variables with values in the Sobolev space $\mathring{H}^1(D)$. Now, moments of random variables with values in appropriate functional spaces come into play. As it turns out it is not sufficient to deal with expectations with respect to the basic probability measure \mathbf{P} , but suitable weightings are required. Given a real-valued measurable random function $\rho : D \times \Omega \rightarrow \mathbb{R}$ with $\rho(\cdot, \cdot) > 0$ a. e. and a. s. which satisfies Assumption 2.1 we introduce the weighted space

$$\mathring{U}_\rho := \left\{ \eta : \Omega \rightarrow \mathring{H}^1(D) \text{ measurable} : \|\eta\|_{\mathring{U}_\rho} < \infty \right\}$$

with

$$\|\eta\|_{\mathring{U}_\rho}^2 := \mathbf{E}_{\mathbf{P}} \left(\int_D |\nabla \eta(x)|^2 \rho(x) dx \right) = \int_\Omega \int_D |\nabla \eta(x, \omega)|^2 \rho(x, \omega) dx d\mathbf{P}(\omega)$$

which is a Hilbert space together with the inner product

$$(\eta_1, \eta_2)_{\mathring{U}_\rho} := \mathbf{E}_{\mathbf{P}} \left(\int_D \nabla \eta_1(x) \cdot \nabla \eta_2(x) \rho(x) dx \right).$$

If the weight function ρ is independent of the spatial argument, i. e., if it is an almost surely positive real-valued random variable $\rho > 0$, then the weighted spaces

$$\begin{aligned} U_\rho^m &:= L^2(\Omega, \mathfrak{A}, \rho d\mathbf{P}; H^m(D)) \\ &:= \left\{ \eta : \Omega \rightarrow H^m(D) \text{ measurable} : \mathbf{E}_{\mathbf{P}} \left(\|\eta\|_{H^m(D)}^2 \rho \right) < \infty \right\}, \quad m \in \mathbb{Z}, \end{aligned}$$

and

$$\begin{aligned} \mathring{U}_\rho^m &:= L^2(\Omega, \mathfrak{A}, \rho d\mathbf{P}; \mathring{H}^m(D)) \\ &:= \left\{ \eta : \Omega \rightarrow \mathring{H}^m(D) \text{ measurable} : \mathbf{E}_{\mathbf{P}} \left(\|\eta\|_{\mathring{H}^m(D)}^2 \rho \right) < \infty \right\}, \quad m \in \mathbb{N}_0 \end{aligned}$$

equipped with the norm $\|\eta\|_{\mathring{U}_\rho^m}^2 := \mathbf{E}_{\mathbf{P}} \left(\|\eta\|_{\mathring{H}^m(D)}^2 \rho \right) = \mathbf{E}_{\mathbf{P}} \left(\left(\sum_{|\alpha| \leq m} \|\partial^\alpha \eta\|_{L^2(D)}^2 \right) \rho \right)$ and the inner product $(\eta_1, \eta_2)_{\mathring{U}_\rho^m} := \mathbf{E}_{\mathbf{P}} \left((\eta_1, \eta_2)_{\mathring{H}^m(D)} \rho \right)$ are also Hilbert spaces. Note that for $\eta \in \mathring{U}_\rho^m$, i. e. for random variables η with values in $\mathring{H}^1(D)$, the seminorm

$$|\eta|_{U_\rho^m} := \mathbf{E}_{\mathbf{P}} \left(|\eta|_{\mathring{H}^m(D)}^2 \rho \right) = \mathbf{E}_{\mathbf{P}} \left(\rho \int_D |\nabla \eta(x)|^2 dx \right)$$

is equivalent to the norm $\|\cdot\|_{U_\varrho^m}$. If the space $L^2(\Omega, \mathfrak{A}, \varrho d\mathbf{P}) := \{\eta : \Omega \rightarrow \mathbb{R} \text{ measurable} : \mathbf{E}_{\mathbf{P}}(|\eta|^2 \varrho) < \infty\}$ is separable, then there exist isomorphisms to the corresponding tensor product spaces, i. e.

$$U_\varrho^m \cong H^m(D) \otimes L^2(\Omega, \mathfrak{A}, \varrho d\mathbf{P}), \quad m \in \mathbb{Z},$$

and

$$\mathring{U}_\varrho^m \cong \mathring{H}^m(D) \otimes L^2(\Omega, \mathfrak{A}, \varrho d\mathbf{P}), \quad m \in \mathbb{N}_0.$$

Furthermore, the dual space of \mathring{U}_ϱ^m can be identified with the space $U_{\varrho^{-1}}^{-m}$. For convenience we denote the unweighted spaces, i. e. $\varrho \equiv 1$, by U^m and \mathring{U}^m . On occasion we may replace the probability measure \mathbf{P} in the definition of U^m and \mathring{U}^m by another probability measure \mathbf{Q} and write $U_{\mathbf{Q}}^m$ and $\mathring{U}_{\mathbf{Q}}^m$, respectively, in this case.

We define the bilinear form

$$a(u, v) := \mathbf{E}_{\mathbf{P}} \left(\int_D \kappa(x) \nabla u(x) \cdot \nabla v(x) dx \right) = \int_\Omega \int_D \kappa(x, \omega) \nabla u(x, \omega) \cdot \nabla v(x, \omega) dx d\mathbf{P}(\omega),$$

and the linear form

$$\ell(v) := \mathbf{E}_{\mathbf{P}} \left(\langle f, v \rangle_{H^{-1}, \mathring{H}^1} \right) = \int_\Omega \langle f(\omega), v(\omega) \rangle_{H^{-1}, \mathring{H}^1} d\mathbf{P}(\omega).$$

Given Assumption 2.1 the weak problem posed in unweighted spaces, i. e., for a given $f \in U^{-1}$ find $u \in \mathring{U}^1$ such that there holds

$$a(u, v) = \ell(v) \quad \text{for all } v \in \mathring{U}^1 \quad (2.5)$$

is in general *not* well-posed. For example, it may happen that there does not even exist a solution in \mathring{U}^1 for a given $f \in U^{-1}$ (see [30]). Note, that this stands in marked contrast to the case when the coefficient κ can be strictly bounded away from zero and infinity. Thus, we have to use an alternative stochastic variational formulation and arrive at a first weak problem posed on the associated energy space as in [2, 22].

Problem 2.4 (Unweighted Weak Formulation) *For a given f find $\hat{u} \in \mathring{U}_\kappa$, such that*

$$a(\hat{u}, v) = \ell(v) \quad \text{for all } v \in \mathring{U}_\kappa.$$

Thereby the term *unweighted* weak formulation refers to the fact that standard *unweighted* (bi)linear forms are used although the solution and test function spaces are weighted spaces. Afterwards we introduce also a *weighted* weak formulation, see Problem 2.6, where the (bi)linear forms are weighted.

To prove existence of a solution to Problem 2.4 we employ weighted spaces with the lower bound κ_{min} in Assumption 2.1. We note that such a lower bound is not uniquely defined and some freedom in its choice is sometimes useful. Due to (2.2) there exist several real-valued random variables $\mu > 0$ a. s. satisfying

$$0 < c\mu(\omega) \leq \kappa(x, \omega) \quad \text{a. e. and a. s.} \quad (2.6)$$

for a constant $c > 0$. The random variable κ_{min} is only a special case satisfying (2.6) with constant $c = 1$. W. l. o. g. we can assume that the random variable μ is also $\sigma(\kappa)$ -measurable.

Theorem 2.5 *If Assumption 2.1 holds and μ satisfies (2.6), then for any $f \in U_{\mu^{-1}}^{-1}$ there exists a unique $\hat{u} \in \mathring{U}_\kappa$ solving Problem 2.4.*

Proof.

Since for $f \in U_{\mu^{-1}}^{-1}$ we have $\ell \in U_{\mu^{-1}}^{-1} \subset (\mathring{U}_\kappa)^*$, the result follows immediately from Riesz' representation theorem. \square

Therefore Problem 2.4 is well-posed in the given weighted spaces. However, we are interested in a solution \hat{u} in the space \mathring{U}^1 rather than the space \mathring{U}_κ . For deterministic lower bounds the space \mathring{U}_κ is continuously embedded in \mathring{U}^1 . Unfortunately, this is in general not true if the coefficient κ can be bounded from below only by a random variable, e. g. μ . Clearly, if we assume a forcing term $f \in U_{\mu^{-2}}^{-1}$, then our discussion of the pathwise variational problem and the corresponding estimate (2.4) above tell us that the weak solution \hat{u} belongs also to the space \mathring{U}^1 . In general, however, we cannot estimate the \mathring{U}^1 -norm in terms of the \mathring{U}_κ -norm.

A possible remedy is to recast Problem 2.4 above. By multiplying the pathwise variational equation (2.3) with the reciprocal of the random variable μ we obtain a random coefficient κ/μ which can now be strictly bounded away from zero by the constant c in inequality (2.6). The corresponding bilinear form a_μ and linear form ℓ_μ are the weighted versions of the bilinear form a and linear form ℓ , respectively, i. e.

$$a_\mu(u, v) := \mathbf{E}_\mathbf{P} \left(\frac{1}{\mu} \int_D \kappa(x) \nabla u(x) \cdot \nabla v(x) dx \right) = \int_\Omega \int_D \frac{\kappa(x, \omega)}{\mu(\omega)} \nabla u(x, \omega) \cdot \nabla v(x, \omega) dx d\mathbf{P}(\omega)$$

and

$$\ell_\mu(v) := \mathbf{E}_\mathbf{P} \left(\frac{1}{\mu} \langle f, v \rangle_{H^{-1}, \mathring{H}^1} \right) = \int_\Omega \frac{1}{\mu(\omega)} \langle f(\omega), v(\omega) \rangle_{H^{-1}, \mathring{H}^1} d\mathbf{P}(\omega).$$

Problem 2.6 (Weighted Weak Formulation) For a given f find $\bar{u} \in \mathring{U}_\mu^\kappa$, such that there holds

$$a_\mu(\bar{u}, v) = \ell_\mu(v) \quad \text{for all } v \in \mathring{U}_\mu^\kappa.$$

Theorem 2.7 If Assumption 2.1 holds and μ satisfies (2.6), then for any $f \in U_{\mu^{-2}}^{-1}$ there exists a unique $\bar{u} \in \mathring{U}_\mu^\kappa$ solving Problem 2.6.

Proof.

Since for any $f \in U_{\mu^{-2}}^{-1}$ there holds $\ell_\mu \in U^{-1} \subset (\mathring{U}_\mu^\kappa)^*$ the result follows immediately from Riesz' representation theorem. \square

Now, the space \mathring{U}_μ^κ is continuously embedded in \mathring{U}^1 . Thus, there exists a constant $C > 0$ such that

$$\|u\|_{U^1} \leq C \|u\|_{\mathring{U}_\mu^\kappa} \quad \text{for all } u \in \mathring{U}_\mu^\kappa.$$

Corollary 2.8 Under Assumption 2.1 together with (2.6) for every $f \in U_{\mu^{-1}}^{-1} \cap U_{\mu^{-2}}^{-1}$ the weak formulations Problem 2.4 and Problem 2.6 have unique solutions \hat{u} and \bar{u} in \mathring{U}^1 , respectively, which are almost surely equal to the pathwise solution \tilde{u} of the pathwise weak formulation Problem 2.3, i. e.,

$$\hat{u} = \bar{u} = \tilde{u} \quad \text{a. s.}$$

as random variables in $\mathring{H}^1(D)$. Moreover, both weak solutions \hat{u} and \bar{u} are $\sigma(\kappa, f)$ -measurable.

Proof.

Analogously to Theorem 3.3 in [30], we can show that both weak solutions \hat{u} and \bar{u} solve the pathwise weak formulation Problem 2.3 almost surely, the solution \tilde{u} of which is unique and $\sigma(\kappa, f)$ -measurable. \square

Therefore, we assume throughout the rest of the paper that $f \in U_{\mu^{-1}}^{-1} \cap U_{\mu^{-2}}^{-1}$ and do not distinguish between the three weak solutions, but refer to *the* weak solution denoted by \hat{u} .

Remark 2.9 *The results in this article hold also for other boundary conditions as long as the bilinear forms a and a_μ form inner products on the corresponding solution spaces.*

3 Stochastic Galerkin Method

The stochastic Galerkin method or stochastic finite element method (SFEM) is a very popular approach for the numerical solution of random equations. It is based on a stochastic variational formulation and uses sequences of finite-dimensional solution and test function spaces in order to obtain approximate solutions. However, to get reliable results a rigorous convergence analysis must be carried out. In the next section we will construct examples showing that the standard stochastic Galerkin approach applied to our model problem (2.1) is not stable and may not lead to a sequence of approximate solutions which converge in the natural norm to the exact solution. Therefore, we present in this section an alternative stochastic Galerkin approach and prove its stability.

The definition of the finite-dimensional solution and test function spaces requires a discretization in the spatial as well as in the stochastic dimension. Therefore, we parameterize the random input data by finitely many or countably many real-valued random variables which are referred to as basic random variables. For example, the popular Karhunen-Loève expansion of random fields (see e.g. [27]) yields such a representation. As observed, for example, by Karniadakis, Xiu and their co-authors in [28, 39, 40, 41, 42, 43] the choice/distribution of the basic random variables affects the approximation quality of the stochastic Galerkin solution. We mention that not only the distribution of the basic random variables is crucial here.

In the following we assume that for our model problem (2.1) a finite or infinite-dimensional random vector $\xi := (\xi_i)_{i \in I_\xi}$ with values in $\mathbb{R}^{|I_\xi|}$ is given such that κ and f are both measurable with respect to $\sigma(\xi)$. Hence, due to the Doob-Dynkin lemma (e.g. in [25]) there exist measurable functions $\kappa_\xi : \mathbb{R}^{|I_\xi|} \rightarrow L^\infty(D)$ and $f_\xi : \mathbb{R}^{|I_\xi|} \rightarrow H^{-1}(D)$ satisfying $\kappa = \kappa_\xi(\xi)$ in $L^\infty(D)$ and $f = f_\xi(\xi)$ in $H^{-1}(D)$ almost surely. Because of the assumptions on f we obtain a weak solution \hat{u} in \mathring{U}^1 . By Corollary 2.8 the weak solution \hat{u} is $\sigma(\kappa, f)$ -measurable and thus it is also $\sigma(\xi)$ -measurable. Consequently there exists a measurable function $\hat{u}_\xi : \mathbb{R}^{|I_\xi|} \rightarrow \mathring{H}^1(D)$, such that $\hat{u} = \hat{u}_\xi(\xi)$ a.s. in $\mathring{H}^1(D)$. In our setting all random variables of interest are $\sigma(\xi)$ -measurable, thus, we suppose w.l.o.g. that $\mathfrak{A} = \sigma(\xi)$ holds. We then identify the space $U^1 = L^2(\Omega, \sigma(\xi), \mathbf{P}; \mathring{H}^1(D))$ as appropriate solution space.

Moreover, we assume that the random vector ξ has the distribution $F_\xi^{\mathbf{P}}$ and each of the basic random variables ξ_i , $i \in I_\xi$, possesses finite moments of arbitrary order, i.e. $\mathbf{E}_{\mathbf{P}}|\xi_i|^n < \infty$ for all $n \in \mathbb{N}$, and has a continuous distribution function $F_{\xi_i}^{\mathbf{P}}$. Here, the distributions $F_\xi^{\mathbf{P}}$ and $F_{\xi_i}^{\mathbf{P}}$ are understood as the distributions of the random variables ξ and ξ_i w.r.t. the probability measure \mathbf{P} , i.e., these are the distributions of ξ and ξ_i as random variables (with values in the spaces $\mathbb{R}^{|I_\xi|}$ and \mathbb{R} , respectively) on the probability space $(\Omega, \mathfrak{A}, \mathbf{P})$. On occasion we will consider these random variables also on probability spaces with other probability measures and thus their distributions change accordingly. For this reason we attach the corresponding probability measure to the distribution symbol of a random variable.

We choose the finite-dimensional solution space

$$U_{N,K,hp} := U_{hp} \otimes U_{N,K} \subset \dot{H}^1(D) \otimes L^2(\Omega, \sigma(\xi), \mathbf{P}) \simeq L^2(\Omega, \sigma(\xi), \mathbf{P}; \dot{H}^1(D)),$$

where U_{hp} is a finite-dimensional subspace of $\dot{H}^1(D)$ obtained by a suitable version (h -, p - or hp -) of the deterministic finite element method and $U_{N,K}$ is a finite-dimensional subspace of $L^2(\Omega, \sigma(\xi_1, \dots, \xi_K), \mathbf{P}) \subset L^2(\Omega, \sigma(\xi), \mathbf{P})$ where $\{1, \dots, K\} \subseteq I_\xi$. Since we want to use generalized polynomial chaos (see e.g. [39, 40]), i.e. polynomials in the underlying basic random variables ξ , we construct the finite dimensional space $U_{N,K}$ as follows

$$U_{N,K} := \text{span} \left\{ \xi^\alpha := \prod_{i \in I_\xi} \xi_i^{\alpha_i}, \alpha \in \Lambda_{N,K} \right\},$$

where $\Lambda_{N,K}$ denotes a finite index set

$$\Lambda_{N,K} \subset \Lambda := \left\{ \alpha \in \mathbb{N}_0^{|I_\xi|} : \alpha \text{ has only finitely many non-zero entries} \right\}.$$

Specifically, we choose either $\Lambda_{N,K} = J_1$, where

$$J_1 := \left\{ \alpha \in \Lambda : \alpha_i = 0 \forall i > K, |\alpha| \leq N \right\}, \quad |\alpha| := \sum_{i \in I_\xi} \alpha_i,$$

selects (complete) multivariate polynomials with bounded total degree, or $\Lambda_{N,K} = J_2$, where

$$J_2 := \left\{ \alpha \in \Lambda : \alpha_i = 0 \forall i > K, \alpha_i \leq N \forall 1 \leq i \leq K \right\}$$

selects (tensor product) multivariate polynomials with uniformly bounded degree of the individual basic random variables. Then, the discrete versions of the unweighted and the weighted weak formulation introduced in Section 2 read as follows.

Problem 3.1 (Discrete Unweighted Weak Formulation) *For a given f find $\hat{u}_{N,K,hp} \in U_{N,K,hp}$, such that there holds*

$$a(\hat{u}_{N,K,hp}, v) = \ell(v) \quad \text{for all } v \in U_{N,K,hp}.$$

Problem 3.2 (Discrete Weighted Weak Formulation) *For a given f find $\bar{u}_{N,K,hp} \in U_{N,K,hp}$, such that there holds*

$$a_\mu(\bar{u}_{N,K,hp}, v) = \ell_\mu(v) \quad \text{for all } v \in U_{N,K,hp}.$$

So Problem 3.1 is exactly that discrete weak problem which arises if we apply the stochastic Galerkin method to the weak problem posed on the unweighted spaces (2.5) no matter if this is well-posed or not. Alternatively, this means, Problem 3.1 corresponds to the discrete weak problem which results from applying the stochastic Galerkin method to the standard weak problem arising in the case of a coefficient κ which can be bounded by constants. Therefore, we call the approach yielding the discrete unweighted weak problem, i.e. Problem 3.1, the *standard* stochastic Galerkin approach. Thus, the solution $\hat{u}_{N,K,hp}$ is called the *standard* stochastic Galerkin solution or *standard* stochastic Galerkin approximation, respectively.

We then make the following observations.

Theorem 3.3 *Let Assumption 2.1 be fulfilled and $f \in U_{\mu^{-2}}^{-1} \cap U_{\mu^{-1}}^{-1}$.*

(i) If $\kappa_{max} \in L^r(\Omega, \mathfrak{A}, \mathbf{P})$ for some $r > 1$, then there exists a unique stochastic Galerkin solution $\hat{u}_{N,K,hp}$ in $U_{N,K,hp}$ solving Problem 3.1. Moreover, $\hat{u}_{N,K,hp}$ is the best approximation in $U_{N,K,hp}$ of the weak solution \hat{u} with respect to the norm of \dot{U}_κ , i. e. $\|\hat{u} - \hat{u}_{N,K,hp}\|_{\dot{U}_\kappa} = \inf_{z \in U_{N,K,hp}} \|\hat{u} - z\|_{\dot{U}_\kappa}$.

(ii) If $\kappa_{max}/\mu \in L^r(\Omega, \mathfrak{A}, \mathbf{P})$ for some $r > 1$, then there exists a unique stochastic Galerkin solution $\bar{u}_{N,K,hp}$ in $U_{N,K,hp}$ solving Problem 3.2. Moreover, $\bar{u}_{N,K,hp}$ is the best approximation in $U_{N,K,hp}$ of the weak solution \hat{u} with respect to the norm of $\dot{U}_{\frac{\kappa}{\mu}}$, i. e. $\|\hat{u} - \bar{u}_{N,K,hp}\|_{\dot{U}_{\frac{\kappa}{\mu}}} = \inf_{z \in U_{N,K,hp}} \|\hat{u} - z\|_{\dot{U}_{\frac{\kappa}{\mu}}}$.

Proof.

In order to prove this result we show that the finite-dimensional subspace $U_{N,K,hp}$ is a subset of \dot{U}_κ and $\dot{U}_{\frac{\kappa}{\mu}}$, respectively. For any $v = \xi^\alpha \varphi$, $\alpha \in \Lambda$ and $\varphi \in U_{hp}$ this can be deduced for (i) from the integrability condition of κ_{max} by

$$\begin{aligned} \|v\|_{\dot{U}_\kappa}^2 &= a(v, v) = \int_{\Omega} \int_D \kappa(x, \omega) |\nabla v(x, \omega)|^2 dx d\mathbf{P}(\omega) \\ &\leq \int_{\Omega} \int_D \kappa_{max}(\omega) \xi(\omega)^{2\alpha} |\nabla \varphi(x)|^2 dx d\mathbf{P}(\omega) \\ &= |\varphi|_{H^1(D)}^2 \mathbf{E}_{\mathbf{P}} \left(\kappa_{max} \xi^{2\alpha} \right) \leq |\varphi|_{H^1(D)}^2 (\mathbf{E}_{\mathbf{P}} \kappa_{max}^r)^{1/r} (\mathbf{E}_{\mathbf{P}} \xi^{2s\alpha})^{1/s} < \infty \end{aligned}$$

and for (ii) from the integrability condition on κ_{max}/μ by

$$\begin{aligned} \|v\|_{\dot{U}_{\frac{\kappa}{\mu}}}^2 &= a_\mu(v, v) = \int_{\Omega} \int_D \frac{\kappa(x, \omega)}{\mu(\omega)} |\nabla v(x, \omega)|^2 dx d\mathbf{P}(\omega) \\ &\leq \int_{\Omega} \int_D \frac{\kappa_{max}}{\mu}(\omega) \xi(\omega)^{2\alpha} |\nabla \varphi(x)|^2 dx d\mathbf{P}(\omega) \\ &= |\varphi|_{H^1(D)}^2 \mathbf{E}_{\mathbf{P}} \left(\frac{\kappa_{max}}{\mu} \xi^{2\alpha} \right) \leq |\varphi|_{H^1(D)}^2 \left(\mathbf{E}_{\mathbf{P}} \left(\frac{\kappa_{max}}{\mu} \right)^r \right)^{1/r} (\mathbf{E}_{\mathbf{P}} \xi^{2s\alpha})^{1/s} < \infty, \end{aligned}$$

respectively, where $1 < s < \infty$ with $\frac{1}{r} + \frac{1}{s} = 1$ in both cases. Thus, there exist unique best approximations, i. e. orthogonal projections $\hat{u}_{N,K,hp} := \Pi_{U_{N,K,hp}} \hat{u}$ onto $U_{N,K,hp}$ in \dot{U}_κ and $\bar{u}_{N,K,hp} := \Pi_{U_{N,K,hp}} \hat{u}$ onto $U_{N,K,hp}$ in $\dot{U}_{\frac{\kappa}{\mu}}$ satisfying

$$a(\hat{u} - \hat{u}_{N,K,hp}, v) = 0 \quad \text{and} \quad a_\mu(\hat{u} - \bar{u}_{N,K,hp}, v) = 0, \quad \text{respectively,}$$

for all $v \in U_{N,K,hp}$. In other words, since $a(\hat{u}, v) = \ell(v)$ and $a_\mu(\bar{u}, v) = \ell_\mu(v)$, we have arrived at

$$a(\hat{u}_{N,K,hp}, v) = \ell(v) \quad \text{and} \quad a_\mu(\bar{u}_{N,K,hp}, v) = \ell_\mu(v), \quad \text{respectively,}$$

for all $v \in U_{N,K,hp}$. □

Notably, the discrete version of the weighted weak problem can also be interpreted as a discrete version of the unweighted variational formulation by using a so-called stochastic Petrov-Galerkin ansatz where the finite-dimensional test function space differs from the finite-dimensional solution space. If we define the weighted finite-dimensional test function space

$$V_{N,K,hp} := \left\{ \frac{u}{\mu} : u \in U_{N,K,hp} \right\},$$

we see that the problem of finding $\bar{u}_{N,K,hp} \in U_{N,K,hp}$ which satisfies

$$a_\mu(\bar{u}_{N,K,hp}, v) = \ell_\mu(v) \quad \text{for all } v \in U_{N,K,hp}$$

is equivalent to finding $\bar{u}_{N,K,hp} \in U_{N,K,hp}$ which satisfies

$$a(\bar{u}_{N,K,hp}, v) = \ell(v) \quad \text{for all } v \in V_{N,K,hp}.$$

This means, the weighting $1/\mu$ of the (bi)linear forms is transferred to the finite-dimensional test function space. Thus, we arrive at the discrete weighted weak problem using the unweighted bilinear form a and linear form ℓ (instead of a_μ and ℓ_μ) on the finite-dimensional solution space $U_{N,K,hp}$ and the finite-dimensional test function space $V_{N,K,hp}$ which equals the space $U_{N,K,hp}$ weighted by $1/\mu$.

Altogether, this implies, that we have identified two different stochastic Galerkin approaches for the very same variational equation: We are looking for a solution in $U_{N,K,hp}$ which satisfies the equation

$$a(u, v) = \ell(v)$$

for all $v \in U_{N,K,hp}$ or for all $v \in V_{N,K,hp}$, respectively. In general this defines two different Galerkin projections and hence two different solutions in the same finite-dimensional space.

If we know in addition that the finite-dimensional solution spaces $(U_{N,K,hp})_{N,K,hp}$ lie dense in the space \mathring{U}_κ or \mathring{U}_μ^κ , respectively, we can conclude that the approximation errors tend to zero, i. e.

$$\|\hat{u} - \hat{u}_{N,K,hp}\|_{\mathring{U}_\kappa} \rightarrow 0 \quad \text{and} \quad \|\hat{u} - \bar{u}_{N,K,hp}\|_{\mathring{U}_\mu^\kappa} \rightarrow 0.$$

In many cases of practical interest, however, we are not interested in the convergence in these weighted norms but in the approximation error in the unweighted, the U^1 -norm which therefore we call also the *natural norm*. Here the weighted weak formulation is advantageous due to the continuous embedding of \mathring{U}_μ^κ in \mathring{U}^1 which allows us to deduce

$$\|\hat{u} - \bar{u}_{N,K,hp}\|_{U^1} \leq C \|\hat{u} - \bar{u}_{N,K,hp}\|_{\mathring{U}_\mu^\kappa} = C \inf_{z \in U_{N,K,hp}} \|\hat{u} - z\|_{\mathring{U}_\mu^\kappa} \quad (3.1)$$

together with an upper estimate of the form

$$\|\hat{u} - \bar{u}_{N,K,hp}\|_{U^1} \leq C \inf_{z \in U_{N,K,hp}} \|\hat{u} - z\|_{U_{\frac{\kappa_{max}}{\mu}}}. \quad (3.2)$$

Hence, we can estimate the approximation error of the stochastic Galerkin projection $\bar{u}_{N,K,hp}$ in U^1 by a best approximation error in a stronger norm related to a weighting by the real-valued random variable κ_{max}/μ . Due to the assumption $\kappa_{max}/\mu \in L^r(\Omega, \mathfrak{A}, \mathbf{P})$ for some $r > 1$ the expectation $c_{\frac{\kappa_{max}}{\mu}} := \mathbf{E}_{\mathbf{P}}(\kappa_{max}/\mu)$ is finite and the measure \mathbf{Q} defined by

$$d\mathbf{Q} = \frac{1}{c_{\frac{\kappa_{max}}{\mu}}} \frac{\kappa_{max}}{\mu} d\mathbf{P}$$

is a probability measure. In the following we consider the function spaces $U_{\mathbf{Q}}^m$ and $\mathring{U}_{\mathbf{Q}}^m$ instead of $U_{\frac{\kappa_{max}}{\mu}}^m$ and $\mathring{U}_{\frac{\kappa_{max}}{\mu}}^m$, $m \in \mathbb{Z}$, which coincide as sets with $U_{\frac{\kappa_{max}}{\mu}}^m$ and $\mathring{U}_{\frac{\kappa_{max}}{\mu}}^m$ but are much easier to handle due to the corresponding probability space $(\Omega, \mathfrak{A}, \mathbf{Q})$ at hand where w. l. o. g. we have assumed $\mathfrak{A} = \sigma(\xi)$.

Corollary 3.4 *If $f \in U_{\mu^{-2}}^{-1} \cap U_{\mu^{-1}}^{-1}$, $\kappa_{max}/\mu \in L^r(\Omega, \mathfrak{A}, \mathbf{P})$ for some $r > 1$ and $\hat{u} \in \mathring{U}_{\mathbf{Q}}^1$, then there holds*

$$\|\hat{u} - \bar{u}_{N,K,hp}\|_{U^1} \leq C \inf_{z \in U_{N,K,hp}} \|\hat{u} - z\|_{U_{\mathbf{Q}}^1} \quad (3.3)$$

for a constant $C > 0$ independent of N, K, h, p and \hat{u} .

Proof.

This follows immediately from the assumptions and the estimates (3.1) and (3.2). \square

Remark 3.5 *If the coefficient κ can be bounded from below by a positive constant, then the finite-dimensional test function space $V_{N,K,hp}$ coincides with the solution space $U_{N,K,hp}$ and thus weighting is not necessary. If, additionally, the coefficient κ can also be bounded from above by a constant, then the result of Corollary 3.4 is equivalent to Céa's lemma.*

3.1 Convergence Analysis

Next, we proceed with a closer look at the estimate of the approximation error (3.3) and identify and analyze different sources of discretization errors. To this end, we introduce three different orthogonal projections and denote by

$$\Pi_{\mathring{U}_{\mathbf{Q},N,K,hp}^1} : \mathring{U}_{\mathbf{Q}}^1 \rightarrow U_{N,K,hp}$$

the orthogonal projection onto $U_{N,K,hp}$,

$$\Pi_{\mathring{U}_{\mathbf{Q},N,K}^1} : \mathring{U}_{\mathbf{Q}}^1 \rightarrow \mathring{H}^1(D) \otimes U_{N,K}$$

the orthogonal projection onto $\mathring{H}^1(D) \otimes U_{N,K}$,

$$\Pi_{\mathring{U}_{\mathbf{Q},K}^1} : \mathring{U}_{\mathbf{Q}}^1 \rightarrow \mathring{H}^1(D) \otimes L^2(\Omega, \sigma(\xi_1, \dots, \xi_K), \mathbf{Q})$$

the orthogonal projection onto

$$\mathring{H}^1(D) \otimes L^2(\Omega, \sigma(\xi_1, \dots, \xi_K), \mathbf{Q}) \simeq L^2(\Omega, \sigma(\xi_1, \dots, \xi_K), \mathbf{Q}; \mathring{H}^1(D)),$$

respectively. All projections are carried out w. r. t. the inner product associated with the $U_{\mathbf{Q}}^1$ -norm. Assuming $\hat{u} \in \mathring{U}_{\mathbf{Q}}^1$ we can estimate the approximation error of the stochastic Petrov-Galerkin approximation $\bar{u}_{N,K,hp}$ to the exact solution by using (3.3) for $z = \Pi_{\mathring{U}_{\mathbf{Q},N,K,hp}^1} \hat{u}$ as follows

$$\begin{aligned} \|\hat{u} - \bar{u}_{N,K,hp}\|_{U^1} \leq C \left[\|\hat{u} - \Pi_{\mathring{U}_{\mathbf{Q},K}^1} \hat{u}\|_{U_{\mathbf{Q}}^1} + \|\Pi_{\mathring{U}_{\mathbf{Q},K}^1} \hat{u} - \Pi_{\mathring{U}_{\mathbf{Q},N,K}^1} \hat{u}\|_{U_{\mathbf{Q}}^1} \right. \\ \left. + \|\Pi_{\mathring{U}_{\mathbf{Q},N,K}^1} \hat{u} - \Pi_{\mathring{U}_{\mathbf{Q},N,K,hp}^1} \hat{u}\|_{U_{\mathbf{Q}}^1} \right]. \end{aligned} \quad (3.4)$$

Hence this error has three components, namely an approximation error induced by discretizing in the spatial dimension

$$e_1 := \|\Pi_{\mathring{U}_{\mathbf{Q},N,K}^1} \hat{u} - \Pi_{\mathring{U}_{\mathbf{Q},N,K,hp}^1} \hat{u}\|_{U_{\mathbf{Q}}^1}$$

and two approximation errors arising from the discretization in the stochastic dimension: We consider only polynomials in the first K basic random variables

$$e_2 := \|\hat{u} - \Pi_{\mathring{U}_{\mathbf{Q},K}^1} \hat{u}\|_{U_{\mathbf{Q}}^1}$$

and bound their polynomial degrees by N

$$e_3 := \|\Pi_{\mathring{U}_{\mathbf{Q},K}^1} \hat{u} - \Pi_{\mathring{U}_{\mathbf{Q},N,K}^1} \hat{u}\|_{U_{\mathbf{Q}}^1}.$$

Again, the first error term e_1 is the error arising from the spatial discretization of the Sobolev space $\dot{H}^1(D)$ by the finite-dimensional FE space U_{hp} . Therefore, the spatial approximation error can be bounded using standard arguments from the theory of the deterministic Finite Element Methods. Here, we employ an hp -version of the FEM (see e.g. [9]). For example, under the assumptions in Section 5.8.3 in [9] and given that the domain D is obtained from the reference domain $\hat{D} := (-1, 1)^d$, $d = 1, 2, 3$, by a smooth, invertible and regular mapping $F : \hat{D} \rightarrow D$ (in the sense of Section 5.8.3 in [9]) there holds the following.

Corollary 3.6 *For $\hat{u} \in U_{\mathbf{Q}}^k \cap \dot{U}_{\mathbf{Q}}^1$, $k \geq 2$, there holds*

$$e_1 = \|\Pi_{\dot{U}_{\mathbf{Q},N,K}^1} \hat{u} - \Pi_{\dot{U}_{\mathbf{Q},N,K,hp}^1} \hat{u}\|_{U_{\mathbf{Q}}^1} \leq \tilde{C} h^{m-1} p^{-(k-1)} \|\hat{u}\|_{U_{\mathbf{Q}}^k}$$

where $m = \min\{k, p+1\}$ and the constant $\tilde{C} > 0$ is independent of N, K, h, p and \hat{u} .

Proof.

Based on the results in [9] (see e.g. equation (5.8.25)) for the deterministic case we find

$$\|\Pi_{\dot{U}_{\mathbf{Q},N,K}^1} \hat{u}(\omega) - \Pi_{\dot{U}_{\mathbf{Q},N,K,hp}^1} \hat{u}(\omega)\|_{H^1(D)} \leq C h^{m-1} p^{-(k-1)} \|\hat{u}(\omega)\|_{H^k(D)} \quad \text{a. s.}$$

with $m = \min\{k, p+1\}$ and a suitable constant $C > 0$ independent of N, K, h, p, \hat{u} and $\omega \in \Omega$. Squaring the formula above and taking the expectation $\mathbf{E}_{\mathbf{Q}}$ w. r. t. the probability measure \mathbf{Q} yields

$$\begin{aligned} \mathbf{E}_{\mathbf{Q}} \left\| \Pi_{\dot{U}_{\mathbf{Q},N,K}^1} \hat{u} - \Pi_{\dot{U}_{\mathbf{Q},N,K,hp}^1} \hat{u} \right\|_{H^1(D)}^2 &\leq C^2 h^{2(m-1)} p^{-2(k-1)} \mathbf{E}_{\mathbf{Q}} \left(\|\hat{u}\|_{H^k(D)}^2 \right) \\ &= C^2 h^{2(m-1)} p^{-2(k-1)} \|\hat{u}\|_{U_{\mathbf{Q}}^k}^2 \end{aligned}$$

with $m = \min\{k, p+1\}$ and a suitable constant $C > 0$ independent of N, K, h, p and \hat{u} . \square

Remark 3.7 *If, in addition, the mapping F is affine we can prove, analogously to [9], the result*

$$e_1 = \|\Pi_{\dot{U}_{\mathbf{Q},N,K}^1} \hat{u} - \Pi_{\dot{U}_{\mathbf{Q},N,K,hp}^1} \hat{u}\|_{U_{\mathbf{Q}}^1} \leq C h^{m-1} p^{-(k-1)} \left(\mathbf{E}_{\mathbf{Q}} |\hat{u}|_{H^{m;k}(D)}^2 \right)^{1/2}$$

with $m = \min\{k, p+1\}$, a constant $C > 0$ and where $|\cdot|_{H^{m;k}(D)}$ denotes the seminorm

$$|u|_{H^{m;k}(D)} = \left(\sum_{j=m}^k \sum_{i=1}^d \left\| \frac{\partial^j}{\partial x_i^j} u \right\|_{L^2(D)}^2 \right)^{1/2}.$$

Clearly, additional smoothness of the weak solution in the spatial argument implies faster convergence. Note that the approximation error e_1 does not depend on the basic random variables.

The second error term usually occurs only if countably many basic random variables are used to represent the random input data. For finitely many basic random variables $\xi = (\xi_i)_{i \in I_{\xi}}$, i.e. $I_{\xi} = \{1, \dots, M\} \subset \mathbb{N}$, the finite-dimensional solution space $U_{N,K,hp}$ is usually defined for $K = M$. Thus, the orthogonal projection of the weak solution \hat{u} onto the space $\dot{H}^1(D) \otimes L^2(\Omega, \sigma(\xi_1, \dots, \xi_K), \mathbf{Q})$ is the weak solution itself and the error e_2 equals zero. For a countable set of basic random variables it can be shown that the error e_2 converges to zero.

Corollary 3.8 *Let $I_{\xi} = \mathbb{N}$, then there holds for $\hat{u} \in \dot{U}_{\mathbf{Q}}^1$*

$$e_2 = \|\hat{u} - \Pi_{\dot{U}_{\mathbf{Q},K}^1} \hat{u}\|_{U_{\mathbf{Q}}^1} \rightarrow 0, \quad K \rightarrow \infty.$$

Proof.

The orthogonal projection $\Pi_{\dot{U}_{\mathbf{Q},K}^1} v$ onto $L^2(\Omega, \sigma(\xi_1, \dots, \xi_K), \mathbf{Q}; \dot{H}^1(D))$ is equal to the conditional expectation of $v \in \dot{U}_{\mathbf{Q}}^1$ w. r. t. the σ -algebra spanned by the random variables ξ_1, \dots, ξ_K (see e. g. [35]). This means

$$\Pi_{\dot{U}_{\mathbf{Q},K}^1} v = \mathbf{E}_{\mathbf{Q}} \left(v \middle| (\xi_1, \dots, \xi_K) \right).$$

Since

$$\sigma \left(\bigcup_{K \geq 1} \sigma(\xi_1, \dots, \xi_K) \right) = \sigma(\xi),$$

it follows (see e. g. [7])

$$\left\| v - \Pi_{\dot{U}_{\mathbf{Q},K}^1} v \right\|_{U_{\mathbf{Q}}^1} = \left\| v - \mathbf{E}_{\mathbf{Q}} \left(v \middle| (\xi_1, \dots, \xi_K) \right) \right\|_{U_{\mathbf{Q}}^1} \rightarrow 0, \quad K \rightarrow \infty$$

for any $v \in L^2(\Omega, \sigma(\xi), \mathbf{Q}; \dot{H}^1(D))$. Due to $\hat{u} \in L^2(\Omega, \sigma(\xi), \mathbf{Q}; \dot{H}^1(D))$ this implies immediately

$$\left\| \hat{u} - \Pi_{\dot{U}_{\mathbf{Q},K}^1} \hat{u} \right\|_{U_{\mathbf{Q}}^1} \rightarrow 0, \quad K \rightarrow \infty$$

which completes the proof. \square

Investigations of the third error term e_3 are based on properties of generalized polynomial chaos. In order to represent an arbitrary $\dot{H}^1(D)$ -valued random variable in $\dot{U}_{\mathbf{Q}}^1 = L^2(\Omega, \sigma(\xi), \mathbf{Q}; \dot{H}^1(D))$ in terms of a generalized polynomial chaos expansion the multivariate polynomials in the basic random variables ξ must be complete in $L^2(\Omega, \sigma(\xi), \mathbf{Q})$. Since $\kappa_{max}/\mu \in L^r(\Omega, \mathfrak{A}, \mathbf{P})$ for some $r > 1$ there holds $\mathbf{E}_{\mathbf{Q}} |\xi_i|^n < \infty$ for all $n \in \mathbb{N}$ and $i \in I_{\xi}$. Thus, there exists a set of multivariate orthonormal polynomials $\{p_{\alpha}(\xi), \alpha \in \Lambda\}$ w. r. t. the probability measure \mathbf{Q} , i. e.

$$\mathbf{E}_{\mathbf{Q}} (p_{\alpha}(\xi) p_{\beta}(\xi)) = \delta_{\alpha, \beta}, \quad \text{for } \alpha, \beta \in \Lambda.$$

Here, the distribution $F_{\xi}^{\mathbf{Q}}$ of the basic random variable $\xi = (\xi_i)_{i \in I_{\xi}}$ as random variable on the probability space $(\Omega, \mathfrak{A}, \mathbf{Q})$ is defined by

$$F_{\xi}^{\mathbf{Q}}(y) = \mathbf{Q}(\xi \leq y) = \mathbf{E}_{\mathbf{Q}} \mathbb{1}_{\{\xi \leq y\}} = \frac{1}{c \frac{\kappa_{max}}{\mu}} \mathbf{E}_{\mathbf{P}} \left(\mathbb{1}_{\{\xi \leq y\}} \frac{\kappa_{max}}{\mu} \right)$$

Since $\frac{\kappa_{max}}{\mu}$ is $\sigma(\xi)$ -measurable, there exists a measurable function $t_{\frac{\kappa_{max}}{\mu}} : \mathbb{R}^{|I_{\xi}|} \rightarrow \mathbb{R}$ with $t_{\frac{\kappa_{max}}{\mu}}(\xi) = \frac{\kappa_{max}}{\mu}$. Therefore, the distribution $F_{\xi}^{\mathbf{Q}}$ of ξ is determined by

$$F_{\xi}^{\mathbf{Q}}(dy) = \frac{1}{c \frac{\kappa_{max}}{\mu}} t_{\frac{\kappa_{max}}{\mu}}(y) F_{\xi}^{\mathbf{P}}(dy).$$

The completeness of the polynomials $\{p_{\alpha}(\xi), \alpha \in \Lambda\}$ in the space $L^2(\Omega, \sigma(\xi), \mathbf{Q})$ is then equivalent to the density of the polynomials $\{p_{\alpha}(y), \alpha \in \Lambda\}$ in $L^2(\mathbb{R}^{|I_{\xi}|}, \mathfrak{B}(\mathbb{R}^{|I_{\xi}|}), F_{\xi}^{\mathbf{Q}}(dy))$. Some necessary conditions to establish this property are discussed in [13]. For example, the polynomials form an orthonormal basis of $L^2(\Omega, \sigma(\xi), \mathbf{Q})$ if for any marginal distribution

$$\begin{aligned} F_{\xi_1, \dots, \xi_K}^{\mathbf{Q}}(y_1, \dots, y_K) &= \int F_{\xi}^{\mathbf{Q}}(y_1, \dots, y_K, (dy_i)_{i > K}) \\ &= \int \frac{1}{c \frac{\kappa_{max}}{\mu}} t_{\frac{\kappa_{max}}{\mu}}(y) F_{\xi}^{\mathbf{P}}(y_1, \dots, y_K, (dy_i)_{i > K}) \end{aligned}$$

$K \in \mathbb{N}$ with $\{1, \dots, K\} \subseteq I_\xi$, there exists a constant $a > 0$ such that

$$\int \exp \left(a \sqrt{\sum_{i=1}^K y_i^2} \right) F_{\xi_1, \dots, \xi_K}^{\mathbf{Q}}(dy_1, \dots, dy_K) < \infty.$$

In summary, this condition means that there exists a constant $a > 0$ satisfying

$$\begin{aligned} \int \exp \left(a \sqrt{\sum_{i \in I_\xi} y_i^2} \right) F_\xi^{\mathbf{Q}}(dy) &= \mathbf{E}_{\mathbf{Q}} \exp \left(a \sqrt{\sum_{i \in I_\xi} \xi_i^2} \right) \\ &= \frac{1}{c_{\frac{\kappa_{max}}{\mu}}} \mathbf{E}_{\mathbf{P}} \left(\exp \left(a \sqrt{\sum_{i \in I_\xi} \xi_i^2} \right) t_{\frac{\kappa_{max}}{\mu}}(\xi) \right) < \infty. \end{aligned}$$

Assuming that the polynomials are dense in $L^2(\Omega, \sigma(\xi), \mathbf{Q})$ the weak solution $\hat{u} \in \mathring{U}_{\mathbf{Q}}^1$ possesses a generalized polynomial chaos expansion, i. e.

$$\hat{u}(x, \omega) = \sum_{\alpha \in \Lambda} \hat{u}_\alpha(x) p_\alpha(\xi(\omega)),$$

since it is $\sigma(\xi)$ -measurable. The generalized polynomial chaos coefficients are given by

$$\hat{u}_\alpha(x) = \mathbf{E}_{\mathbf{Q}}(\hat{u}(x, \cdot) p_\alpha(\xi)) = \mathbf{E}_{\mathbf{Q}}(\hat{u}_\xi(\xi) p_\alpha(\xi)), \quad \alpha \in \Lambda.$$

Notably, the orthogonal projections $\Pi_{\mathring{U}_{\mathbf{Q},K}^1} \hat{u}$ and $\Pi_{\mathring{U}_{\mathbf{Q},N,K}^1} \hat{u}$ admit a generalized polynomial chaos representation, namely,

$$\Pi_{\mathring{U}_{\mathbf{Q},K}^1} \hat{u}(x, \omega) = \sum_{d(\alpha) \leq K} \hat{u}_\alpha(x) p_\alpha(\xi(\omega)), \quad d(\alpha) := \max\{i \in \mathbb{N} : \alpha_i \neq 0\}$$

and

$$\Pi_{\mathring{U}_{\mathbf{Q},N,K}^1} \hat{u}(x, \omega) = \sum_{\alpha \in \Lambda_{N,K}} \hat{u}_\alpha(x) p_\alpha(\xi(\omega)).$$

Summarizing these results we arrive at the following corollary.

Corollary 3.9 *If the polynomials $\{p_\alpha(\xi), \alpha \in \Lambda\}$ form an orthonormal basis of $L^2(\Omega, \sigma(\xi), \mathbf{Q})$, and $\hat{u} \in \mathring{U}_{\mathbf{Q}}^1$, then the approximation error e_3 converges to zero, i. e.*

$$e_3 = \|\Pi_{\mathring{U}_{\mathbf{Q},K}^1} \hat{u} - \Pi_{\mathring{U}_{\mathbf{Q},N,K}^1} \hat{u}\|_{U_{\mathbf{Q}}^1} \rightarrow 0 \quad \text{when } N, K \rightarrow \infty.$$

In particular, if the polynomials $\{p_\alpha(\xi), \alpha \in \Lambda : d(\alpha) \leq K\}$ form an orthonormal basis of $L^2(\Omega, \sigma(\xi_1, \dots, \xi_K), \mathbf{Q})$ for any $K \in \mathbb{N}$ with $\{1, \dots, K\} \subseteq I_\xi$, we have for any K

$$e_3 = \|\Pi_{\mathring{U}_{\mathbf{Q},K}^1} \hat{u} - \Pi_{\mathring{U}_{\mathbf{Q},N,K}^1} \hat{u}\|_{U_{\mathbf{Q}}^1} \rightarrow 0 \quad \text{for } N \rightarrow \infty.$$

Remark 3.10 *A necessary condition for the completeness of the polynomials $\{p_\alpha(\xi), \alpha \in \Lambda\}$ in $L^2(\Omega, \sigma(\xi), \mathbf{Q})$ is their completeness in the space $L^2(\Omega, \sigma(\xi), \mathbf{P})$.*

Combining the results of the Corollaries 3.6, 3.8 and 3.9 we can now formulate the precise conditions for which the sequence of stochastic Petrov-Galerkin solutions $\bar{u}_{N,K,hp}$ converges to the weak solution \hat{u} in \mathring{U}^1 .

Corollary 3.11 For $\hat{u} \in U_{\mathbf{Q}}^k \cap \hat{U}_{\mathbf{Q}}^1$, $k \geq 2$, if the stochastic Petrov-Galerkin solutions $\bar{u}_{N,K,hp}$ exist and the orthonormal polynomials $\{p_\alpha(\xi), \alpha \in \Lambda\}$ are complete in $L^2(\Omega, \sigma(\xi), \mathbf{Q})$, then the sequence of stochastic Petrov-Galerkin solutions $\bar{u}_{N,K,hp}$ converge in \hat{U}^1 to the weak solution \hat{u} .

Proof.

The result follows immediately from the estimate (3.4) and the Corollaries 3.4, 3.6, 3.8 and 3.9. \square

This means that for the stochastic Petrov-Galerkin solutions $\bar{u}_{N,K,hp}$ associated with the weighted test function space we can establish relatively weak conditions which guarantee convergence in U^1 . We would like to emphasize that for standard stochastic Galerkin solutions $\hat{u}_{N,K,hp}$ associated with unweighted test functions, we need stronger assumptions to prove convergence in U^1 . To illustrate this point we refer to Example 1 in Section 4 ahead where a convergent sequence of stochastic Petrov-Galerkin solutions $\bar{u}_{N,K,hp}$ is constructed whereas the sequence of standard stochastic Galerkin solutions $\hat{u}_{N,K,hp}$ fails to converge. In ongoing research we are also studying the basic random variables' impact on the approximation quality and the rate of convergence of the stochastic Petrov-Galerkin solutions $\bar{u}_{N,K,hp}$.

3.2 Assembly of the stochastic Galerkin equations and computational aspects

The use of weighted test functions affects the assembly of the associated stochastic Galerkin system. In what follows we outline the differences to the unweighted case (see e.g. [18, 12, 26]). Specifically, we address the calculation of the stochastic Petrov-Galerkin solution $\bar{u}_{N,K,hp}$ and associated characteristics, for example, the expectation function $\mathbf{E}_{\mathbf{P}}\bar{u}_{N,K,hp}$ and second order moment function $\mathbf{E}_{\mathbf{P}}\bar{u}_{N,K,hp}^2$.

Let $\kappa_{max}/\mu \in L^r(\Omega, \mathfrak{A}, \mathbf{P})$ for some $r > 1$. Assume $\{q_\alpha(\xi)\varphi_i(x), \alpha \in \Lambda_{N,K}, i \in I_{hp}\}$ is a basis of the space $U_{N,K,hp}$ where the sets $\{\varphi_i(x), i \in I_{hp}\}$ and $\{q_\alpha(\xi), \alpha \in \Lambda_{N,K}\}$ are bases of the spaces U_{hp} and $U_{N,K}$, respectively. Then a basis of the finite-dimensional test function $V_{N,K,hp}$ space is given by

$$\left\{ \frac{q_\alpha(\xi)}{\mu} \varphi_i(x), \alpha \in \Lambda_{N,K}, i \in I_{hp} \right\}.$$

Hence, the stochastic Petrov-Galerkin solution $\bar{u}_{N,K,hp} = \sum_{\substack{\alpha \in \Lambda_{N,K} \\ i \in I_{hp}}} u_{\alpha i} \varphi_i q_\alpha(\xi)$ satisfies the linear

system of equations

$$A\underline{u} = \underline{f}$$

where

$$A := \left(a(q_\alpha \varphi_i, q_\beta \varphi_j / \mu) \right)_{\beta j, \alpha i}, \quad \underline{f} := \left(\ell(\varphi_j q_\beta / \mu) \right)_{\beta j} \quad \text{and} \quad \underline{u} := \left(u_{\alpha i} \right)_{\alpha i}.$$

Using the parameterized input data $\kappa_\xi(\xi)$ with $\kappa_\xi : \mathbb{R}^{|\mathcal{I}_\xi|} \rightarrow L^\infty(D)$ and $f_\xi(\xi)$ with

$f_\xi : \mathbb{R}^{|\mathcal{I}_\xi|} \rightarrow H^{-1}(D)$ we obtain

$$\begin{aligned} a(\varphi_i q_\alpha, \varphi_j q_\beta / \mu) &= \mathbf{E}_{\mathbf{P}} \left(\left(\int_D [\kappa_\xi(\xi)](x) \nabla \varphi_i(x) \cdot \nabla \varphi_j(x) dx \right) q_\alpha(\xi) q_\beta(\xi) / \mu \right) \\ &= \int_D \mathbf{E}_{\mathbf{P}} \left(\frac{[\kappa_\xi(\xi)](x)}{\mu} q_\alpha(\xi) q_\beta(\xi) \right) \nabla \varphi_i(x) \cdot \nabla \varphi_j(x) dx \end{aligned}$$

and

$$\ell(\varphi_j q_\beta / \mu) = \mathbf{E}_{\mathbf{P}} \left(\langle f_\xi(\xi), \varphi_j \rangle_{H^{-1}, \dot{H}^1} q_\beta(\xi) / \mu \right) = \left\langle \mathbf{E}_{\mathbf{P}} \left(\frac{f_\xi(\xi)}{\mu} q_\beta(\xi) \right), \varphi_j \right\rangle_{H^{-1}, \dot{H}^1}$$

for any $i, j \in I_{hp}$, $\alpha, \beta \in \Lambda_{N,K}$.

It is important to choose suitable polynomials $\{q_\alpha(\xi), \alpha \in \Lambda_{N,K}\}$ so that the computation of the expectations in the linear system is actually feasible. To this end, we discuss two options.

First, it is possible to employ the polynomials $\{\bar{p}_\alpha(\xi), \alpha \in \Lambda\}$ which form an orthonormal basis of $L^2(\Omega, \sigma(\xi), \mathbf{P})$. Thus, evaluation of the expected values

$$\mathbf{E}_{\mathbf{P}} \left(\kappa_\xi(\xi) \bar{p}_\alpha(\xi) \bar{p}_\beta(\xi) / \mu \right) \quad \text{and} \quad \mathbf{E}_{\mathbf{P}} \left(f_\xi(\xi) \bar{p}_\beta(\xi) / \mu \right)$$

requires the calculation of the generalized polynomial chaos expansions of $\kappa_\xi(\xi)/\mu$ and $f_\xi(\xi)/\mu$ or their generalized polynomial chaos coefficients, respectively. The corresponding characteristics of the stochastic Petrov-Galerkin solution $\bar{u}_{N,K,hp}$, i. e. expected value and second order moment function

$$\mathbf{E}_{\mathbf{P}} \bar{u}_{N,K,hp} \quad \text{and} \quad \mathbf{E}_{\mathbf{P}} \bar{u}_{N,K,hp}^2$$

are determined as usual by

$$\mathbf{E}_{\mathbf{P}} \bar{u}_{N,K,hp} = \mathbf{E}_{\mathbf{P}} \left(\sum_{\substack{\alpha \in \Lambda_{N,K} \\ i \in I_{hp}}} u_{\alpha i} \varphi_i \bar{p}_\alpha(\xi) \right) = \sum_{\substack{\alpha \in \Lambda_{N,K} \\ i \in I_{hp}}} u_{\alpha i} \varphi_i \underbrace{\mathbf{E}_{\mathbf{P}} (\bar{p}_\alpha(\xi))}_{=\delta_{\alpha,0}} = \sum_{i \in I_{hp}} u_{0i} \varphi_i$$

and

$$\begin{aligned} \mathbf{E}_{\mathbf{P}} \bar{u}_{N,K,hp}^2 &= \mathbf{E}_{\mathbf{P}} \left(\sum_{\substack{\alpha, \beta \in \Lambda_{N,K} \\ i, j \in I_{hp}}} u_{\alpha i} u_{\beta j} \varphi_i \varphi_j \bar{p}_\alpha(\xi) \bar{p}_\beta(\xi) \right) \\ &= \sum_{\substack{\alpha, \beta \in \Lambda_{N,K} \\ i, j \in I_{hp}}} u_{\alpha i} \varphi_i u_{\beta j} \varphi_j \underbrace{\mathbf{E}_{\mathbf{P}} (\bar{p}_\alpha(\xi) \bar{p}_\beta(\xi))}_{=\delta_{\alpha, \beta}} \\ &= \sum_{\alpha \in \Lambda_{N,K}} \sum_{i, j \in I_{hp}} u_{\alpha i} u_{\alpha j} \varphi_i \varphi_j = \sum_{\alpha \in \Lambda_{N,K}} \left(\sum_{i \in I_{hp}} u_{\alpha i} \varphi_i \right)^2. \end{aligned}$$

A second option can be obtained if the expectation of the random variable μ^{-1} is finite and $\mathbf{E}_{\mathbf{P}} |\xi_i|^n \mu^{-1} < \infty$ holds for all $n \in \mathbb{N}$ and $i \in I_\xi$. For example, this is satisfied if $H^{-1}(D) \subset U_{\mu^{-2}}^{-1}$. Then, w.l.o.g. we can assume that the expectation of μ^{-1} is equal to one. In this case the polynomials $\{\tilde{p}_\alpha(\xi), \alpha \in \Lambda\}$ which are orthonormal in $L^2(\Omega, \mathfrak{A}, \mu^{-1} d\mathbf{P})$ can serve as basis polynomials $q_\alpha(\xi)$. Then, the computation of the expected values $\mathbf{E}_{\mathbf{P}} \left(\kappa_\xi(\xi) \tilde{p}_\alpha(\xi) \tilde{p}_\beta(\xi) / \mu \right)$ and $\mathbf{E}_{\mathbf{P}} \left(f_\xi(\xi) \tilde{p}_\beta(\xi) / \mu \right)$ requires only the generalized

polynomial chaos expansions of κ_ξ and f_ξ in the polynomials $\{\tilde{p}_\alpha(\xi), \alpha \in \Lambda\}$. However, evaluating the first and second order moment function of the stochastic Petrov-Galerkin solution is more complicated than in the first option since the polynomials $\{\tilde{p}_\alpha(\xi), \alpha \in \Lambda\}$ do not satisfy $\mathbf{E}_\mathbf{P}(\tilde{p}_\alpha(\xi)\tilde{p}_\beta(\xi)) = C_\alpha\delta_{\alpha,\beta}$. Thus, we have to compute

$$\mathbf{E}_\mathbf{P}\bar{u}_{N,K,hp} = \mathbf{E}_\mathbf{P}\left(\sum_{\substack{\alpha \in \Lambda_{N,K} \\ i \in I_{hp}}} u_{\alpha i} \varphi_i \tilde{p}_\alpha(\xi)\right) = \sum_{\substack{\alpha \in \Lambda_{N,K} \\ i \in I_{hp}}} u_{\alpha i} \varphi_i \mathbf{E}_\mathbf{P} \tilde{p}_\alpha(\xi)$$

and

$$\mathbf{E}_\mathbf{P}\bar{u}_{N,K,hp}^2 = \mathbf{E}_\mathbf{P}\left(\sum_{\substack{\alpha, \beta \in \Lambda_{N,K} \\ i, j \in I_{hp}}} u_{\alpha i} u_{\beta j} \varphi_i \varphi_j \tilde{p}_\alpha(\xi) \tilde{p}_\beta(\xi)\right) = \sum_{\substack{\alpha, \beta \in \Lambda_{N,K} \\ i, j \in I_{hp}}} u_{\alpha i} \varphi_i u_{\beta j} \varphi_j \mathbf{E}_\mathbf{P}(\tilde{p}_\alpha(\xi) \tilde{p}_\beta(\xi)).$$

Suppose now, that the generalized polynomial chaos expansions of the polynomials $\tilde{p}_\alpha(\xi)$, $\alpha \in \Lambda_{N,K}$, are given in terms of sums of the polynomials $\{\bar{p}_\alpha(\xi), \alpha \in \Lambda\}$, i. e., we know the generalized polynomial chaos coefficients $\tilde{c}_\beta^{(\alpha)}$ satisfying

$$\tilde{p}_\alpha(\xi) = \sum_{\beta \in \Lambda_{N,K}} \tilde{c}_\beta^{(\alpha)} \bar{p}_\beta(\xi).$$

These coefficients are sometimes called *connection coefficients* in the literature [23, 29, 31]. We then obtain the stochastic Petrov-Galerkin solution in terms of the polynomials $\{\bar{p}_\alpha(\xi), \alpha \in \Lambda\}$ by rearranging

$$\bar{u}_{N,K,hp} = \sum_{\substack{\alpha \in \Lambda_{N,K} \\ i \in I_{hp}}} u_{\alpha i} \varphi_i \tilde{p}_\alpha(\xi) = \sum_{\substack{\alpha \in \Lambda_{N,K} \\ i \in I_{hp}}} u_{\alpha i} \varphi_i \sum_{\beta \in \Lambda_{N,K}} \tilde{c}_\beta^{(\alpha)} \bar{p}_\beta(\xi) = \sum_{\beta \in \Lambda_{N,K}} \bar{p}_\beta(\xi) \left[\sum_{\substack{\alpha \in \Lambda_{N,K} \\ i \in I_{hp}}} \tilde{c}_\beta^{(\alpha)} u_{\alpha i} \varphi_i \right].$$

The expected value and second order moment function can be computed as follows

$$\mathbf{E}_\mathbf{P}\bar{u}_{N,K,hp} = \sum_{\substack{\alpha \in \Lambda_{N,K} \\ i \in I_{hp}}} \tilde{c}_0^{(\alpha)} u_{\alpha i} \varphi_i \quad \text{and} \quad \mathbf{E}_\mathbf{P}\bar{u}_{N,K,hp}^2 = \sum_{\beta \in \Lambda_{N,K}} \left[\sum_{\substack{\alpha \in \Lambda_{N,K} \\ i \in I_{hp}}} \tilde{c}_\beta^{(\alpha)} u_{\alpha i} \varphi_i \right]^2.$$

If both sets of polynomials are basis polynomials of the finite-dimensional subspace $U_{N,K}$, then the resulting stochastic Galerkin solutions coincide, of course. Depending on the problem, however, it can happen that the two options differ with respect to computational aspects and performance.

Example 3.12 Assume that for a given boundary-value problem the basic random variable is a single standard Gaussian random variable ξ and that the random variable $1/\mu = t_{\mu-1}(\xi) = \exp\left(\frac{\varepsilon}{2}\xi^2\right)/\sigma$ with $\sigma = 1/\sqrt{1-\varepsilon}$ for $0 < \varepsilon < 1$ serves as an admissible weighting random variable. Then the basic random variable ξ has also a Gaussian distribution on the probability space $(\Omega, \mathfrak{A}, \mu^{-1}d\mathbf{P})$ and the corresponding density function $f_\xi^{\mu^{-1}d\mathbf{P}}$ is given by

$$\begin{aligned} f_\xi^{\mu^{-1}d\mathbf{P}}(y) &= t_{\mu-1}(y) f_\xi^\mathbf{P}(y) = \frac{1}{\sqrt{2\pi}\sigma} \exp\left(\frac{\varepsilon}{2}y^2\right) \exp\left(-\frac{y^2}{2}\right) \\ &= \frac{1}{\sqrt{2\pi}\sigma} \exp\left(-\frac{y^2(1-\varepsilon)}{2}\right) = \frac{1}{\sqrt{2\pi}\sigma} \exp\left(-\frac{y^2}{2\sigma^2}\right), \quad y \in \mathbb{R}. \end{aligned}$$

The associated orthonormal polynomials $\{\tilde{p}_n(\xi), n \in \mathbb{N}_0\}$ in $L^2(\Omega, \sigma(\xi), \mu^{-1}d\mathbf{P})$ are the normalized probabilistic Hermite polynomials $\{He_n(\xi), n \in \mathbb{N}_0\}$ scaled in the argument, i. e.

$$\tilde{p}_n(\xi) = \frac{1}{\sqrt{n!}} He_n\left(\frac{\xi}{\sigma}\right), \quad n \in \mathbb{N}_0.$$

If we assemble the stochastic Galerkin system by the first option described above, the original Hermite polynomials

$$\bar{p}_n(\xi) = \frac{He_n(\xi)}{\sqrt{n!}}, \quad n = 0, \dots, N$$

are used to span the finite-dimensional subspace $U_{N,K}$ with $K = 1$. Therefore, we have to calculate for instance the polynomial chaos coefficients of $\kappa_\xi(\xi)/\mu$, i. e.

$$\mathbf{E}_{\mathbf{P}} \left(\frac{\kappa_\xi(\xi)}{\mu} \frac{He_n(\xi)}{\sqrt{n!}} \right), \quad n \in \mathbb{N}_0.$$

Using the second proposed option we have to calculate the generalized polynomial chaos coefficients of κ and f w. r. t. the polynomials $\{\tilde{p}_n(\xi), n \in \mathbb{N}_0\}$. For the coefficient κ this requires evaluation of the polynomial chaos coefficients of $\kappa_\xi(\sigma\xi)$ by the change of variables $z = y/\sigma$

$$\begin{aligned} \mathbf{E}_{\mathbf{P}} \left(\frac{\kappa_\xi(\xi)}{\mu} \frac{He_n(\xi/\sigma)}{\sqrt{n!}} \right) &= \int_{\mathbb{R}} \kappa_\xi(y) \frac{He_n(y/\sigma)}{\sqrt{n!}} t_{\mu^{-1}}(y) f_\xi^{\mathbf{P}}(y) dy = \int_{\mathbb{R}} \kappa_\xi(y) \frac{He_n(y/\sigma)}{\sqrt{n!}} f_\xi^{\mu^{-1}d\mathbf{P}}(y) dy \\ &= \int_{\mathbb{R}} \kappa_\xi(\sigma z) \frac{He_n(z)}{\sqrt{n!}} f_\xi^{\mathbf{P}}(z) dz = \mathbf{E}_{\mathbf{P}} \left(\kappa_\xi(\sigma\xi) \frac{He_n(\xi)}{\sqrt{n!}} \right), \quad n \in \mathbb{N}_0. \end{aligned}$$

In order to obtain the first and second order moments of the approximate solution, we need the generalized polynomial chaos coefficients $c_i^{(n)}$ satisfying

$$\frac{He_n(\xi/\sigma)}{\sqrt{n!}} = \sum_{i=0}^n c_i^{(n)} \frac{He_i(\xi)}{\sqrt{i!}} \quad n \in \mathbb{N}_0.$$

Fortunately, in this case analytic expressions for these coefficients are available:

$$c_i^{(n)} = \begin{cases} \frac{\sqrt{n!}}{\sqrt{i!} \left(\frac{n-i}{2}\right)!} \sigma^{-n} \left(\frac{1-\sigma^2}{2}\right)^{(n-i)/2}, & n+i \text{ even}, \\ 0, & \text{otherwise}, \end{cases} \quad i = 0, \dots, n, \quad n \in \mathbb{N}_0. \quad \blacksquare$$

If the multivariate orthonormal polynomials $\{\bar{p}_\alpha(\xi), \alpha \in \Lambda\}$ or $\{\tilde{p}_\alpha(\xi), \alpha \in \Lambda\}$ can be represented as tensor products of univariate orthonormal polynomials, then the assembly of the stochastic Galerkin system simplifies significantly. To answer this question, we need to investigate if the basic random variables $(\xi_i)_{i \in I_\xi}$ are independent as random variables on the probability spaces $(\Omega, \mathfrak{A}, \mathbf{P})$ or $(\Omega, \mathfrak{A}, \mu^{-1}d\mathbf{P})$, respectively. Let $t_{\mu^{-1}} : \mathbb{R}^{|I_\xi|} \rightarrow \mathbb{R}$ be a measurable function such that there holds

$$t_{\mu^{-1}}(\xi) = \mu^{-1}.$$

Then the distribution $F_\xi^{\mu^{-1}d\mathbf{P}}$ of ξ on $(\Omega, \mathfrak{A}, \mu^{-1}d\mathbf{P})$ is

$$F_\xi^{\mu^{-1}d\mathbf{P}} = t_{\mu^{-1}}(y) F_\xi^{\mathbf{P}}(dy).$$

Analogously, the distribution $F_{\xi_i}^{\mu^{-1}d\mathbf{P}}$ of each random variable $\xi_i, i \in I_\xi$, is given by

$$F_{\xi_i}^{\mu^{-1}d\mathbf{P}}(dy_i) = t_{\mu^{-1},i}(y_i) F_{\xi_i}^{\mathbf{P}}(dy_i),$$

where $t_{\mu^{-1},i} : \mathbb{R} \rightarrow \mathbb{R}$ is the measurable function satisfying

$$t_{\mu^{-1},i}(\xi_i) = \mathbf{E}_{\mathbf{P}} \left(\mu^{-1} | \xi_i \right), \quad i \in I_\xi.$$

Then, the independence of the basic random variables $(\xi_i)_{i \in I_\xi}$ on the respective probability spaces holds if and only if the associated distribution $F_\xi^{\mathbf{P}}$ or $F_\xi^{\mu^{-1}d\mathbf{P}}$ is the product of the distributions of the individual random variables ξ_i , $i \in I_\xi$, i. e.,

$$F_\xi^{\mathbf{P}}(dy) = \prod_{i \in I_\xi} F_{\xi_i}^{\mathbf{P}}(dy_i) \quad \text{or} \quad F_\xi^{\mu^{-1}d\mathbf{P}}(dy) = \prod_{i \in I_\xi} F_{\xi_i}^{\mu^{-1}d\mathbf{P}}(dy_i),$$

respectively. If, for instance, the random variables $(\xi_i)_{i \in I_\xi}$ are independent on the probability space $(\Omega, \mathfrak{A}, \mathbf{P})$ and the function $t_{\mu^{-1}}$ can be represented as product

$$t_{\mu^{-1}}(y) = \prod_{i \in I_\xi} t_{\mu^{-1},i}(y_i),$$

then the random variables $(\xi_i)_{i \in I_\xi}$ are also independent on the probability space $(\Omega, \mathfrak{A}, \mu^{-1}d\mathbf{P})$.

4 Examples

In this section we present two examples of boundary-value problems which illustrate the different convergence behavior of the standard stochastic Galerkin approach and the stochastic Petrov-Galerkin approach, respectively. Because the difference is related to the stochastic aspects of the problems, we construct very simple one-dimensional boundary-value problems where the random coefficients depend on a single basic random variable. In both examples we will observe that

- the standard stochastic Galerkin approach which is usually used (see e. g. [26, 38]) fails to provide a sequence of approximate solutions converging to the exact solution in the natural norm,
- the stochastic Petrov-Galerkin approach with a suitable weighted test function space yields a sequence of approximate solutions converging to the exact solution in the natural norm.

Both results, non-convergence for the standard stochastic Galerkin approach and convergence for the stochastic Petrov-Galerkin approach, respectively, are illustrated numerically. In addition, we prove non-convergence of the standard stochastic Galerkin approach for Example 1 analytically.

In both examples we study a one-dimensional boundary-value problem of the type

$$-(\kappa(x, \omega)u'(x, \omega))' = f(x, \omega), \quad x \in (0, 1), \quad \omega \in \Omega,$$

with homogeneous Dirichlet boundary conditions

$$u(0, \omega) = u(1, \omega) = 0, \quad \omega \in \Omega,$$

or mixed boundary conditions

$$u(0, \omega) = 0 \quad \text{and} \quad \kappa(1, \omega)u'(1, \omega) = F(\omega), \quad \omega \in \Omega.$$

Though we do not consider mixed boundary conditions in our model problem (2.1) in Section 1, the theory developed in the preceding sections still applies. To eliminate the spatial discretization error as far as possible we use a single Gauss-Lobatto-Legendre spectral element of suitable degree p for the spatial discretization. Thus, we are able to observe the effect of the different discretizations of the stochastic space. In the experiments we always

compare the corresponding relative approximation error in the U^1 -norm – the natural norm, i. e.,

$$\frac{\|\hat{u} - \hat{u}_{N,K,hp}\|_{U^1}}{\|\hat{u}\|_{U^1}} \quad \text{and} \quad \frac{\|\hat{u} - \bar{u}_{N,K,hp}\|_{U^1}}{\|\hat{u}\|_{U^1}},$$

respectively. Furthermore, since we consider a single basic random variable, we drop the index K in the notation of the finite-dimensional subspaces and stochastic Galerkin solutions for the sake of convenience.

Example 1. We consider the boundary-value problem on the domain $D = (0, 1)$ with random coefficient $\kappa(x, \omega) = \exp(\xi(\omega))$ and random forcing term $f(x, \omega) = \exp(\xi(\omega))|\xi(\omega) - 1|$ where $\xi \sim \mathcal{N}(0, 1)$ is a standard Gaussian random variable. This means, we are looking for the solution of

$$\begin{aligned} -(\exp(\xi(\omega))u'(x, \omega))' &= \exp(\xi(\omega))|\xi(\omega) - 1|, & x \in (0, 1), \omega \in \Omega, \\ u(0, \omega) &= u(1, \omega) = 0, & \omega \in \Omega. \end{aligned}$$

Since the coefficient κ is strongly measurable in $L^\infty(D)$ and there holds

$$0 < \kappa_{min} := \exp(\xi) \leq \kappa \leq \exp(\xi) =: \kappa_{max} < \infty$$

we can choose $\mu = \kappa_{min}$. Since $f \in U_{\exp(-2\xi)}^{-1} \cap U_{\exp(-\xi)}^{-1}$ we can also conclude that there exists a unique weak solution $\hat{u} \in \dot{U}^1$ given by

$$\hat{u}(x, \omega) = [\hat{u}_\xi(\xi(\omega))](x) = \frac{1}{2}|\xi(\omega) - 1|(x - x^2), \quad x \in (0, 1), \omega \in \Omega.$$

This is of course also a pathwise strong solution.

For the standard stochastic Galerkin approach we choose identical finite-dimensional solution and test function spaces $U_{N,hp} = U_{hp} \otimes U_N$ where

$$U_N = \text{span} \left\{ \frac{\text{He}_n(\xi)}{\sqrt{n!}}, n = 0, \dots, N \right\}$$

is spanned by the first $N + 1$ probabilistic Hermite polynomials $\{\text{He}_n(\xi), n \in \mathbb{N}_0\}$. The relative approximation error

$$err = \frac{\|\hat{u} - \hat{u}_{N,hp}\|_{U^1}}{\|\hat{u}\|_{U^1}}$$

of the corresponding standard stochastic Galerkin solutions $\hat{u}_{N,hp}$ is shown in Figure 1. Here, we use a single Gauss-Lobatto-Legendre spectral element of degree $p = 3$ for the spatial discretization. We clearly see that the approximation error does not converge to zero for increasing orders of the chaos polynomials. In fact, we can prove that the sequence of standard stochastic Galerkin solutions $\hat{u}_{N,hp}$ does not converge in \dot{U}^1 to the exact solution for $N \rightarrow \infty$. To see this, we note that due to the special structure of the boundary-value problem the standard stochastic Galerkin solution

$$\hat{u}_{N,hp}(x, \xi) = \hat{u}_{hp}(x)\hat{u}_N(\xi)$$

is the product of the deterministic function $\hat{u}_{hp}(x)$ which satisfies the equation

$$\int_D \hat{u}'_{hp}(x)v'_{hp}(x) dx = \int_D v_{hp}(x) dx \quad \text{for all } v_{hp} \in U_{hp} \quad (4.1)$$




Figure 1: Relative approximation error err of the standard stochastic Galerkin solution $\hat{u}_{N,hp}$ for different orders N of polynomial chaos.

and the random variable $\hat{u}_N(\xi)$ satisfying the equation

$$\mathbf{E}_{\mathbf{P}} \left(\exp(\xi) \hat{u}_N(\xi) \frac{\text{He}_m(\xi)}{\sqrt{m!}} \right) = \mathbf{E}_{\mathbf{P}} \left(\exp(\xi) |\xi - 1| \frac{\text{He}_m(\xi)}{\sqrt{m!}} \right) \quad \text{for all } m = 0, \dots, N. \quad (4.2)$$

We denote $\hat{u}_D(x) := \frac{1}{2}(x - x^2)$ and $\hat{u}_\Omega(\xi) := |\xi - 1|$ and obtain the following lower bound

$$\begin{aligned} \|\hat{u} - \hat{u}_{N,hp}\|_{U^1} &= \|\hat{u}_D \hat{u}_\Omega(\xi) - \hat{u}_{hp} \hat{u}_N(\xi)\|_{U^1} = \|\hat{u}_\Omega(\xi)(\hat{u}_D - \hat{u}_{hp}) + \hat{u}_{hp}(\hat{u}_\Omega(\xi) - \hat{u}_N(\xi))\|_{U^1} \\ &\geq \|\hat{u}_{hp}(\hat{u}_\Omega(\xi) - \hat{u}_N(\xi))\|_{U^1} - \|\hat{u}_\Omega(\xi)(\hat{u}_D - \hat{u}_{hp})\|_{U^1} \\ &= \|\hat{u}_{hp}\|_{H^1(D)} \|\hat{u}_\Omega(\xi) - \hat{u}_N(\xi)\|_{L^2(\Omega, \mathfrak{A}, \mathbf{P})} - \|\hat{u}_\Omega(\xi)\|_{L^2(\Omega, \mathfrak{A}, \mathbf{P})} \|\hat{u}_D - \hat{u}_{hp}\|_{H^1(D)}. \end{aligned}$$

In view of

$$\|\hat{u}_D - \hat{u}_{hp}\|_{H^1(D)} \rightarrow 0 \quad \text{and} \quad \|\hat{u}_\Omega(\xi)\|_{L^2(\Omega, \mathfrak{A}, \mathbf{P})} < \infty$$

the second error term $(\|\hat{u}_\Omega(\xi)\|_{L^2(\Omega, \mathfrak{A}, \mathbf{P})} \|\hat{u}_D - \hat{u}_{hp}\|_{H^1(D)})$ converges to zero and the norm of $\|\hat{u}_{hp}\|_{H^1(D)}$ converges to $\|\hat{u}_D\|_{H^1(D)}$. However, the stochastic part $\hat{u}_N(\xi)$ does not converge to $\hat{u}_\Omega(\xi)$ in $L^2(\Omega, \mathfrak{A}, \mathbf{P})$ as shown in detail in the Appendix A. In fact, there exists a subsequence $(\hat{u}_{N_k}(\xi))_{k \in \mathbb{N}_0}$ of $(\hat{u}_N(\xi))_{N \in \mathbb{N}_0}$ whose second order moments tend to infinity which implies that the error $\|\hat{u}_\Omega(\xi) - \hat{u}_{N_k}(\xi)\|_{L^2(\Omega, \mathfrak{A}, \mathbf{P})}$ goes to infinity, i. e.,

$$\|\hat{u}_\Omega(\xi) - \hat{u}_{N_k}(\xi)\|_{L^2(\Omega, \mathfrak{A}, \mathbf{P})} \rightarrow \infty, \quad k \rightarrow \infty.$$

Of course, this also implies that the corresponding approximation error err goes to infinity, i. e.,

$$\|\hat{u} - \hat{u}_{N_k,hp}\|_{U^1} \rightarrow \infty, \quad k, p \rightarrow \infty.$$

Thus, the sequence of standard stochastic Galerkin solutions $\hat{u}_{N,hp}$ does not converge to the exact solution \hat{u} in \hat{U}^1 .

Alternatively, we employ the proposed stochastic Petrov-Galerkin approach. W.l.o.g. we can normalize the random variable μ such that the expectation of its reciprocal μ^{-1} is one, i. e., we take

$$\mu := \exp(1/2) \exp(\xi).$$

We define the weighted finite-dimensional test function space

$$V_{N,hp} = \{u \exp(-\xi), u \in U_{N,hp}\} = \text{span} \left\{ u_{hp} \frac{\text{He}_n(\xi)}{\sqrt{n!}} \exp(-\xi), u_{hp} \in U_{hp}, n = 0, \dots, N \right\}.$$

Since $\kappa_{max} = \exp(-1/2)\mu$ the probability measure \mathbf{Q} coincides with the probability measure \mathbf{P} and thus $\hat{U}^1 = \hat{U}_{\mathbf{Q}}^1$. Hence, based on Corollary 3.4, we obtain for the corresponding stochastic Galerkin solution $\bar{u}_{N,hp}$ the estimate

$$\|\hat{u} - \bar{u}_{N,hp}\|_{U^1} \leq C \inf_{z \in \hat{U}_{N,hp}} \|\hat{u} - z\|_{U^1}$$

where exactly the same norms appear on the left and right hand side of the inequality. Due to $\hat{u} \in U^k \cap \hat{U}^1$ for any $k \geq 1$ and the completeness of the Hermite polynomials in the space $L^2(\Omega, \sigma(\xi), \mathbf{P})$ the sequence of the stochastic Petrov-Galerkin solutions $\bar{u}_{N,hp}$ converges to the exact solution. This is illustrated by the approximation error err on the left side of Figure 2. Here we have again used a single Gauss-Lobatto-Legendre spectral element of




Figure 2: Left: Relative approximation error err of the stochastic Petrov-Galerkin solution $\bar{u}_{N,hp}$ for different orders N of polynomial chaos. Right: Second order moment function of the exact solution \hat{u} and the standard stochastic Galerkin solution $\hat{u}_{N,hp}$ and stochastic Petrov-Galerkin solution $\bar{u}_{N,hp}$ for polynomial chaos of order $N = 40$.

degree $p = 3$ for the spatial discretization. Again, any solution $\bar{u}_{N,hp}$ is the product of a deterministic function \bar{u}_{hp} and a random variable $\bar{u}_N(\xi)$ satisfying analogous equations as (4.1) and (4.2) for \hat{u}_{hp} and $\hat{u}_N(\xi)$. Note that in this case, the stochastic part $\bar{u}_N(\xi)$ is the *exact* polynomial chaos approximation of order N of $\hat{u}_\Omega(\xi)$, hence we compute the best approximation with respect to the norm of the space $L^2(\Omega, \sigma(\xi), \mathbf{P})$. For example, for $N = 0$ the expectation of the stochastic Petrov-Galerkin solution $\bar{u}_{0,hp}$ is the exact expectation of \hat{u} as function of the spatial argument $x \in D$. On the right side of Figure 2 (using a single Gauss-Lobatto-Legendre spectral element of degree $p = 50$ for the spatial discretization) the second order moment function of the exact solution (blue- \square line)

$$\mathbf{E}_{\mathbf{P}} \hat{u}^2(x, \cdot) = \frac{1}{2}(x - x^2)^2, \quad x \in D,$$

is compared with the second order moment function of the stochastic Galerkin solutions $\hat{u}_{N,hp}$ (green-+ line) and $\bar{u}_{N,hp}$ (red- \times line) for generalized polynomial chaos order $N = 40$. As expected, the second order moment function of the stochastic Petrov-Galerkin solution $\bar{u}_{N,hp}$ approximates well the second order moment function of the exact solution whereas the second order moment function of the standard stochastic Galerkin solution $\hat{u}_{N,hp}$ differs substantially. ■

Our second example is again a one-dimensional boundary-value problem but with a more sophisticated random coefficient, a deterministic forcing term and mixed boundary conditions.

Example 2. We study the following boundary-value problem

$$\begin{aligned} -\left(\exp\left(-\xi^2(\omega)x/10\right)u'(x,\omega)\right)' &= f(x), & x \in (0,1), \omega \in \Omega, \\ u(0,\omega) &= 0, & \omega \in \Omega, \\ \exp\left(-\xi^2(\omega)/10\right)u'(1,\omega) &= \exp\left(-6\xi^2(\omega)\right), & \omega \in \Omega, \end{aligned}$$

with random coefficient $\kappa(x,\omega) = \exp(-\xi^2(\omega)x/10)$, deterministic forcing term $f \in C[0,1] \subset H^{-1}(D)$ and random boundary value $F(\omega) = \exp(-6\xi^2(\omega))$ where the random variable ξ is again standard Gaussian. The coefficient κ depends on $\omega \in \Omega$ as well as on the spatial argument $x \in D$, is strongly measurable and bounded by the random variables

$$\kappa_{min} := \exp(-\xi^2/10) \quad \text{and} \quad \kappa_{max} \equiv 1,$$

respectively, satisfying

$$0 < \kappa_{min} \leq \kappa \leq \kappa_{max} < \infty \quad \text{a. e. and a. s.}$$

The exact solution \hat{u} of the boundary-value problem is explicitly given by

$$\hat{u}(x,\omega) = \int_0^x \exp\left(\xi^2(\omega)y/10\right) \left(\exp\left(-6\xi^2(\omega)\right) + \int_y^1 f(z) dz\right) dy.$$

Alternatively, any random variable of the form $\exp\left(-\frac{\varepsilon}{10}\xi^2\right)$ with $\varepsilon \geq 1$ is also a lower bound for the coefficient κ . Since we want to work with random variables for which the reciprocal has expectation one and is integrable of an order greater than one we consider

$$\mu_\varepsilon := \sqrt{\frac{5}{5-\varepsilon}} \exp\left(-\frac{\varepsilon}{10}\xi^2\right), \quad 1 \leq \varepsilon < 5.$$

Then, for $\varepsilon \leq 3$ the expectation

$$\mathbf{E}_{\mathbf{P}}\left(|\hat{u}|_{H^1}^2 \mu_\varepsilon^{-1}\right) = \sqrt{\frac{5-\varepsilon}{5}} \mathbf{E}_{\mathbf{P}}\left(|\hat{u}|_{H^1}^2 \exp\left(\frac{\varepsilon}{10}\xi^2\right)\right)$$

is also finite. Note that for $\varepsilon > 3$ the latter property is in general not satisfied. For $1 \leq \varepsilon \leq 3$ there exists a unique weak solution \hat{u} in U^1 , especially in $U_{\mu_\varepsilon}^1$. Therefore, any μ_ε with $1 \leq \varepsilon \leq 3$ can serve as a suitable weighting random variable. This allows us to define the weighted test function spaces

$$V_{N,hp}^{(\varepsilon)} = \{u/\mu_\varepsilon, u \in U_{N,hp}\}$$

where

$$U_{N,hp} = U_{hp} \otimes U_N \quad \text{with} \quad U_N = \text{span}\{\xi^n, n = 0, \dots, N\}$$

is the finite-dimensional solution space. Now, any sequence of the stochastic Petrov-Galerkin solutions $(\bar{u}_{N,hp}^{(\varepsilon)})_{N,hp}$ with $1 \leq \varepsilon \leq 3$ converges to the exact solution \hat{u} in U^1 , since there holds $\hat{u} \in U_{\mu_\varepsilon}^{\kappa_{max}} \cap U_{\mu_\varepsilon}^{\kappa_{max}}$ for any $k \geq 1$ and the corresponding polynomials are dense in $L^2(\Omega, \sigma(\xi), \kappa_{max}/\mu_\varepsilon d\mathbf{P})$. For the constant forcing term $f \equiv 1$ we compute the relative approximation errors err of the stochastic Petrov-Galerkin solutions $\bar{u}_{N,hp}^{(\varepsilon)}$ for different values ε , namely $\varepsilon = 1, 2, 3$, using a single Gauss-Lobatto-Legendre spectral element of degree $p = 50$ for the spatial discretization. The results are presented in Figure 3.




Figure 3: Relative approximation error err as function of the generalized polynomial chaos order N for different weights μ_ε , $\varepsilon = 0, 1, 2, 3$.

Obviously the error decays according to the theory developed in the preceding sections. For comparison purposes the relative approximation error of the standard stochastic Galerkin solution $\hat{u}_{N,hp}$ for the unweighted test function space, i. e. $\varepsilon = 0$, is also shown in Figure 3. After a period of decrease the approximation error increases suggesting that this sequence of standard stochastic Galerkin solutions does not converge to the exact solution \hat{u} in U^1 . ■

5 Summary

In this work we have presented two possible stochastic variational formulations – an unweighted and a weighted one – for elliptic partial differential equations with random coefficients where the coefficients are bounded strictly away from zero and infinity by random variables. For each formulation we have established a solution theory and applied the stochastic Galerkin method for the numerical solution. It turns out that the stochastic Galerkin method employed on the unweighted weak formulation corresponds to the standard stochastic Galerkin approach whereas the stochastic Galerkin method employed on the weighted weak formulation can be interpreted as a stochastic Petrov-Galerkin approach since the test function space does no longer coincide with the solution space but corresponds to the solution space weighted by a suitable random variable. We have analyzed the convergence of these two different approximations to the exact solution. We have presented an example where the standard stochastic Galerkin approach produces approximations which do not converge to the exact solution in the natural norm. This does not mean, however, that this approach never yields convergent approximate solutions to the weak problem. The convergence of the standard stochastic Galerkin approximation depends on the underlying boundary-value problem and must be proved separately. As possible remedy we have presented the stochastic Petrov-Galerkin approach with a modified test function space. We have established conditions for the convergence of the stochastic Petrov-Galerkin solutions in the natural norm and presented examples where this approach produces convergent approximations despite the failure of the standard stochastic Galerkin approach.

A On the convergence of $\hat{u}_N(\xi)$ in Example 1

We consider the algebraic equation

$$\kappa u = f$$

for a given random coefficient κ with

$$\kappa(\omega) = \exp(\xi(\omega))$$

and random right side f with

$$f(\omega) = \exp(\xi(\omega))|\xi(\omega) - 1|$$

where the random variable ξ is standard Gaussian. Clearly, the exact solution is

$$u(\omega) = u_\xi(\xi(\omega)) = |\xi(\omega) - 1| \in L^2(\Omega, \sigma(\xi), \mathbf{P}).$$

As discussed in Example 1 of Section 4 the finite-dimensional solution and test function space of the standard stochastic Galerkin approach is the subspace

$$U_N = \text{span} \left\{ \frac{\text{He}_n(\xi)}{\sqrt{n!}}, n = 0, \dots, N \right\}$$

spanned by the probabilistic Hermite polynomials. Thus, the corresponding solution $u_N \in U_N$ solving

$$\mathbf{E}_{\mathbf{P}} \left(\exp(\xi) u_N \frac{\text{He}_m(\xi)}{\sqrt{m!}} \right) = \mathbf{E}_{\mathbf{P}} \left(\exp(\xi) |\xi - 1| \frac{\text{He}_m(\xi)}{\sqrt{m!}} \right) \quad \text{for all } m = 0, \dots, N \quad (\text{A.1})$$

is equal to the stochastic part \hat{u}_N of the stochastic Galerkin solution $\hat{u}_{N, hp}$ of Example 1. Since $u_N \in U_N$ satisfies (A.1) there holds

$$\mathbf{E}_{\mathbf{P}} (\kappa u_N v) - \mathbf{E}_{\mathbf{P}} (f v) = \mathbf{E}_{\mathbf{P}} (\kappa (u_N - u) v) = 0 \quad \text{for all } v \in U_N.$$

Thus, the stochastic Galerkin solution u_N is the best approximation to the exact solution $u \in L^2(\Omega, \sigma(\xi), \kappa d\mathbf{P})$ in the subspace U_N . In other words, u_N is the orthogonal projection of u onto U_N in $L^2(\Omega, \sigma(\xi), \kappa d\mathbf{P})$, i. e. with respect to the weighted inner product $\langle v_1, v_2 \rangle_\kappa := \mathbf{E}_{\mathbf{P}} (\kappa v_1 v_2)$. To prove that the solution u_N does not converge in quadratic mean to the exact solution u we consider the orthonormal basis $\{\tilde{p}_n(\xi), n \in \mathbb{N}_0\}$ of the weighted space $L^2(\Omega, \sigma(\xi), \exp(\xi) d\mathbf{P})$ which is given by

$$\tilde{p}_n(\xi) := \frac{e^{-1/4} \text{He}_n(\xi - 1)}{\sqrt{n!}}, \quad n \in \mathbb{N}_0.$$

Hence, the stochastic Galerkin solution u_N can be written as

$$u_N = \sum_{n=0}^N \langle u, \tilde{p}_n(\xi) \rangle_\kappa \tilde{p}_n(\xi).$$

Assume that the polynomial chaos coefficients $c_i^{(n)}, i = 0, \dots, n, n \in \mathbb{N}_0$, in the expansions

$$\tilde{p}_n(\xi) = \sum_{i=0}^n c_i^{(n)} \frac{\text{He}_i(\xi)}{\sqrt{i!}},$$

are known, then the stochastic Galerkin solution of order N with respect to the Hermite polynomials takes on the form

$$u_N = \sum_{n=0}^N \langle u, \tilde{p}_n(\xi) \rangle_\kappa \tilde{p}_n(\xi) = \sum_{n=0}^N \langle u, \tilde{p}_n(\xi) \rangle_\kappa \sum_{i=0}^n c_i^{(n)} \frac{\text{He}_i(\xi)}{\sqrt{i!}} = \sum_{i=0}^N \frac{\text{He}_i(\xi)}{\sqrt{i!}} \sum_{n=i}^N \langle u, \tilde{p}_n(\xi) \rangle_\kappa c_i^{(n)}.$$

If the stochastic Galerkin solutions u_N converge to the exact solution, their second order moments must converge to the exact solution's second order moment since

$$\left| \|u\|_{L^2(\Omega, \mathfrak{A}, \mathbf{P})} - \|u_N\|_{L^2(\Omega, \mathfrak{A}, \mathbf{P})} \right| \leq \|u - u_N\|_{L^2(\Omega, \mathfrak{A}, \mathbf{P})} \rightarrow 0, \quad N \rightarrow \infty,$$

i. e., there must hold

$$\|u_N\|_{L^2(\Omega, \mathfrak{A}, \mathbf{P})}^2 = \mathbf{E}_{\mathbf{P}} (u_N)^2 = \sum_{i=0}^N \left(\sum_{n=i}^N \langle u, \tilde{p}_n(\xi) \rangle_\kappa c_i^{(n)} \right)^2 \rightarrow \|u\|_{L^2(\Omega, \mathfrak{A}, \mathbf{P})}^2 = \mathbf{E}_{\mathbf{P}} u^2 < \infty, \quad N \rightarrow \infty.$$

However, we show that there exists a subsequence of stochastic Galerkin solutions $(u_N)_{N \in \mathbb{N}}$ where the second order moments diverge to infinity.

To see this, we evaluate the generalized polynomial chaos coefficients $\langle u, \tilde{p}_n(\xi) \rangle_\kappa$ for $n \in \mathbb{N}_0$ by the change of variables $y = x - 1$ yielding

$$\begin{aligned} \langle u, \tilde{p}_n(\xi) \rangle_\kappa &= \int_{-\infty}^{\infty} u_\xi(x) \frac{e^{-1/4} \text{He}_n(x-1)}{\sqrt{n!}} \exp(x) \frac{\exp(-\frac{1}{2}x^2)}{\sqrt{2\pi}} dx \\ &= e^{1/4} \int_{-\infty}^{\infty} u_\xi(x) \frac{\text{He}_n(x-1)}{\sqrt{n!}} \frac{\exp(-\frac{(x-1)^2}{2})}{\sqrt{2\pi}} dx \\ &= e^{1/4} \int_{-\infty}^{\infty} u_\xi(y+1) \frac{\text{He}_n(y)}{\sqrt{n!}} \frac{\exp(-\frac{y^2}{2})}{\sqrt{2\pi}} dy = e^{1/4} \mathbf{E}_{\mathbf{P}} \left(u_\xi(\xi+1) \frac{\text{He}_n(\xi)}{\sqrt{n!}} \right) \\ &= e^{1/4} \mathbf{E}_{\mathbf{P}} \left(|\xi| \frac{\text{He}_n(\xi)}{\sqrt{n!}} \right) = \begin{cases} e^{1/4} \frac{(-1)^{n/2+1} \sqrt{n!}}{\sqrt{2\pi} 2^{n/2-1} (\frac{n}{2})! (n-1)}, & n \text{ even.} \\ 0, & \text{otherwise.} \end{cases} \end{aligned}$$

Expanding the polynomials $\tilde{p}_n(\xi)$ in polynomial chaos leads us to the sums

$$\tilde{p}_n(\xi) = \frac{e^{-1/4} \text{He}_n(\xi-1)}{\sqrt{n!}} = \sum_{i=0}^n c_i^{(n)} \frac{\text{He}_i(\xi)}{\sqrt{i!}}$$

where the coefficients are given by

$$c_i^{(n)} = e^{-1/4} \frac{(-1)^{n-i} \sqrt{n!}}{(n-i)! \sqrt{i!}}$$

due to Formula 22.12.8 in [1]. Combining these two results we obtain

$$\begin{aligned} \langle u, \tilde{p}_n(\xi) \rangle_\kappa c_i^{(n)} &= \begin{cases} e^{1/4} \frac{(-1)^{n/2+1} \sqrt{n!}}{\sqrt{2\pi} 2^{n/2-1} (\frac{n}{2})! (n-1)} e^{-1/4} \frac{(-1)^{n-i} \sqrt{n!}}{(n-i)! \sqrt{i!}}, & n \text{ even,} \\ 0, & \text{otherwise,} \end{cases} \\ &= \begin{cases} \frac{(-1)^{3n/2+1-i} n!}{\sqrt{2\pi} 2^{n/2-1} (\frac{n}{2})! (n-1) (n-i)! \sqrt{i!}}, & n \text{ even,} \\ 0, & \text{otherwise.} \end{cases} \end{aligned}$$

Now, we investigate the second order moments of the subsequence $(u_{2N})_{N \in \mathbb{N}_0}$ of the stochastic Galerkin solutions, that is,

$$\mathbf{E}_{\mathbf{P}}(u_{2N})^2 = \sum_{i=0}^{2N} \left[\sum_{n=i}^{2N} \langle u, \tilde{p}_n(\xi) \rangle_{\kappa} c_i^{(n)} \right]^2.$$

In order to show that this sequence goes to infinity for $N \rightarrow \infty$ we consider a single term in this sum, namely the summand for $i = 2N - 3$:

$$\begin{aligned} & \left[\sum_{n=2N-3}^{2N} \langle u, \tilde{p}_{2n}(\xi) \rangle_{\kappa} c_i^{(n)} \right]^2 \\ &= \left[\langle u, \tilde{p}_{2N-2}(\xi) \rangle_{\kappa} c_{2N-3}^{(2N-2)} + \langle u, \tilde{p}_{2N}(\xi) \rangle_{\kappa} c_{2N-3}^{(2N)} \right]^2 \\ &= \left[\frac{(-1)^{N+1}(2N-2)!}{\sqrt{2\pi}2^{N-2}(N-1)!(2N-3)\sqrt{(2N-3)!}} + \frac{(-1)^N(2N)!}{\sqrt{2\pi}2^{N-1}N!(2N-1)3!\sqrt{(2N-3)!}} \right]^2 \\ &= \left[\frac{(2N-2)!}{\sqrt{2\pi}2^{N-1}(N-1)!\sqrt{(2N-3)!}} \left(\frac{(-1)^{N+1}2}{2N-3} + \frac{(-1)^N}{3} \right) \right]^2 \\ &= \frac{(2N-2)!^2}{2\pi 2^{2N-2}(N-1)!^2(2N-3)!} \left(\frac{4}{(2N-3)^2} - \frac{4}{3(2N-3)} + \frac{1}{9} \right) \\ &= \frac{(2N-2)!2^{-(2N-2)}}{\pi(N-1)!^2} \left(\frac{4(N-1)}{(2N-3)^2} - \frac{4(N-1)}{3(2N-3)} + \frac{N-1}{9} \right). \end{aligned}$$

Then, applying the Stirling's formula (see e. g. [1]) yields

$$\underbrace{\frac{(2N-2)!2^{-(2N-2)}}{\pi(N-1)!^2}}_{\simeq \frac{1}{\sqrt{N}}} \underbrace{\left(\frac{4(N-1)}{(2N-3)^2} - \frac{4(N-1)}{3(2N-3)} + \frac{N-1}{9} \right)}_{\simeq N} \simeq \sqrt{N}$$

when $N \rightarrow \infty$ and thus we obtain

$$\begin{aligned} \mathbf{E}_{\mathbf{P}}(u_{2N})^2 &= \sum_{i=0}^{2N} \left[\sum_{n=i}^{2N} \langle u, \tilde{p}_n(\xi) \rangle_{\kappa} c_i^{(n)} \right]^2 \\ &\geq \frac{(2N-2)!2^{-(2N-2)}}{\pi(N-1)!^2} \left(\frac{4(N-1)}{(2N-3)^2} - \frac{4(N-1)}{3(2N-3)} + \frac{N-1}{9} \right) \rightarrow \infty \end{aligned}$$

for $N \rightarrow \infty$. This implies that the sequence of stochastic Galerkin solutions $(u_N)_{N \in \mathbb{N}_0}$ fails to converge in quadratic mean to the exact solution u .

References

- [1] M. Abramowitz and I. A. Stegun, *Handbook of Mathematical Functions*, DOVER Publications, Inc, New York, 1965.
- [2] I. Babuška, F. Nobile, and R. Tempone, *A Stochastic Collocation Method for Elliptic Partial Differential Equations with Random Input Data*, SIAM J. Numer. Anal. **45** (2007), no. 3, 1005–1034.

- [3] I. Babuška, R. Tempone, and G. E. Zouraris, *Galerkin Finite Element Approximations of Stochastic Elliptic Partial Differential Equations*, SIAM J. Numer. Anal. **42** (2004), no. 2, 800–825.
- [4] ———, *Solving elliptic boundary value problems with uncertain coefficients by the finite element method: the stochastic formulation*, Comput. Methods Appl. Mech. Engrg. **194** (2005), 1251–1294.
- [5] M. Bieri, R. Andreev, and C. Schwab, *Sparse tensor discretization of elliptic SPDEs*, SIAM J. Sci. Comput. **31** (2009), no. 6, 4281–4304.
- [6] M. Bieri and C. Schwab, *Sparse high order FEM for elliptic sPDEs*, Comput. Methods Appl. Mech. Engrg. **198** (2009), no. 13-14, 1149–1170.
- [7] A. Bobrowski, *Functional Analysis for Probability and Stochastic Processes*, Cambridge University Press, Cambridge UK, 2005.
- [8] S. C. Brenner and L. R. Scott, *The Mathematical Theory of Finite Element Methods*, 2nd ed., Texts in Applied Mathematics, vol. 15, Springer-Verlag, New York, 2002.
- [9] C. Canuto, M. Y. Hussaini, A. Quarteroni, and T. A. Zang, *Spectral Methods: Fundamentals in Single Domains*, Springer-Verlag, Berlin Heidelberg, 2006.
- [10] A. Cohen, R. DeVore, and C. Schwab, *Convergence Rates of Best N -term Galerkin Approximations for a Class of Elliptic sPDEs*, Foundations of Computational Mathematics **10** (2010), no. 6, 615–646.
- [11] M. K. Deb, I. Babuška, and J. T. Oden, *Solution of stochastic partial differential equations using Galerkin finite element techniques*, Comput. Methods Appl. Mech. Engrg. **190** (2001), 6359–6372.
- [12] M. Eiermann, O. G. Ernst, and E. Ullmann, *Computational aspects of the stochastic finite element method*, Computing and Visualization in Science **10** (2007), no. 1, 3–15.
- [13] O. Ernst, A. Mugler, E. Ullmann, and H. J. Starkloff, *On the convergence of generalized polynomial chaos*, ESAIM: Mathematical Modelling and Numerical Analysis **46** (2012), no. 2, 317–339.
- [14] R. V. Field and M. Grigoriu, *Convergence Properties of Polynomial Chaos Approximations for L^2 -Random Variables*, Sandia Report SAND2007-1262, 2007.
- [15] P. Frauenfelder, C. Schwab, and R. A. Todor, *Finite elements for elliptic problems with stochastic coefficients*, Computer Methods in Applied Mechanics and Engineering **194** (2005), no. 2-5, 205–228.
- [16] R. A. Freeze, *A Stochastic-Conceptual Analysis of One-Dimensional Groundwater Flow in Nonuniform Homogeneous Media*, Water Resources Research (1975), 725–741.
- [17] J. Galvis and M. Sarkis, *Approximating infinity-dimensional stochastic Darcy's Equations without uniform ellipticity*, SIAM J. Numer. Anal. **47** (2009), no. 5, 3624–3651.
- [18] R. Ghanem, *Ingredients for a general purpose stochastic finite elements implementation*, Comput. Methods Appl. Mech. Engrg. **168** (1999), 19–34.

- [19] ———, *Stochastic Finite Elements with Multiple Random Non-Gaussian Properties*, Journal of Engineering Mechanics **125** (1999), no. 1, 26–40.
- [20] R. Ghanem and S. Dham, *Stochastic Finite Element Analysis for Multiphase Flow in Heterogeneous Porous Media*, Transport in Porous Media **32** (1998), 239–262.
- [21] R. Ghanem and P. D. Spanos, *Stochastic Finite Elements: A Spectral Approach*, Springer-Verlag, New York, 1991.
- [22] C. J. Gittelsohn, *Stochastic Galerkin discretization of the log-normal isotropic diffusion problem*, Math. Models Methods Appl. Sci. **20** (2010), no. 2, 237–263.
- [23] E. Godoy, A. Ronveaux, A. Zarzo, and I. Area, *Minimal recurrence relations for connection coefficients between classical orthogonal polynomials: Continuous case*, Journal of Computational and Applied Mathematics **84** (1997), 257–275.
- [24] E. Hille and R. S. Phillips, *Functional Analysis and Semi-Groups*, Colloquium Publications, vol. 31, American Mathematical Society, 1957.
- [25] O. Kallenberg, *Foundations of modern probability*, Springer, Berlin et al., 2001.
- [26] O. P. Le Maître and O. M. Knio, *Spectral Methods for Uncertainty Quantification. Scientific Computation: With Applications to Computational Fluid Dynamics*, 1 ed., Springer-Verlag, s.l, 2010.
- [27] M. Loève, *Probability Theory II. 4th Edition*, Springer-Verlag, New York, Heidelberg, Berlin, 1978.
- [28] D. Lucor, C. H. Su, and G. E. Karniadakis, *Generalized polynomial chaos and random oscillators*, Int. J. Numer. Meth. Engng. **60** (2004), 571–596.
- [29] P. Maroni and Z. Rocha, *Connection coefficients between orthogonal polynomials and the canonical sequence: an approach based on symbolic computation*, Numer. Algor. (Numerical Algorithms) **47** (2008), no. 3, 291–314.
- [30] A. Mugler and H. J. Starkloff, *On elliptic partial differential equations with random coefficients*, Stud. Univ. Babeş-Bolyai Math. **56** (2011), no. 2, 473–487.
- [31] A. Narayan and J. Hesthaven, *Computation of connection coefficients and measure modifications for orthogonal polynomials*, BIT Numerical Mathematics (2011), 1–27, 10.1007/s10543-011-0363-z.
- [32] W. Nowak, *Geostatistical Methods for the Identification of Flow and Transport Parameters in the Subsurface*, Ph.D. thesis, Universität Stuttgart, 2005.
- [33] C. Schwab and C. J. Gittelsohn, *Sparse tensor discretizations of high-dimensional parametric and stochastic PDEs*, Acta Numerica **20** (2011), 291–467.
- [34] R. A. Todor and C. Schwab, *Convergence rates for sparse chaos approximations of elliptic problems with stochastic coefficients*, IMA Journal of Numerical Analysis **27** (2007), 232–261.
- [35] H. Umegaki and A. T. Bharucha-Reid, *Banach Space-Valued Random Variables and Tensor Products of Banach Spaces*, Journal of Mathematical Analysis and Applications **31** (1970), 49–67.

- [36] X. Wan and G. E. Karniadakis, *Beyond Wiener Askey Expansions: Handling Arbitrary PDFs*, Journal of Scientific Computing **27** (2006), 455–464.
- [37] N. Wiener, *Homogeneous Chaos*, American Journal of Mathematics **60** (1938), no. 4, 897–936.
- [38] D. Xiu, *Numerical methods for stochastic computations: A spectral method approach*, Princeton Univ. Press, Princeton and NJ, 2010.
- [39] D. Xiu and G. E. Karniadakis, *Modeling uncertainty in steady state diffusion problems via generalized polynomial chaos*, Comput. Methods Appl. Mech. Engrg. **191** (2002), no. 43, 4927–4948.
- [40] ———, *The Wiener-Askey Polynomial Chaos for Stochastic Differential Equations*, SIAM J. Sci. Comput. **24** (2002), no. 2, 619–644.
- [41] ———, *A new stochastic approach to transient heat conduction modeling with uncertainty*, International Journal of Heat and Mass Transfer **46** (2003), no. 24, 4681–4693.
- [42] ———, *Modeling uncertainty in flow simulations via generalized polynomial chaos*, Journal of Computational Physics **187** (2003), 137–167.
- [43] D. Xiu, D. Lucor, C. H. Su, and G. E. Karniadakis, *Performance Evaluation of Generalized Polynomial Chaos*, Computational Science — ICCS 2003 (Peter M.A. Sloot, David Abramson, Alexander V. Bogdanov, Jack J. Dongarra, Albert Y. Zomaya, and Yuriy E. Gorbachev, eds.), Lecture Notes in Computer Science, vol. 2660, Springer-Verlag, 2003.
- [44] D. Zhang, *Stochastic Methods for Flow in Porous Media. Coping with Uncertainties*, Academic Press, San Diego, CA, 2002.

Preprint Series DFG-SPP 1324

<http://www.dfg-spp1324.de>

Reports

- [1] R. Ramlau, G. Teschke, and M. Zhariy. A Compressive Landweber Iteration for Solving Ill-Posed Inverse Problems. Preprint 1, DFG-SPP 1324, September 2008.
- [2] G. Plonka. The Easy Path Wavelet Transform: A New Adaptive Wavelet Transform for Sparse Representation of Two-dimensional Data. Preprint 2, DFG-SPP 1324, September 2008.
- [3] E. Novak and H. Woźniakowski. Optimal Order of Convergence and (In-) Tractability of Multivariate Approximation of Smooth Functions. Preprint 3, DFG-SPP 1324, October 2008.
- [4] M. Espig, L. Grasedyck, and W. Hackbusch. Black Box Low Tensor Rank Approximation Using Fibre-Crosses. Preprint 4, DFG-SPP 1324, October 2008.
- [5] T. Bonesky, S. Dahlke, P. Maass, and T. Raasch. Adaptive Wavelet Methods and Sparsity Reconstruction for Inverse Heat Conduction Problems. Preprint 5, DFG-SPP 1324, January 2009.
- [6] E. Novak and H. Woźniakowski. Approximation of Infinitely Differentiable Multivariate Functions Is Intractable. Preprint 6, DFG-SPP 1324, January 2009.
- [7] J. Ma and G. Plonka. A Review of Curvelets and Recent Applications. Preprint 7, DFG-SPP 1324, February 2009.
- [8] L. Denis, D. A. Lorenz, and D. Tiede. Greedy Solution of Ill-Posed Problems: Error Bounds and Exact Inversion. Preprint 8, DFG-SPP 1324, April 2009.
- [9] U. Friedrich. A Two Parameter Generalization of Lions' Nonoverlapping Domain Decomposition Method for Linear Elliptic PDEs. Preprint 9, DFG-SPP 1324, April 2009.
- [10] K. Bredies and D. A. Lorenz. Minimization of Non-smooth, Non-convex Functionals by Iterative Thresholding. Preprint 10, DFG-SPP 1324, April 2009.
- [11] K. Bredies and D. A. Lorenz. Regularization with Non-convex Separable Constraints. Preprint 11, DFG-SPP 1324, April 2009.

- [12] M. Döhler, S. Kunis, and D. Potts. Nonequispaced Hyperbolic Cross Fast Fourier Transform. Preprint 12, DFG-SPP 1324, April 2009.
- [13] C. Bender. Dual Pricing of Multi-Exercise Options under Volume Constraints. Preprint 13, DFG-SPP 1324, April 2009.
- [14] T. Müller-Gronbach and K. Ritter. Variable Subspace Sampling and Multi-level Algorithms. Preprint 14, DFG-SPP 1324, May 2009.
- [15] G. Plonka, S. Tenorth, and A. Iske. Optimally Sparse Image Representation by the Easy Path Wavelet Transform. Preprint 15, DFG-SPP 1324, May 2009.
- [16] S. Dahlke, E. Novak, and W. Sickel. Optimal Approximation of Elliptic Problems by Linear and Nonlinear Mappings IV: Errors in L_2 and Other Norms. Preprint 16, DFG-SPP 1324, June 2009.
- [17] B. Jin, T. Khan, P. Maass, and M. Pidcock. Function Spaces and Optimal Currents in Impedance Tomography. Preprint 17, DFG-SPP 1324, June 2009.
- [18] G. Plonka and J. Ma. Curvelet-Wavelet Regularized Split Bregman Iteration for Compressed Sensing. Preprint 18, DFG-SPP 1324, June 2009.
- [19] G. Teschke and C. Borries. Accelerated Projected Steepest Descent Method for Nonlinear Inverse Problems with Sparsity Constraints. Preprint 19, DFG-SPP 1324, July 2009.
- [20] L. Grasedyck. Hierarchical Singular Value Decomposition of Tensors. Preprint 20, DFG-SPP 1324, July 2009.
- [21] D. Rudolf. Error Bounds for Computing the Expectation by Markov Chain Monte Carlo. Preprint 21, DFG-SPP 1324, July 2009.
- [22] M. Hansen and W. Sickel. Best m-term Approximation and Lizorkin-Triebel Spaces. Preprint 22, DFG-SPP 1324, August 2009.
- [23] F.J. Hickernell, T. Müller-Gronbach, B. Niu, and K. Ritter. Multi-level Monte Carlo Algorithms for Infinite-dimensional Integration on \mathbb{R}^N . Preprint 23, DFG-SPP 1324, August 2009.
- [24] S. Dereich and F. Heidenreich. A Multilevel Monte Carlo Algorithm for Lévy Driven Stochastic Differential Equations. Preprint 24, DFG-SPP 1324, August 2009.
- [25] S. Dahlke, M. Fornasier, and T. Raasch. Multilevel Preconditioning for Adaptive Sparse Optimization. Preprint 25, DFG-SPP 1324, August 2009.

- [26] S. Dereich. Multilevel Monte Carlo Algorithms for Lévy-driven SDEs with Gaussian Correction. Preprint 26, DFG-SPP 1324, August 2009.
- [27] G. Plonka, S. Tenorth, and D. Roşca. A New Hybrid Method for Image Approximation using the Easy Path Wavelet Transform. Preprint 27, DFG-SPP 1324, October 2009.
- [28] O. Koch and C. Lubich. Dynamical Low-rank Approximation of Tensors. Preprint 28, DFG-SPP 1324, November 2009.
- [29] E. Faou, V. Gradinaru, and C. Lubich. Computing Semi-classical Quantum Dynamics with Hagedorn Wavepackets. Preprint 29, DFG-SPP 1324, November 2009.
- [30] D. Conte and C. Lubich. An Error Analysis of the Multi-configuration Time-dependent Hartree Method of Quantum Dynamics. Preprint 30, DFG-SPP 1324, November 2009.
- [31] C. E. Powell and E. Ullmann. Preconditioning Stochastic Galerkin Saddle Point Problems. Preprint 31, DFG-SPP 1324, November 2009.
- [32] O. G. Ernst and E. Ullmann. Stochastic Galerkin Matrices. Preprint 32, DFG-SPP 1324, November 2009.
- [33] F. Lindner and R. L. Schilling. Weak Order for the Discretization of the Stochastic Heat Equation Driven by Impulsive Noise. Preprint 33, DFG-SPP 1324, November 2009.
- [34] L. Kämmerer and S. Kunis. On the Stability of the Hyperbolic Cross Discrete Fourier Transform. Preprint 34, DFG-SPP 1324, December 2009.
- [35] P. Cerejeiras, M. Ferreira, U. Kähler, and G. Teschke. Inversion of the noisy Radon transform on $SO(3)$ by Gabor frames and sparse recovery principles. Preprint 35, DFG-SPP 1324, January 2010.
- [36] T. Jahnke and T. Udrescu. Solving Chemical Master Equations by Adaptive Wavelet Compression. Preprint 36, DFG-SPP 1324, January 2010.
- [37] P. Kittipoom, G. Kutyniok, and W.-Q Lim. Irregular Shearlet Frames: Geometry and Approximation Properties. Preprint 37, DFG-SPP 1324, February 2010.
- [38] G. Kutyniok and W.-Q Lim. Compactly Supported Shearlets are Optimally Sparse. Preprint 38, DFG-SPP 1324, February 2010.

- [39] M. Hansen and W. Sickel. Best m -Term Approximation and Tensor Products of Sobolev and Besov Spaces – the Case of Non-compact Embeddings. Preprint 39, DFG-SPP 1324, March 2010.
- [40] B. Niu, F.J. Hickernell, T. Müller-Gronbach, and K. Ritter. Deterministic Multi-level Algorithms for Infinite-dimensional Integration on $\mathbb{R}^{\mathbb{N}}$. Preprint 40, DFG-SPP 1324, March 2010.
- [41] P. Kittipoom, G. Kutyniok, and W.-Q Lim. Construction of Compactly Supported Shearlet Frames. Preprint 41, DFG-SPP 1324, March 2010.
- [42] C. Bender and J. Steiner. Error Criteria for Numerical Solutions of Backward SDEs. Preprint 42, DFG-SPP 1324, April 2010.
- [43] L. Grasedyck. Polynomial Approximation in Hierarchical Tucker Format by Vector-Tensorization. Preprint 43, DFG-SPP 1324, April 2010.
- [44] M. Hansen und W. Sickel. Best m -Term Approximation and Sobolev-Besov Spaces of Dominating Mixed Smoothness - the Case of Compact Embeddings. Preprint 44, DFG-SPP 1324, April 2010.
- [45] P. Binev, W. Dahmen, and P. Lamby. Fast High-Dimensional Approximation with Sparse Occupancy Trees. Preprint 45, DFG-SPP 1324, May 2010.
- [46] J. Ballani and L. Grasedyck. A Projection Method to Solve Linear Systems in Tensor Format. Preprint 46, DFG-SPP 1324, May 2010.
- [47] P. Binev, A. Cohen, W. Dahmen, R. DeVore, G. Petrova, and P. Wojtaszczyk. Convergence Rates for Greedy Algorithms in Reduced Basis Methods. Preprint 47, DFG-SPP 1324, May 2010.
- [48] S. Kestler and K. Urban. Adaptive Wavelet Methods on Unbounded Domains. Preprint 48, DFG-SPP 1324, June 2010.
- [49] H. Yserentant. The Mixed Regularity of Electronic Wave Functions Multiplied by Explicit Correlation Factors. Preprint 49, DFG-SPP 1324, June 2010.
- [50] H. Yserentant. On the Complexity of the Electronic Schrödinger Equation. Preprint 50, DFG-SPP 1324, June 2010.
- [51] M. Guillemard and A. Iske. Curvature Analysis of Frequency Modulated Manifolds in Dimensionality Reduction. Preprint 51, DFG-SPP 1324, June 2010.
- [52] E. Herrholz and G. Teschke. Compressive Sensing Principles and Iterative Sparse Recovery for Inverse and Ill-Posed Problems. Preprint 52, DFG-SPP 1324, July 2010.

- [53] L. Kämmerer, S. Kunis, and D. Potts. Interpolation Lattices for Hyperbolic Cross Trigonometric Polynomials. Preprint 53, DFG-SPP 1324, July 2010.
- [54] G. Kutyniok and W.-Q Lim. Shearlets on Bounded Domains. Preprint 54, DFG-SPP 1324, July 2010.
- [55] A. Zeiser. Wavelet Approximation in Weighted Sobolev Spaces of Mixed Order with Applications to the Electronic Schrödinger Equation. Preprint 55, DFG-SPP 1324, July 2010.
- [56] G. Kutyniok, J. Lemvig, and W.-Q Lim. Compactly Supported Shearlets. Preprint 56, DFG-SPP 1324, July 2010.
- [57] A. Zeiser. On the Optimality of the Inexact Inverse Iteration Coupled with Adaptive Finite Element Methods. Preprint 57, DFG-SPP 1324, July 2010.
- [58] S. Jokar. Sparse Recovery and Kronecker Products. Preprint 58, DFG-SPP 1324, August 2010.
- [59] T. Aboiyar, E. H. Georgoulis, and A. Iske. Adaptive ADER Methods Using Kernel-Based Polyharmonic Spline WENO Reconstruction. Preprint 59, DFG-SPP 1324, August 2010.
- [60] O. G. Ernst, A. Mugler, H.-J. Starkloff, and E. Ullmann. On the Convergence of Generalized Polynomial Chaos Expansions. Preprint 60, DFG-SPP 1324, August 2010.
- [61] S. Holtz, T. Rohwedder, and R. Schneider. On Manifolds of Tensors of Fixed TT-Rank. Preprint 61, DFG-SPP 1324, September 2010.
- [62] J. Ballani, L. Grasedyck, and M. Kluge. Black Box Approximation of Tensors in Hierarchical Tucker Format. Preprint 62, DFG-SPP 1324, October 2010.
- [63] M. Hansen. On Tensor Products of Quasi-Banach Spaces. Preprint 63, DFG-SPP 1324, October 2010.
- [64] S. Dahlke, G. Steidl, and G. Teschke. Shearlet Coorbit Spaces: Compactly Supported Analyzing Shearlets, Traces and Embeddings. Preprint 64, DFG-SPP 1324, October 2010.
- [65] W. Hackbusch. Tensorisation of Vectors and their Efficient Convolution. Preprint 65, DFG-SPP 1324, November 2010.
- [66] P. A. Cioica, S. Dahlke, S. Kinzel, F. Lindner, T. Raasch, K. Ritter, and R. L. Schilling. Spatial Besov Regularity for Stochastic Partial Differential Equations on Lipschitz Domains. Preprint 66, DFG-SPP 1324, November 2010.

- [67] E. Novak and H. Woźniakowski. On the Power of Function Values for the Approximation Problem in Various Settings. Preprint 67, DFG-SPP 1324, November 2010.
- [68] A. Hinrichs, E. Novak, and H. Woźniakowski. The Curse of Dimensionality for Monotone and Convex Functions of Many Variables. Preprint 68, DFG-SPP 1324, November 2010.
- [69] G. Kutyniok and W.-Q. Lim. Image Separation Using Shearlets. Preprint 69, DFG-SPP 1324, November 2010.
- [70] B. Jin and P. Maass. An Analysis of Electrical Impedance Tomography with Applications to Tikhonov Regularization. Preprint 70, DFG-SPP 1324, December 2010.
- [71] S. Holtz, T. Rohwedder, and R. Schneider. The Alternating Linear Scheme for Tensor Optimisation in the TT Format. Preprint 71, DFG-SPP 1324, December 2010.
- [72] T. Müller-Gronbach and K. Ritter. A Local Refinement Strategy for Constructive Quantization of Scalar SDEs. Preprint 72, DFG-SPP 1324, December 2010.
- [73] T. Rohwedder and R. Schneider. An Analysis for the DIIS Acceleration Method used in Quantum Chemistry Calculations. Preprint 73, DFG-SPP 1324, December 2010.
- [74] C. Bender and J. Steiner. Least-Squares Monte Carlo for Backward SDEs. Preprint 74, DFG-SPP 1324, December 2010.
- [75] C. Bender. Primal and Dual Pricing of Multiple Exercise Options in Continuous Time. Preprint 75, DFG-SPP 1324, December 2010.
- [76] H. Harbrecht, M. Peters, and R. Schneider. On the Low-rank Approximation by the Pivoted Cholesky Decomposition. Preprint 76, DFG-SPP 1324, December 2010.
- [77] P. A. Cioica, S. Dahlke, N. Döhring, S. Kinzel, F. Lindner, T. Raasch, K. Ritter, and R. L. Schilling. Adaptive Wavelet Methods for Elliptic Stochastic Partial Differential Equations. Preprint 77, DFG-SPP 1324, January 2011.
- [78] G. Plonka, S. Tenorth, and A. Iske. Optimal Representation of Piecewise Hölder Smooth Bivariate Functions by the Easy Path Wavelet Transform. Preprint 78, DFG-SPP 1324, January 2011.

- [79] A. Mugler and H.-J. Starkloff. On Elliptic Partial Differential Equations with Random Coefficients. Preprint 79, DFG-SPP 1324, January 2011.
- [80] T. Müller-Gronbach, K. Ritter, and L. Yaroslavtseva. A Derandomization of the Euler Scheme for Scalar Stochastic Differential Equations. Preprint 80, DFG-SPP 1324, January 2011.
- [81] W. Dahmen, C. Huang, C. Schwab, and G. Welper. Adaptive Petrov-Galerkin methods for first order transport equations. Preprint 81, DFG-SPP 1324, January 2011.
- [82] K. Grella and C. Schwab. Sparse Tensor Spherical Harmonics Approximation in Radiative Transfer. Preprint 82, DFG-SPP 1324, January 2011.
- [83] D.A. Lorenz, S. Schiffler, and D. Tiede. Beyond Convergence Rates: Exact Inversion With Tikhonov Regularization With Sparsity Constraints. Preprint 83, DFG-SPP 1324, January 2011.
- [84] S. Dereich, M. Scheutzow, and R. Schottstedt. Constructive quantization: Approximation by empirical measures. Preprint 84, DFG-SPP 1324, January 2011.
- [85] S. Dahlke and W. Sickel. On Besov Regularity of Solutions to Nonlinear Elliptic Partial Differential Equations. Preprint 85, DFG-SPP 1324, January 2011.
- [86] S. Dahlke, U. Friedrich, P. Maass, T. Raasch, and R.A. Ressel. An adaptive wavelet method for parameter identification problems in parabolic partial differential equations. Preprint 86, DFG-SPP 1324, January 2011.
- [87] A. Cohen, W. Dahmen, and G. Welper. Adaptivity and Variational Stabilization for Convection-Diffusion Equations. Preprint 87, DFG-SPP 1324, January 2011.
- [88] T. Jahnke. On Reduced Models for the Chemical Master Equation. Preprint 88, DFG-SPP 1324, January 2011.
- [89] P. Binev, W. Dahmen, R. DeVore, P. Lamby, D. Savu, and R. Sharpley. Compressed Sensing and Electron Microscopy. Preprint 89, DFG-SPP 1324, March 2011.
- [90] P. Binev, F. Blanco-Silva, D. Blom, W. Dahmen, P. Lamby, R. Sharpley, and T. Vogt. High Quality Image Formation by Nonlocal Means Applied to High-Angle Annular Dark Field Scanning Transmission Electron Microscopy (HAADF-STEM). Preprint 90, DFG-SPP 1324, March 2011.
- [91] R. A. Ressel. A Parameter Identification Problem for a Nonlinear Parabolic Differential Equation. Preprint 91, DFG-SPP 1324, May 2011.

- [92] G. Kutyniok. Data Separation by Sparse Representations. Preprint 92, DFG-SPP 1324, May 2011.
- [93] M. A. Davenport, M. F. Duarte, Y. C. Eldar, and G. Kutyniok. Introduction to Compressed Sensing. Preprint 93, DFG-SPP 1324, May 2011.
- [94] H.-C. Kreuzler and H. Yserentant. The Mixed Regularity of Electronic Wave Functions in Fractional Order and Weighted Sobolev Spaces. Preprint 94, DFG-SPP 1324, June 2011.
- [95] E. Ullmann, H. C. Elman, and O. G. Ernst. Efficient Iterative Solvers for Stochastic Galerkin Discretizations of Log-Transformed Random Diffusion Problems. Preprint 95, DFG-SPP 1324, June 2011.
- [96] S. Kunis and I. Melzer. On the Butterfly Sparse Fourier Transform. Preprint 96, DFG-SPP 1324, June 2011.
- [97] T. Rohwedder. The Continuous Coupled Cluster Formulation for the Electronic Schrödinger Equation. Preprint 97, DFG-SPP 1324, June 2011.
- [98] T. Rohwedder and R. Schneider. Error Estimates for the Coupled Cluster Method. Preprint 98, DFG-SPP 1324, June 2011.
- [99] P. A. Cioica and S. Dahlke. Spatial Besov Regularity for Semilinear Stochastic Partial Differential Equations on Bounded Lipschitz Domains. Preprint 99, DFG-SPP 1324, July 2011.
- [100] L. Grasedyck and W. Hackbusch. An Introduction to Hierarchical (H-) Rank and TT-Rank of Tensors with Examples. Preprint 100, DFG-SPP 1324, August 2011.
- [101] N. Chegini, S. Dahlke, U. Friedrich, and R. Stevenson. Piecewise Tensor Product Wavelet Bases by Extensions and Approximation Rates. Preprint 101, DFG-SPP 1324, September 2011.
- [102] S. Dahlke, P. Oswald, and T. Raasch. A Note on Quarkonial Systems and Multi-level Partition of Unity Methods. Preprint 102, DFG-SPP 1324, September 2011.
- [103] A. Uschmajew. Local Convergence of the Alternating Least Squares Algorithm For Canonical Tensor Approximation. Preprint 103, DFG-SPP 1324, September 2011.
- [104] S. Kvaal. Multiconfigurational time-dependent Hartree method for describing particle loss due to absorbing boundary conditions. Preprint 104, DFG-SPP 1324, September 2011.

- [105] M. Guillemard and A. Iske. On Groupoid C^* -Algebras, Persistent Homology and Time-Frequency Analysis. Preprint 105, DFG-SPP 1324, September 2011.
- [106] A. Hinrichs, E. Novak, and H. Woźniakowski. Discontinuous information in the worst case and randomized settings. Preprint 106, DFG-SPP 1324, September 2011.
- [107] M. Espig, W. Hackbusch, A. Litvinenko, H. Matthies, and E. Zander. Efficient Analysis of High Dimensional Data in Tensor Formats. Preprint 107, DFG-SPP 1324, September 2011.
- [108] M. Espig, W. Hackbusch, S. Handschuh, and R. Schneider. Optimization Problems in Contracted Tensor Networks. Preprint 108, DFG-SPP 1324, October 2011.
- [109] S. Dereich, T. Müller-Gronbach, and K. Ritter. On the Complexity of Computing Quadrature Formulas for SDEs. Preprint 109, DFG-SPP 1324, October 2011.
- [110] D. Belomestny. Solving optimal stopping problems by empirical dual optimization and penalization. Preprint 110, DFG-SPP 1324, November 2011.
- [111] D. Belomestny and J. Schoenmakers. Multilevel dual approach for pricing American style derivatives. Preprint 111, DFG-SPP 1324, November 2011.
- [112] T. Rohwedder and A. Uschmajew. Local convergence of alternating schemes for optimization of convex problems in the TT format. Preprint 112, DFG-SPP 1324, December 2011.
- [113] T. Görner, R. Hielscher, and S. Kunis. Efficient and accurate computation of spherical mean values at scattered center points. Preprint 113, DFG-SPP 1324, December 2011.
- [114] Y. Dong, T. Görner, and S. Kunis. An iterative reconstruction scheme for photoacoustic imaging. Preprint 114, DFG-SPP 1324, December 2011.
- [115] L. Kämmerer. Reconstructing hyperbolic cross trigonometric polynomials by sampling along generated sets. Preprint 115, DFG-SPP 1324, February 2012.
- [116] H. Chen and R. Schneider. Numerical analysis of augmented plane waves methods for full-potential electronic structure calculations. Preprint 116, DFG-SPP 1324, February 2012.
- [117] J. Ma, G. Plonka, and M.Y. Hussaini. Compressive Video Sampling with Approximate Message Passing Decoding. Preprint 117, DFG-SPP 1324, February 2012.

- [118] D. Heinen and G. Plonka. Wavelet shrinkage on paths for scattered data denoising. Preprint 118, DFG-SPP 1324, February 2012.
- [119] T. Jahnke and M. Kreim. Error bound for piecewise deterministic processes modeling stochastic reaction systems. Preprint 119, DFG-SPP 1324, March 2012.
- [120] C. Bender and J. Steiner. A-posteriori estimates for backward SDEs. Preprint 120, DFG-SPP 1324, April 2012.
- [121] M. Espig, W. Hackbusch, A. Litvinenko, H.G. Matthies, and P. Wöhnert. Efficient low-rank approximation of the stochastic Galerkin matrix in tensor formats. Preprint 121, DFG-SPP 1324, May 2012.
- [122] O. Bokanowski, J. Garcke, M. Griebel, and I. Klompaker. An adaptive sparse grid semi-Lagrangian scheme for first order Hamilton-Jacobi Bellman equations. Preprint 122, DFG-SPP 1324, June 2012.
- [123] A. Mugler and H.-J. Starkloff. On the convergence of the stochastic Galerkin method for random elliptic partial differential equations. Preprint 123, DFG-SPP 1324, June 2012.