


DFG-Schwerpunktprogramm 1324

„Extraktion quantifizierbarer Information aus komplexen Systemen“

On the $L_q(L_p)$ -regularity and Besov smoothness of
stochastic parabolic equations on bounded Lipschitz
domains

P.A. Cioica, K.-H. Kim, K. Lee, F. Lindner

Preprint 130



Edited by

AG Numerik/Optimierung
Fachbereich 12 - Mathematik und Informatik
Philipps-Universität Marburg
Hans-Meerwein-Str.
35032 Marburg


DFG-Schwerpunktprogramm 1324

„Extraktion quantifizierbarer Information aus komplexen Systemen“

On the $L_q(L_p)$ -regularity and Besov smoothness of
stochastic parabolic equations on bounded Lipschitz
domains

P.A. Cioica, K.-H. Kim, K. Lee, F. Lindner

Preprint 130



The consecutive numbering of the publications is determined by their chronological order.

The aim of this preprint series is to make new research rapidly available for scientific discussion. Therefore, the responsibility for the contents is solely due to the authors. The publications will be distributed by the authors.

On the $L_q(L_p)$ -regularity and Besov smoothness of stochastic parabolic equations on bounded Lipschitz domains*

Petru A. Cioica, Kyeong-Hun Kim, Kijung Lee, Felix Lindner

December 4, 2012

Abstract

We investigate the regularity of linear stochastic parabolic equations with zero Dirichlet boundary condition on bounded Lipschitz domains $\mathcal{O} \subseteq \mathbb{R}^d$ with both theoretical and numerical purpose. We use N.V. Krylov's framework of stochastic parabolic weighted Sobolev spaces $\mathfrak{H}_{p,\theta}^{\gamma,q}(\mathcal{O}, T)$. The summability parameters p and q in space and time may differ. Existence and uniqueness of solutions in these spaces is established and the Hölder regularity in time is analysed. Moreover, we prove a general embedding of weighted $L_p(\mathcal{O})$ -Sobolev spaces into the scale of Besov spaces $B_{\tau,\tau}^\alpha(\mathcal{O})$, $1/\tau = \alpha/d + 1/p$, $\alpha > 0$. This leads to a Hölder-Besov regularity result for the solution process. The regularity in this Besov scale determines the order of convergence that can be achieved by certain nonlinear approximation schemes.

Keywords: Stochastic partial differential equations, Lipschitz domain, $L_q(L_p)$ -theory, weighted Sobolev spaces, Besov spaces, quasi-Banach spaces, embedding theorems, Hölder regularity in time, nonlinear approximation, wavelets, adaptive numerical methods, square root of Laplacian operator.

AMS 2010 Subject Classification: Primary: 60H15; secondary: 46E35, 35R60.

1 Introduction

Let $\mathcal{O} \subseteq \mathbb{R}^d$ be a bounded Lipschitz domain, $T \in (0, \infty)$ and let $(w_t^k)_{t \in [0, T]}$, $k \in \mathbb{N}$, be independent one-dimensional standard Wiener processes defined on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$. We are interested in the regularity of the solutions to parabolic stochastic partial differential equations (SPDEs, for short) with zero Dirichlet boundary condition of the form

$$\left. \begin{aligned} du &= (a^{ij}u_{x^i x^j} + f)dt + (\sigma^{ik}u_{x^i} + g^k)dw_t^k && \text{on } \Omega \times [0, T] \times \mathcal{O}, \\ u &= 0 && \text{on } \Omega \times (0, T] \times \partial\mathcal{O}, \\ u(0, \cdot) &= u_0 && \text{on } \Omega \times \mathcal{O}, \end{aligned} \right\} \quad (1.1)$$

*This work has been supported by the Deutsche Forschungsgemeinschaft (DFG, grants DA 360/13-1, DA 360/13-2, DA 360/11-1, DA 360/11-2, SCHI 419/5-1, SCHI 419/5-2) and a doctoral scholarship of the Philipps-Universität Marburg. The research of the third author has been also supported by Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, Science and Technology (2011-0005597).

where the indices i and j run from 1 to d and the index k runs through $\mathbb{N} = \{1, 2, \dots\}$. Here and in the sequel we use the summation convention on the repeated indices i, j, k . The coefficients a^{ij} and σ^{ik} depend on $(\omega, t) \in \Omega \times [0, T]$. The force terms f and g^k depend on $(\omega, t, x) \in \Omega \times [0, T] \times \mathcal{O}$. By the nature of the problem, in particular by the bad contribution of the infinitesimal differences of the Wiener processes, the second spatial derivatives of the solution may blow up at the boundary $\partial\mathcal{O}$ even if the boundary is smooth, see, e.g., [28]. Hence, a natural way to deal with problems of type (1.1) is to consider u as a stochastic process with values in weighted Sobolev spaces on \mathcal{O} that allow the derivatives of functions from these spaces to blow up near the boundary. This approach has been initiated and developed by N.V. Krylov and collaborators, first as an L_2 -theory for smooth domains \mathcal{O} (see [28]), then as an L_p -theory ($p \geq 2$) for the half space ([33, 34]), for smooth domains ([23, 27]), and for general bounded domains allowing Hardy's inequality such as bounded Lipschitz domains ([26]). Existence and uniqueness of solutions have been established within specific stochastic parabolic weighted Sobolev spaces, denoted by $\mathfrak{H}_{p,\theta}^\gamma(\mathcal{O}, T)$ in [26]. These spaces consist of elements \tilde{u} of the form $d\tilde{u} = \tilde{f}dt + \tilde{g}^k dw_t^k$, where \tilde{u} , \tilde{f} and \tilde{g}^k , considered as stochastic processes with values in certain weighted $L_p(\mathcal{O})$ -Sobolev spaces, are L_p -integrable w.r.t. $\mathbb{P} \otimes dt$. We refer to Section 3 for the exact definition.

In this article we treat regularity issues concerning the solution u of problem (1.1) which arise, besides others, in the context of adaptive numerical approximation methods.

The starting point of our considerations was the question whether we can improve the Besov regularity results in [4] in time direction. In [4] the spatial regularity of u is measured in the scale of Besov spaces

$$B_{\tau,\tau}^\alpha(\mathcal{O}), \quad \frac{1}{\tau} = \frac{\alpha}{d} + \frac{1}{p}, \quad \alpha > 0, \quad (*)$$

where $p \geq 2$ is fixed. Note that for $\alpha > (p-1)d/p$ the sumability parameter τ becomes less than one, so that in this case $B_{\tau,\tau}^\alpha(\mathcal{O})$ is not a Banach space but a quasi-Banach space. It is a known result from approximation theory that the smoothness of a target function $f \in L_p(\mathcal{O})$ within the scale (*) determines the rate of convergence that can be achieved by adaptive and other nonlinear approximation methods if the approximation error is measured in $L_p(\mathcal{O})$; see [5, Chapter 4], [13] or the introduction of [4]. Based on the L_p -theory in [26], it is shown in [4] that the solution u to problem (1.1) satisfies

$$u \in L_\tau(\Omega \times [0, T], \mathcal{P}, \mathbb{P} \otimes dt; B_{\tau,\tau}^\alpha(\mathcal{O})), \quad \frac{1}{\tau} = \frac{\alpha}{d} + \frac{1}{p}, \quad (1.2)$$

for certain $\alpha > 0$ depending on the smoothness of u_0 , f and g^k , $k \in \mathbb{N}$. In general, the spatial regularity of u in the Sobolev scale $W_p^s(\mathcal{O})$, $s \geq 0$, which determines the order of convergence for uniform approximation methods in $L_p(\mathcal{O})$, is strictly less than the spatial regularity of u in the scale (*). It can be due to, e.g., the irregular behaviour of the noise at the boundary or the irregularities of the boundary itself; see [37, Chapter 4] for the latter case. This justifies the use of nonlinear approximation methods such as adaptive wavelet methods for the numerical treatment of SPDEs, cf. [3, 2]. The proof of (1.2) relies on characterizations of Besov spaces by wavelet expansions and on weighted Sobolev norm estimates for u , resulting from the solvability of the problem (1.1) within the spaces $\mathfrak{H}_{p,\theta}^\gamma(\mathcal{O}, T)$.

An obvious approach to improve (1.2) with respect to regularity in time is to try to combine the existing Hölder estimates in time for the elements of the spaces $\mathfrak{H}_{p,\theta}^\gamma(\mathcal{O}, T)$ (see [26, Theorem

2.9]) with the wavelet arguments in [4]. However, it turns out that a satisfactory result requires a more subtle strategy in three different aspects.

Firstly, we need an extension of the L_p -theory in [26] to an $L_q(L_p)$ -theory for SPDEs dealing with stochastic parabolic weighted Sobolev spaces $\mathfrak{H}_{p,\theta}^{\gamma,q}(\mathcal{O}, T)$ with possibly different summability parameters q and p in time and space respectively. These spaces consist of elements \tilde{u} of the form $d\tilde{u} = \tilde{f}dt + \tilde{g}^k dw_t^k$, where \tilde{u} , \tilde{f} and \tilde{g}^k , considered as stochastic processes with values in suitable weighted $L_p(\mathcal{O})$ -Sobolev spaces, are L_q -integrable w.r.t. $\mathbb{P} \otimes dt$. Such an extension is needed to obtain better Hölder estimates in time in a second step. Satisfactory existence and uniqueness results concerning solutions in the spaces $\mathfrak{H}_{p,\theta}^{\gamma,q}(\mathcal{O}, T)$ have been established in [25] for domains \mathcal{O} with C^1 -boundary. Unfortunately, the techniques used there do not work on general Lipschitz domains. Also, the $L_q(L_p)$ -results that have been obtained in [47] within the semigroup approach to SPDEs do not directly suit our purpose: On the one hand, for general Lipschitz domains \mathcal{O} the domains of the fractional powers of the leading linear differential operator cannot be characterized in terms of Sobolev or Besov spaces as in the case of a smooth domains \mathcal{O} ; see, e.g., the introduction of [4] for details. On the other hand, even in the case of a smooth domain \mathcal{O} we need regularity in terms *weighted* Sobolev spaces to obtain the optimal regularity in the scale (*).

Secondly, once we have established the solvability of SPDEs within the spaces $\mathfrak{H}_{p,\theta}^{\gamma,q}(\mathcal{O}, T)$, we have to exploit the $L_q(L_p)$ -regularity of the solution and derive improved results on the Hölder regularity in time for large q . For $\mathcal{O} = \mathbb{R}_+^d$ this has been done by Krylov [32]. It takes quite delicate arguments to apply these results to the case of bounded Lipschitz domains via a boundary flattening argument.

Thirdly, in order to obtain a reasonable Hölder-Besov regularity result, it is necessary to generalize the wavelet arguments applied in [4] to a wider range of smoothness parameters. This requires more sophisticated estimates.

In this article we tackle and solve the tasks described above. We organize the article as follows. In Section 2 we recall the definition and basic properties of the (deterministic) weighted Sobolev spaces $H_{p,\theta}^\gamma(G)$ introduced in [38] (see also [43, Chapter 6]) on general domains $G \subseteq \mathbb{R}^d$ with non-empty boundary. In Section 3 we give the definition of the spaces $\mathfrak{H}_{p,\theta}^{\gamma,q}(G, T)$ and specify the concept of a solution for equations of type (1.1) in these spaces. Moreover, we show that if we have a solution $u \in \mathfrak{H}_{p,\theta}^{\gamma,q}(G, T)$ with low regularity $\gamma \geq 0$, but f and the g^k 's have high $L_q(L_p)$ -regularity, then we can lift up the regularity of the solution (Theorem 3.8). In this sense the spaces $\mathfrak{H}_{p,\theta}^{\gamma,q}(G, T)$ are the right ones for our regularity analysis of SPDEs.

Section 4 is devoted to the solvability of Eq. (1.1) in $\mathfrak{H}_{p,\theta}^{\gamma,q}(\mathcal{O}, T)$, $\mathcal{O} \subseteq \mathbb{R}^d$ being a bounded Lipschitz domain. The focus lies on the case $q > p \geq 2$ and we restrict our considerations to equations with additive noise, i.e. $\sigma^{ik} \equiv 0$. In Subsection 4.2 we consider equations on domains with small Lipschitz constants and derive a result for general integrability parameters $q \geq p \geq 2$ (Theorem 4.2). We use an $L_q(L_p)$ -regularity result for deterministic parabolic equations from [16] and an estimate for stochastic integrals in UMD spaces from [46] to obtain a certain low $L_q(L_p)$ -regularity of the solution. Then the regularity is lifted up with the help of Theorem 3.8. In Subsection 4.2, we consider the stochastic heat equation on general bounded Lipschitz domains. Here we use the results from [47] on maximal L_q -regularity of stochastic evolution equations (see also [48] and [46]) to derive existence and uniqueness of a solution with low regularity. A main ingredient will be to prove that the domain of the square root of the weak Dirichlet Laplacian on $L_p(\mathcal{O})$ coincides with the closure of the test functions $C_0^\infty(\mathcal{O})$ in the $L_p(\mathcal{O})$ -Sobolev space of order one (Lemma 4.5). This stays true only for a certain range of $p \in [2, p_0)$ with $p_0 > 3$. Thus, so does

our result (Theorem 4.4). In a second step, we again lift up the regularity by using Theorem 3.8. In both settings we derive suitable a-priori estimates.

In Section 5 we present our result on the Hölder regularity in time of the elements of $\mathfrak{H}_{p,\theta}^{\gamma,q}(\mathcal{O}, T)$ (Theorem 5.1). It is an extension of the Hölder estimates in time for the elements of $\mathfrak{H}_{p,\theta}^{\gamma,q}(T) = \mathfrak{H}_{p,\theta}^{\gamma,q}(\mathbb{R}_+^d, T)$ in [31] to the case of bounded Lipschitz domains. The implications for the Hölder regularity of the solutions of SPDEs are described in Theorem 5.3.

In Section 6 we pave the way for the analysis of the spatial regularity of the solutions of SPDEs in the scale (*). We discuss the relationship between the weighted Sobolev spaces $H_{p,\theta}^\gamma(\mathcal{O})$ and Besov spaces. Our main result in this section, Theorem 6.9, is a general embedding of the spaces $H_{p,d-\nu p}^\gamma(\mathcal{O})$, $\gamma, \nu > 0$, into the Besov scale (*). Its proof is an extension of the wavelet arguments in the proof of [4, Theorem 3.1], where only integer valued smoothness parameters γ are considered. It can also be seen as an extension of and a supplement to the Besov regularity results for deterministic elliptic equations in [10] and [7, 8, 9, 11]. To the best of our knowledge, no such general embedding has been proven before. In the course of the discussion we also enlighten the fact that, for the relevant range of parameters γ and ν , the spaces $H_{p,d-\nu p}^\gamma(\mathcal{O})$ act like Besov spaces $B_{p,p}^{\gamma \wedge \nu}(\mathcal{O})$ with zero trace on the boundary (Remark 6.7).

In Section 7 the results of the previous sections are combined in order to determine the Hölder-Besov regularity of the elements of the stochastic parabolic spaces $\mathfrak{H}_{p,\theta}^{\gamma,q}(\mathcal{O}, T)$ and of the solutions of SPDEs within these spaces. The related result in [4] is significantly improved in several aspects; see Remark 7.3 for a detailed comparison. We obtain an estimate of the form

$$\mathbb{E} \|u\|_{C^\kappa([0,T]; B_{\tau,\tau}^\alpha(\mathcal{O}))}^q \leq C \|u\|_{\mathfrak{H}_{p,\theta}^{\gamma,q}(\mathcal{O}, T)}^q, \quad \frac{1}{\tau} = \frac{\alpha}{d} + \frac{1}{p},$$

for certain α depending on the smoothness and weight parameters γ and θ and for certain κ depending on q and α (Theorem 7.4). Using the a-priori estimates from Section 4, the right hand side of the above inequality can be estimated by suitable norms of f and g if u is the solution to the corresponding SPDE (Theorem 7.5).

Let us mention the related work [1] on the Besov regularity for the deterministic heat equation. The authors study the regularity of temperatures in terms of anisotropic Besov spaces of type $B_{\tau,\tau}^{\alpha/2,\alpha}((0, T) \times \mathcal{O})$, $1/\tau = \alpha/d + 1/p$. However, the range of admissible values for the parameter τ is a priori restricted to $(1, \infty)$, so that α is always less than $d(1 - 1/p)$. In our article the parameter τ in (*) may be any positive number, including in particular the case where τ is less than 1 and where $B_{\tau,\tau}^\alpha(\mathcal{O})$ is not a Banach space but a quasi-Banach space.

Notation and Conventions. Throughout this paper, \mathcal{O} always denotes a bounded Lipschitz domain in \mathbb{R}^d , $d \geq 1$, as specified in Definition 2.5 below. General subsets of \mathbb{R}^d are denoted by G . We write ∂G for their boundary (if it is not empty) and G° for the interior. $\mathbb{N} := \{1, 2, \dots\}$ denotes the set of strictly positive integers whereas $\mathbb{N}_0 := \mathbb{N} \cup \{0\}$. Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a complete probability space and $\{\mathcal{F}_t, t \geq 0\}$ be an increasing filtration of σ -fields $\mathcal{F}_t \subset \mathcal{F}$, each of which contains all $(\mathcal{F}, \mathbb{P})$ -null sets. By \mathcal{P} we denote the predictable σ -field generated by $\{\mathcal{F}_t, t \geq 0\}$ and we assume that $\{(w_t^1)_{t \in [0,T]}, (w_t^2)_{t \in [0,T]}, \dots\}$ are independent one-dimensional Wiener processes w.r.t. $\{\mathcal{F}_t, t \geq 0\}$. For $\kappa \in (0, 1)$ and a quasi-Banach space $(X, \|\cdot\|_X)$ we denote by $C^\kappa([0, T]; X)$ the Hölder space of continuous X -valued functions on $[0, T]$ with finite norm $\|\cdot\|_{C^\kappa([0,T]; X)}$ defined

by

$$\begin{aligned}
[u]_{C^\kappa([0,T];X)} &:= \sup_{s,t \in [0,T]} \frac{\|u(t) - u(s)\|_X}{|t - s|^\kappa}, \\
\|u\|_{C([0,T];X)} &:= \sup_{t \in [0,T]} \|u(t)\|_X, \\
\|u\|_{C^\kappa([0,T];X)} &= \|u\|_{C([0,T];X)} + [u]_{C^\kappa([0,T];X)}.
\end{aligned}$$

For $1 < p < \infty$, $L_p(A, \Sigma, \mu; X)$ denotes the space of μ -strongly measurable and p -Bochner integrable functions with values in X on a σ -finite measure space (A, Σ, μ) , endowed with the usual L_p -Norm. We write $L_p(G)$ instead of $L_p(G, \mathcal{B}(G), \lambda^d; \mathbb{R})$ if $G \in \mathcal{B}(\mathbb{R}^d)$, where $\mathcal{B}(G)$ and $\mathcal{B}(\mathbb{R}^d)$ are the Borel- σ -fields on G and \mathbb{R}^d . Recall the Hilbert space $\ell_2 := \ell_2(\mathbb{N}) = \{\mathbf{a} = (\mathbf{a}^1, \mathbf{a}^2, \dots) : |\mathbf{a}|_{\ell_2} = (\sum_k |\mathbf{a}^k|^2)^{1/2} < \infty\}$ with the inner product $\langle \mathbf{a}, \mathbf{b} \rangle_{\ell_2} = \sum_i \mathbf{a}^k \mathbf{b}^k$, for $\mathbf{a}, \mathbf{b} \in \ell_2$. The notation $C_0^\infty(G)$ is used for the space of infinitely differentiable test functions with compact support in a domain $G \subseteq \mathbb{R}^d$. For any distribution f on G and any $\varphi \in C_0^\infty(G)$, (f, φ) denotes the application of f to φ . Furthermore, for any multi-index $\alpha = (\alpha_1, \dots, \alpha_d) \in \mathbb{N}_0^d$, we write $D^\alpha f = \frac{\partial^{|\alpha|} f}{\partial x_1^{\alpha_1} \dots \partial x_d^{\alpha_d}}$ for the corresponding (generalized) derivative w.r.t. $x = (x_1, \dots, x_d) \in G$, where $|\alpha| = \alpha_1 + \dots + \alpha_d$. By making slight abuse of notation, for $m \in \mathbb{N}_0$, we write $D^m f$ for any (generalized) m -th order derivative of f and for the vector of all m -th order derivatives of f . E.g. if we write $D^m f \in X$, where X is a function space on G , we mean $D^\alpha f \in X$ for all $\alpha \in \mathbb{N}_0^d$ with $|\alpha| = m$. We also use the notation $f_{x^i x^j} = \frac{\partial^2 f}{\partial x^i \partial x^j}$, $f_{x^i} = \frac{\partial f}{\partial x^i}$. The notation f_x (respectively f_{xx}) is used synonymously for $Df := D^1 f$ (respectively for $D^2 f$), whereas $\|f_x\|_X := \sum_i \|u_{x^i}\|_X$ (respectively $\|f_{xx}\|_X := \sum_{i,j} \|f_{x^i x^j}\|_X$). Moreover, $\Delta f := \sum_i f_{x^i x^i}$, whenever it makes sense. Given $p \in [1, \infty)$ and $m \in \mathbb{N}_0$, $W_p^m(G)$ denotes the classical Sobolev space consisting of all $f \in L_p(G)$ such that $|f|_{W_p^m(G)} := \sup_{\alpha \in \mathbb{N}_0^d, |\alpha|=m} \|D^\alpha f\|_{L_p(G)}$ is finite. It is normed via $\|f\|_{W_p^m(G)} := \|f\|_{L_p(G)} + |f|_{W_p^m(G)}$. The closure of $C_0^\infty(\mathcal{O})$ in $W_p^1(\mathcal{O})$ is denoted by $\dot{W}_p^1(\mathcal{O})$ and is normed by $\|f\|_{\dot{W}_p^1(\mathcal{O})} := (\sum_i \|f_{x^i}\|_{L_p(\mathcal{O})}^p)^{1/p}$. If we have two quasi-normed spaces $(X_i, \|\cdot\|_{X_i})$, $i = 1, 2$, $X_1 \hookrightarrow X_2$ means that X_1 is continuously linearly embedded in X_2 . For a compatible couple (X_1, X_2) of quasi-Banach spaces, $[X_1, X_2]_\eta$ denotes the interpolation space of exponent $\eta \in (0, 1)$ arising from the complex interpolation method. In general, N will denote a positive finite constant, which may differ from line to line. The notation $N = N(a_1, a_2, \dots)$ is used to emphasize the dependence of the constant N on the set of parameters $\{a_1, a_2, \dots\}$. In general, this set will *not* contain all the parameters N depends on. $A \sim B$ means that A and B are equivalent.

2 Weighted Sobolev spaces

We start by recalling the definition and some basic properties of the (deterministic and stationary) weighted Sobolev spaces $H_{p,\theta}^\gamma(G)$ introduced in [38]. These spaces will serve as state spaces for the solution processes $u = (u(t))_{t \in [0,T]}$ to SPDEs of type (1.1) and they will play a fundamental role in all the forthcoming sections.

For $p \in (1, \infty)$ and $\gamma \in \mathbb{R}$, let $H_p^\gamma := H_p^\gamma(\mathbb{R}^d) := (1 - \Delta)^{-\gamma/2} L_p(\mathbb{R}^d)$ be the spaces of Bessel potentials, endowed with the norm

$$\|u\|_{H_p^\gamma} := \|(1 - \Delta)^{\gamma/2} u\|_{L_p(\mathbb{R}^d)} := \|\mathcal{F}^{-1}[(1 + |\xi|^2)^{\gamma/2} \mathcal{F}(u)(\xi)]\|_{L_p(\mathbb{R}^d)},$$

where \mathcal{F} denotes the Fourier transform. It is well known that if γ is a nonnegative integer, then

$$H_p^\gamma = \{u \in L_p : D^\alpha u \in L_p \text{ for all } \alpha \in \mathbb{N}_0^d \text{ with } |\alpha| \leq \gamma\}.$$

Let $G \subseteq \mathbb{R}^d$ be an arbitrary domain with non-empty boundary ∂G . We denote by $\rho(x) := \text{dist}(x, \partial G)$ the distance of a point $x \in G$ to the boundary ∂G . Furthermore, we fix a bounded infinitely differentiable function ψ defined on G such that for all $x \in G$,

$$\rho(x) \leq N\psi(x), \quad \rho(x)^{m-1}|D^m\psi(x)| \leq N(m) < \infty \text{ for all } m \in \mathbb{N}_0, \quad (2.1)$$

where N and $N(m)$ do not depend on $x \in G$. For a detailed construction of such a function see, e.g., [43, Chapter 3, Section 3.2.3]. Let $\zeta \in C_0^\infty(\mathbb{R}_+)$ be a non-negative function satisfying

$$\sum_{n \in \mathbb{Z}} \zeta(e^{n+t}) > c > 0 \text{ for all } t \in \mathbb{R}. \quad (2.2)$$

Note that any non-negative smooth function $\zeta \in C_0^\infty(\mathbb{R}_+)$ with $\zeta > 0$ on $[e^{-1}, e]$ satisfies (2.2). For $x \in G$ and $n \in \mathbb{Z}$, define

$$\zeta_n(x) := \zeta(e^n\psi(x)).$$

Then, there exists $k_0 > 0$ such that, for all $n \in \mathbb{Z}$, $\text{supp } \zeta_n \subset G_n := \{x \in G : e^{-n-k_0} < \rho(x) < e^{-n+k_0}\}$, i.e., $\zeta_n \in C_0^\infty(G_n)$. Moreover, $|D^m\zeta_n(x)| \leq N(\zeta, m)e^{mn}$ for all $x \in G$ and $m \in \mathbb{N}_0$, and $\sum_{n \in \mathbb{Z}} \zeta_n(x) \geq \delta > 0$ for all $x \in G$. For $p \in (1, \infty)$ and $\gamma, \theta \in \mathbb{R}$, we denote by $H_{p,\theta}^\gamma(G)$ the space of all distributions u on G such that

$$\|u\|_{H_{p,\theta}^\gamma(G)}^p := \sum_{n \in \mathbb{Z}} e^{n\theta} \|\zeta_{-n}(e^n \cdot) u(e^n \cdot)\|_{H_p^\gamma}^p < \infty.$$

It is well-known that

$$L_{p,\theta}(G) := H_{p,\theta}^0(G) = L_p(G, \rho^{\theta-d} dx),$$

and that, if γ is a positive integer,

$$H_{p,\theta}^\gamma(G) = \{u \in L_{p,\theta}(G) : \rho^{|\alpha|} D^\alpha u \in L_{p,\theta}(G) \text{ for all } \alpha \in \mathbb{N}_0^d \text{ with } |\alpha| \leq \gamma\},$$

$$\|u\|_{H_{p,\theta}^\gamma(G)}^p \sim \sum_{|\alpha| \leq \gamma} \int_G |\rho^{|\alpha|} D^\alpha u|^p \rho^{\theta-d} dx;$$

see, e.g., [38, Proposition 2.2]. This is the reason why the space $H_{p,\theta}^\gamma(G)$ is called weighted Sobolev space of order γ , with summability parameter p and weight parameter θ .

For $p \in (1, \infty)$ and $\gamma \in \mathbb{R}$ we write $H_p^\gamma(\ell_2)$ for the collection of all sequences $g = (g^1, g^2, \dots)$ of distributions on \mathbb{R}^d with $g^k \in H_p^\gamma$ for each $k \in \mathbb{N}$ and

$$\|g\|_{H_p^\gamma(\ell_2)} := \|g\|_{H_p^\gamma(\mathbb{R}^d; \ell_2)} := \|(1 - \Delta)^{\gamma/2} g|_{\ell_2}\|_{L_p} := \left\| \left(\sum_{k=1}^{\infty} |(1 - \Delta)^{\gamma/2} g^k|^2 \right)^{1/2} \right\|_{L_p} < \infty.$$

Analogously, for $\theta \in \mathbb{R}$, a sequence $g = (g^1, g^2, \dots)$ of distributions on G is in $H_{p,\theta}^\gamma(G; \ell_2)$ if, and only if, $g^k \in H_{p,\theta}^\gamma(G)$ for each $k \in \mathbb{N}$ and

$$\|g\|_{H_{p,\theta}^\gamma(G; \ell_2)}^p := \sum_{n \in \mathbb{Z}} e^{n\theta} \|\zeta_{-n}(e^n \cdot) g(e^n \cdot)\|_{H_p^\gamma}^p < \infty.$$

Now we present some useful properties of the space $H_{p,\theta}^\gamma(G)$ taken from [38], see also [29, 30].

Lemma 2.1. (i) The space $C_0^\infty(G)$ is dense in $H_{p,\theta}^\gamma(G)$.

(ii) Assume that $\gamma - d/p = m + \nu$ for some $m \in \mathbb{N}_0$, $\nu \in (0, 1]$ and that $i, j \in \mathbb{N}_0^d$ are multi-indices such that $|i| \leq m$ and $|j| = m$. Then for any $u \in H_{p,\theta}^\gamma(G)$, we have

$$\begin{aligned} \psi^{|i|+\theta/p} D^i u &\in C(G), \quad \psi^{m+\nu+\theta/p} D^j u \in C^\nu(G), \\ |\psi^{|i|+\theta/p} D^i u|_{C(G)} + [\psi^{m+\nu+\theta/p} D^j u]_{C^\nu(G)} &\leq N \|u\|_{H_{p,\theta}^\gamma(G)}. \end{aligned}$$

(iii) $u \in H_{p,\theta}^\gamma(G)$ if, and only if, $u, \psi u_x \in H_{p,\theta}^{\gamma-1}(G)$ and

$$\|u\|_{H_{p,\theta}^\gamma(G)} \leq N \|\psi u_x\|_{H_{p,\theta}^{\gamma-1}(G)} + N \|u\|_{H_{p,\theta}^{\gamma-1}(G)} \leq N \|u\|_{H_{p,\theta}^\gamma(G)}.$$

Also, $u \in H_{p,\theta}^\gamma(G)$ if, and only if, $u, (\psi u)_x \in H_{p,\theta}^{\gamma-1}(G)$ and

$$\|u\|_{H_{p,\theta}^\gamma(G)} \leq N \|(\psi u)_x\|_{H_{p,\theta}^{\gamma-1}(G)} + N \|u\|_{H_{p,\theta}^{\gamma-1}(G)} \leq N \|u\|_{H_{p,\theta}^\gamma(G)}.$$

(iv) For any $\nu, \gamma \in \mathbb{R}$, $\psi^\nu H_{p,\theta}^\gamma(G) = H_{p,\theta-p\nu}^\gamma(G)$ and

$$\|u\|_{H_{p,\theta-p\nu}^\gamma(G)} \leq N \|\psi^{-\nu} u\|_{H_{p,\theta}^\gamma(G)} \leq N \|u\|_{H_{p,\theta-p\nu}^\gamma(G)}.$$

(v) If $\gamma \in (\gamma_0, \gamma_1)$ then, for any $\varepsilon > 0$, there exists a constant $N = N(\gamma_0, \gamma_1, \theta, p, \varepsilon)$, such that

$$\|u\|_{H_{p,\theta}^\gamma(G)} \leq \varepsilon \|u\|_{H_{p,\theta}^{\gamma_1}(G)} + N(\gamma_0, \gamma_1, \theta, p, \varepsilon) \|u\|_{H_{p,\theta}^{\gamma_0}(G)}.$$

Also, if $\theta \in (\theta_0, \theta_1)$ then, for any $\varepsilon > 0$, there exists a constant $N = N(\theta_0, \theta_1, \gamma, p, \varepsilon)$, such that

$$\|u\|_{H_{p,\theta}^\gamma(G)} \leq \varepsilon \|u\|_{H_{p,\theta_0}^\gamma(G)} + N(\theta_0, \theta_1, \gamma, p, \varepsilon) \|u\|_{H_{p,\theta_1}^\gamma(G)}.$$

(vi) There exists a constant $c_0 > 0$ depending on p, θ, γ and the function ψ such that, for all $c \geq c_0$, the operator $\psi^2 \Delta - c$ is a homeomorphism from $H_{p,\theta}^{\gamma+1}(G)$ to $H_{p,\theta}^{\gamma-1}(G)$.

Remark 2.2. Assertions (vi) and (iv) in Lemma 2.1 imply the following: If $u \in H_{p,\theta-p}^\gamma(G)$ and $\Delta u \in H_{p,\theta+p}^\gamma(G)$, then $u \in H_{p,\theta-p}^\gamma(G)$ and there exists a constant N , which does not depend on u , such that

$$\|u\|_{H_{p,\theta-p}^{\gamma+2}(G)} \leq N \|\Delta u\|_{H_{p,\theta+p}^\gamma(G)} + N \|u\|_{H_{p,\theta-p}^\gamma(G)}.$$

A proof of the following equivalent characterization of the weighted Sobolev spaces $H_{p,\theta}^\gamma(G)$ can be found in [38, Proposition 2.2].

Lemma 2.3. Let $\{\xi_n : n \in \mathbb{Z}\} \subseteq C_0^\infty(G)$ be such that for all $n \in \mathbb{Z}$ and $m \in \mathbb{N}_0$,

$$|D^m \xi_n| \leq N(m) c^{nm} \quad \text{and} \quad \text{supp } \xi_n \subseteq \{x \in G : c^{-n-k_0} < \rho(x) < c^{-n+k_0}\} \quad (2.3)$$

for some $c > 1$ and $k_0 > 0$, where the constant $N(m)$ does not depend on $n \in \mathbb{Z}$ and $x \in G$. Then, for any $u \in H_{p,\theta}^\gamma(G)$,

$$\sum_{n \in \mathbb{Z}} c^{n\theta} \|\xi_{-n}(c^n \cdot) u(c^n \cdot)\|_{H_{p,\theta}^\gamma}^p \leq N \|u\|_{H_{p,\theta}^\gamma}^p.$$

If in addition

$$\sum_{n \in \mathbb{Z}} \xi_n(x) \geq \delta > 0 \quad \text{for all } x \in G \quad (2.4)$$

then the converse inequality also holds.

Remark 2.4. (i) *It is easy to check that both*

$$\{\xi_n^{(1)} := e^{-n}(\zeta_n)_{x^i} : n \in \mathbb{Z}\} \quad \text{and} \quad \{\xi_n^{(2)} := e^{-2n}(\zeta_n)_{x^i x^j} : n \in \mathbb{Z}\}$$

satisfy (2.3) with $c := e$. Therefore,

$$\sum_{n \in \mathbb{Z}} e^{n\theta} \left(\|e^n(\zeta_{-n})_{x^i}(e^n \cdot)u(e^n \cdot)\|_{H_p^\gamma}^p + \|e^{2n}(\zeta_{-n})_{x^i x^j}(e^n \cdot)u(e^n \cdot)\|_{H_p^\gamma}^p \right) \leq N \|u\|_{H_{p,\theta}^\gamma(G)}^p.$$

(ii) *Given $k_1 \geq 1$, fix a function $\tilde{\zeta} \in C_0^\infty(\mathbb{R}_+)$ with*

$$\tilde{\zeta}(t) = 1 \quad \text{for all } t \in \left[\frac{1}{N} 2^{-k_1}, N(0) 2^{k_1} \right],$$

where N and $N(0)$ are as in (2.1). Then, the sequence $\{\xi_n : n \in \mathbb{Z}\} \subseteq C_0^\infty(G)$ defined by

$$\xi_n := \tilde{\zeta}(2^n \psi(\cdot)), \quad n \in \mathbb{Z},$$

fulfils the conditions (2.3) and (2.4) from Lemma 2.3 with $c = 2$ and a suitable $k_0 > 0$. Furthermore,

$$\xi_n(x) = 1 \quad \text{for all } x \in \rho^{-1}(2^{-n}[2^{-k_1}, 2^{k_1}]).$$

In this paper, \mathcal{O} **will always denote a bounded Lipschitz domain in \mathbb{R}^d** . More precisely:

Definition 2.5. *We call a bounded domain $\mathcal{O} \subset \mathbb{R}^d$ a Lipschitz domain if, and only if, for any $x_0 = (x_0^1, x_0') \in \partial\mathcal{O}$, there exists a Lipschitz continuous function $\mu_0 : \mathbb{R}^{d-1} \rightarrow \mathbb{R}$ such that, upon relabeling and reorienting the coordinate axes if necessary, we have*

(i) $\mathcal{O} \cap B_{r_0}(x_0) = \{x = (x^1, x') \in B_{r_0}(x_0) : x^1 > \mu_0(x')\}$, and

(ii) $|\mu_0(x') - \mu_0(y')| \leq K_0|x' - y'|$, for any $x', y' \in \mathbb{R}^{d-1}$,

where r_0, K_0 are independent of x_0 .

Remark 2.6. *Recall that for a bounded Lipschitz domain $\mathcal{O} \subseteq \mathbb{R}^d$,*

$$\dot{W}_p^1(\mathcal{O}) = H_{p,d-p}^1(\mathcal{O})$$

with equivalent norms. This follows from [35, Theorem 9.7] and Poincaré's inequality.

3 Stochastic parabolic weighted Sobolev spaces and SPDEs

In this section, we first introduce the stochastic parabolic spaces $\mathfrak{H}_{p,\theta}^{\gamma,q}(G, T)$ for arbitrary domains $G \subseteq \mathbb{R}^d$ with non-empty boundary in analogy to the spaces $\mathfrak{H}_{p,\theta}^{\gamma,q}(T) = \mathfrak{H}_{p,\theta}^{\gamma,q}(\mathbb{R}_+^d, T)$ from [31, 32]. Then we show that they are suitable to serve as solution spaces for equations of type (1.1) in the following sense: If we have a solution $u \in \mathfrak{H}_{p,\theta}^{\gamma,q}(G, T)$ with low regularity $\gamma \geq 0$, but f and the g^k 's have high $L_q(L_p)$ -regularity, then we can lift up the regularity of the solution (Theorem 3.8).

Definition 3.1. Let G be a domain in \mathbb{R}^d with non-empty boundary. For $p, q \in (1, \infty)$, $\gamma, \theta \in \mathbb{R}$ and $T \in (0, \infty)$ we define

$$\begin{aligned}\mathbb{H}_{p,\theta}^{\gamma,q}(G, T) &:= L_q(\Omega \times [0, T], \mathcal{P}, \mathbb{P} \otimes dt; H_{p,\theta}^\gamma(G)), \\ \mathbb{H}_{p,\theta}^{\gamma,q}(G, T; \ell_2) &:= L_q(\Omega \times [0, T], \mathcal{P}, \mathbb{P} \otimes dt; H_{p,\theta}^\gamma(G; \ell_2)), \\ U_{p,\theta}^{\gamma,q}(G) &:= L_q(\Omega, \mathcal{F}_0, \mathbb{P}; \psi^{1-2/q} H_{p,\theta}^{\gamma-2/q}(G)).\end{aligned}$$

If $p = q$ we also write $\mathbb{H}_{p,\theta}^\gamma(G, T)$, $\mathbb{H}_{p,\theta}^\gamma(G, T; \ell_2)$ and $U_{p,\theta}^\gamma(G)$ instead of $\mathbb{H}_{p,\theta}^{\gamma,p}(G, T)$, $\mathbb{H}_{p,\theta}^{\gamma,p}(G, T; \ell_2)$ and $U_{p,\theta}^{\gamma,p}(G)$ respectively.

From now on let

$$p \in [2, \infty), \quad q \in [2, \infty), \quad \gamma \in \mathbb{R}, \quad \theta \in \mathbb{R}.$$

Definition 3.2. Let G be a domain in \mathbb{R}^d with non-empty boundary. We write $u \in \mathfrak{H}_{p,\theta}^{\gamma,q}(G, T)$ if, and only if, $u \in \mathbb{H}_{p,\theta-p}^{\gamma,q}(G, T)$, $u(0, \cdot) \in U_{p,\theta}^{\gamma,q}(G)$, and there exist some $f \in \mathbb{H}_{p,\theta+p}^{\gamma-2,q}(G, T)$ and $g \in \mathbb{H}_{p,\theta}^{\gamma-1,q}(G, T; \ell_2)$ such that

$$du = f dt + g^k dw_t^k$$

in the sense of distributions. That is, for any $\varphi \in C_0^\infty(G)$, with probability one, the equality

$$(u(t, \cdot), \varphi) = (u(0, \cdot), \varphi) + \int_0^t (f(s, \cdot), \varphi) ds + \sum_{k=1}^{\infty} \int_0^t (g^k(s, \cdot), \varphi) dw_s^k$$

holds for all $t \in [0, T]$, where the series is assumed to converge uniformly on $[0, T]$ in probability. In this situation we write $\mathbb{D}u := f$ and $\mathbb{S}u := g$. The norm in $\mathfrak{H}_{p,\theta}^{\gamma,q}(G, T)$ is defined as

$$\|u\|_{\mathfrak{H}_{p,\theta}^{\gamma,q}(G, T)} := \|u\|_{\mathbb{H}_{p,\theta-p}^{\gamma,q}(G, T)} + \|\mathbb{D}u\|_{\mathbb{H}_{p,\theta+p}^{\gamma-2,q}(G, T)} + \|\mathbb{S}u\|_{\mathbb{H}_{p,\theta}^{\gamma-1,q}(G, T; \ell_2)} + \|u(0, \cdot)\|_{U_{p,\theta}^{\gamma,q}(G)}.$$

If $p = q$ we also write $\mathfrak{H}_{p,\theta}^\gamma(G, T)$ instead of $\mathfrak{H}_{p,\theta}^{\gamma,p}(G, T)$.

Remark 3.3. Replacing G by \mathbb{R}^d and omitting the weight parameter θ and the weight function ψ in the definitions above, one obtains the spaces $\mathbb{H}_p^{\gamma,q}(T) = \mathbb{H}_p^{\gamma,q}(\mathbb{R}^d, T)$, $\mathbb{H}_p^{\gamma,q}(T; \ell_2) = \mathbb{H}_p^{\gamma,q}(\mathbb{R}^d, T; \ell_2)$, $U_p^{\gamma,q} = U_p^{\gamma,q}(\mathbb{R}^d)$, and $\mathcal{H}_p^{\gamma,q}(T)$ as introduced in [32, Definition 3.5]. The latter are denoted by $\mathcal{H}_p^{\gamma,q}(T)$ in [31]; if $q = p$ they coincide with the spaces $\mathcal{H}_p^\gamma(T)$ introduced in [29, Definition 3.1].

We consider initial value problems of the form

$$du = (a^{ij} u_{x_i x_j} + f) dt + (\sigma^{ik} u_{x_i} + g^k) dw_t^k, \quad u(0, \cdot) = u_0, \quad (3.1)$$

on an arbitrary domain $G \subseteq \mathbb{R}^d$ with non-empty boundary. We use the following solution concept.

Definition 3.4. We say that a stochastic process $u \in \mathfrak{H}_{p,\theta}^{\gamma,q}(G, T)$ is a solution of Eq. (3.1) if, and only if,

$$u(0, \cdot) = u_0, \quad \mathbb{D}u = a^{ij} u_{x_i x_j} + f, \quad \text{and} \quad \mathbb{S}u = (\sigma^{ik} u_{x_i} + g^k)_{k \in \mathbb{N}},$$

in the sense of Definition 3.2.

Remark 3.5. Here and in the sequel we use the summation convention on the repeated indices i, j, k . The question, in which sense, for a bounded Lipschitz domain $\mathcal{O} \subseteq \mathbb{R}^d$, the elements of $\mathfrak{H}_{p,\theta}^{\gamma,q}(\mathcal{O}, T)$ fulfil a zero Dirichlet boundary condition as in Eq. (1.1), will be answered in Remark 6.7.

We make the following assumptions on the coefficients in Eq. (3.1). Throughout this paper, whenever we will talk about this equation, we will assume that they are fulfilled.

Assumption 3.6. (i) The coefficients $a^{ij} = a^{ij}(\omega, t)$ and $\sigma^{ik} = \sigma^{ik}(\omega, t)$ are predictable. They do not depend on $x \in G$. Furthermore, $a^{ij} = a^{ji}$ for $i, j \in \{1, \dots, d\}$.

(ii) There exist constants $\delta_0, K > 0$ such that for any $(\omega, t) \in \Omega \times [0, T]$ and $\lambda \in \mathbb{R}^d$,

$$\delta_0 |\lambda|^2 \leq \bar{a}^{ij}(\omega, t) \lambda^i \lambda^j \leq K |\lambda|^2,$$

where $\bar{a}^{ij}(\omega, t) := a^{ij}(\omega, t) - \frac{1}{2}(\sigma^{i\cdot}(\omega, t), \sigma^{j\cdot}(\omega, t))_{\ell_2}$, with $\sigma^{i\cdot}(\omega, t) = (\sigma^{ik}(\omega, t))_{k \in \mathbb{N}} \in \ell_2$.

We will use the following result taken from [31, Lemma 2.3].

Lemma 3.7. Let $p \geq 2$, $m \in \mathbb{N}$, and, for $i = 1, 2, \dots, m$,

$$\lambda_i \in (0, \infty), \quad \gamma_i \in \mathbb{R}, \quad u^{(i)} \in \mathcal{H}_p^{\gamma_i+2}(T), \quad u^{(i)}(0, \cdot) = 0.$$

Denote $\Lambda_i := (\lambda_i - \Delta)^{\gamma_i/2}$. Then

$$\begin{aligned} \mathbb{E} \left[\int_0^T \prod_{i=1}^m \|\Lambda_i \Delta u^{(i)}\|_{L_p}^p dt \right] &\leq N \sum_{i=1}^m \mathbb{E} \left[\int_0^T \left(\|\Lambda_i f^{(i)}\|_{L_p}^p + \|\Lambda_i g_x^{(i)}\|_{L_p(\ell_2)}^p \right) \prod_{\substack{j=1 \\ j \neq i}}^m \|\Lambda_j \Delta u^{(j)}\|_{L_p}^p dt \right] \\ &\quad + N \sum_{1 \leq i < j \leq m} \mathbb{E} \left[\int_0^T \|\Lambda_i g_x^{(i)}\|_{L_p(\ell_2)}^p \|\Lambda_j g_x^{(j)}\|_{L_p(\ell_2)}^p \prod_{\substack{k=1 \\ k \neq i, j}}^m \|\Lambda_k \Delta u^{(k)}\|_{L_p}^p dt \right], \end{aligned}$$

where $f^{(i)} := \mathbb{D}u^{(i)} - a^{rs} u_{x^r x^s}^{(i)}$, $g^{(i)k} := \mathbb{S}^k u^{(i)} - \sigma^{rk} u_{x^r}^{(i)}$ and $L_p(\ell_2) := H_p^0(\ell_2)$. The constant N depends only on m, d, p, δ_0 , and K .

Now we are able to prove that if we have a solution $u \in \mathfrak{H}_{p,\theta}^{\gamma+1,q}(G, T)$ to Eq. (3.1) and if the regularity of the forcing terms f and g is high then we can lift the regularity of the solution. Note that in the next theorem there is no restriction, neither on the shape of the domain $G \subseteq \mathbb{R}^d$ nor on the parameters $\theta, \gamma \in \mathbb{R}$.

Theorem 3.8. Let $G \subseteq \mathbb{R}^d$ be an arbitrary domain with non-empty boundary. Let $\gamma \in \mathbb{R}$, $p \geq 2$ and $q = pm$ for some $m \in \mathbb{N}$. Let $f \in \mathbb{H}_{p,\theta+p}^{\gamma,q}(G, T)$, $g \in \mathbb{H}_{p,\theta}^{\gamma+1,q}(G, T; \ell_2)$ and let $u \in \mathfrak{H}_{p,\theta}^{\gamma+1,q}(G, T)$ be a solution to Eq. (3.1) with $u_0 = 0$. Then $u \in \mathfrak{H}_{p,\theta}^{\gamma+2,q}(G, T)$, and

$$\|u\|_{\mathbb{H}_{p,\theta-p}^{\gamma+2,q}(G,T)}^q \leq N \left(\|u\|_{\mathbb{H}_{p,\theta-p}^{\gamma+1,q}(G,T)}^q + \|f\|_{\mathbb{H}_{p,\theta+p}^{\gamma,q}(G,T)}^q + \|g\|_{\mathbb{H}_{p,\theta}^{\gamma+1,q}(G,T;\ell_2)}^q \right),$$

where the constant $N \in (0, \infty)$ does not depend on u, f and g .

Proof. The case $m = 1$, i.e., $p = q$ is covered by [26, Lemma 3.2]. Therefore, let $m \geq 2$. According to Remark 2.2 it is enough to show that

$$\|\Delta u\|_{\mathbb{H}_{p,\theta+p}^{\gamma,q}(G,T)}^q \leq N \left(\|u\|_{\mathbb{H}_{p,\theta-p}^{\gamma+1,q}(G,T)}^q + \|f\|_{\mathbb{H}_{p,\theta+p}^{\gamma,q}(G,T)}^q + \|g\|_{\mathbb{H}_{p,\theta}^{\gamma+1,q}(G,T;\ell_2)}^q \right).$$

Using the definition of weighted Sobolev spaces from Section 2, we observe that

$$\begin{aligned} \|\Delta u\|_{\mathbb{H}_{p,\theta+p}^{\gamma,q}(G,T)}^q &= \mathbb{E} \left[\int_0^T \left(\sum_{n \in \mathbb{Z}} e^{n(\theta+p)} \|(\zeta_{-n} \Delta u(t))(e^{n \cdot})\|_{H_p^\gamma}^p \right)^m dt \right] \\ &\leq N \mathbb{E} \left[\int_0^T \left(\sum_{n \in \mathbb{Z}} e^{n(\theta+p)} \left(\|\Delta(\zeta_{-n} u(t))(e^{n \cdot})\|_{H_p^\gamma}^p \right. \right. \right. \\ &\quad \left. \left. \left. + \|(\Delta \zeta_{-n} u(t))(e^{n \cdot})\|_{H_p^\gamma}^p + \|(\zeta_{-n} x u_x(t))(e^{n \cdot})\|_{H_p^\gamma}^p \right) \right)^m dt \right]. \end{aligned}$$

(Here $\zeta_{-n} u_x$ is meant to be a scalar product in \mathbb{R}^d .) Now we can use Jensen's inequality and Remark 2.4(i) to obtain

$$\begin{aligned} \|\Delta u\|_{\mathbb{H}_{p,\theta+p}^{\gamma,q}(G,T)}^q &\leq N \mathbb{E} \left[\int_0^T \left(\sum_{n \in \mathbb{Z}} e^{n(\theta+p)} \|\Delta(\zeta_{-n} u(t))(e^{n \cdot})\|_{H_p^\gamma}^p \right)^m \right. \\ &\quad \left. + \|u(t)\|_{H_{p,\theta-p}^\gamma(G)}^q + \|u_x(t)\|_{H_{p,\theta}^\gamma(G)}^q dt \right]. \end{aligned}$$

An application of Lemma 2.1(iii) and (iv) leads to

$$\|\Delta u\|_{\mathbb{H}_{p,\theta+p}^{\gamma,q}(G,T)}^q \leq N \mathbb{E} \left[\int_0^T \left(\sum_{n \in \mathbb{Z}} e^{n(\theta+p)} \|\Delta(\zeta_{-n} u(t))(e^{n \cdot})\|_{H_p^\gamma}^p \right)^m dt \right] + N \|u\|_{\mathbb{H}_{p,\theta-p}^{\gamma+1,q}(G,T)}^q.$$

Therefore, it is enough to estimate the first term on the right hand side,

$$\begin{aligned} &\mathbb{E} \left[\int_0^T \left(\sum_{n \in \mathbb{Z}} e^{n(\theta+p)} \|\Delta(\zeta_{-n} u(t))(e^{n \cdot})\|_{H_p^\gamma}^p \right)^m dt \right] \\ &= \mathbb{E} \left[\int_0^T \sum_{n_1, \dots, n_m \in \mathbb{Z}} e^{(\sum_{i=1}^m n_i)(\theta+p)} \prod_{i=1}^m \|\Delta(\zeta_{-n_i} u(t))(e^{n_i \cdot})\|_{H_p^\gamma}^p dt \right]. \end{aligned}$$

Tonelli's theorem together with the relation

$$\|u(c \cdot)\|_{H_p^\gamma}^p = c^{p\gamma-d} \|(c^{-2} - \Delta)^{\gamma/2} u\|_{L_p}^p \quad \text{for } c \in (0, \infty), \quad (3.2)$$

applied to $\Delta u^{(n_i)}$ with $u^{(n)} := \zeta_{-n} u$ for $n \in \mathbb{Z}$, show that we only have to handle

$$\sum_{n_1, \dots, n_m \in \mathbb{Z}} e^{(\sum_{i=1}^m n_i)(\theta+p+p\gamma-d)} \mathbb{E} \left[\int_0^T \prod_{i=1}^m \|(e^{-2n_i} - \Delta)^{\gamma/2} \Delta u^{(n_i)}(t)\|_{L_p}^p dt \right].$$

Note that since $u \in \mathfrak{H}_{p,\theta}^{\gamma+1,q}(G, T)$ solves Eq. (3.1) with vanishing initial value, $u^{(n)}$ is a solution of the equation

$$dv = (a^{rs}v_{x^r x^s} + f^{(n)})dt + (\sigma^{rk}v_{x^r} + g^{(n)k})dw_t^k, \quad v(0, \cdot) = 0,$$

on \mathbb{R}^d , where $f^{(n)} = -2a^{rs}(\zeta_{-n})_{x^s}u_{x^r} - a^{rs}(\zeta_{-n})_{x^r x^s}u + \zeta_{-n}f$ and $g^{(n)k} = -\sigma^{rk}(\zeta_{-n})_{x^r}u + \zeta_{-n}g^k$. Furthermore, applying [29, Theorem 4.10], we have $u^{(n)} \in \mathcal{H}_p^{\gamma+2}(T)$. Thus, we can use Lemma 3.7 to obtain

$$\mathbb{E} \left[\int_0^T \prod_{i=1}^m \|(e^{-2n_i} - \Delta)^{\gamma/2} \Delta u^{(n_i)}(t)\|_{L_p}^p dt \right] \leq N \sum_{i=1}^m (I_{n_i} + II_{n_i}) + N \sum_{1 \leq i < j \leq m} III_{n_i n_j}$$

where we denote

$$\begin{aligned} I_{n_i} &:= \mathbb{E} \left[\int_0^T \|\Lambda_{n_i} f^{(n_i)}(t)\|_{L_p}^p \prod_{\substack{j=1 \\ j \neq i}}^m \|\Lambda_{n_j} \Delta u^{(n_j)}(t)\|_{L_p}^p dt \right], \\ II_{n_i} &:= \mathbb{E} \left[\int_0^T \|\Lambda_{n_i} g_x^{(n_i)}(t)\|_{L_p(\ell_2)}^p \prod_{\substack{j=1 \\ j \neq i}}^m \|\Lambda_{n_j} \Delta u^{(n_j)}(t)\|_{L_p}^p dt \right], \\ III_{n_i n_j} &:= \mathbb{E} \left[\int_0^T \|\Lambda_{n_i} g_x^{(n_i)}(t)\|_{L_p(\ell_2)}^p \|\Lambda_{n_j} g_x^{(n_j)}(t)\|_{L_p(\ell_2)}^p \prod_{\substack{k=1 \\ k \neq i, j}}^m \|\Lambda_{n_k} \Delta u^{(n_k)}(t)\|_{L_p}^p dt \right], \end{aligned}$$

with $\Lambda_n := (e^{-2n} - \Delta)^{\gamma/2}$. Thus, it is enough to find a proper estimate for

$$\sum_{n_1, \dots, n_m \in \mathbb{Z}} e^{(\sum_{i=1}^m n_i)(\theta + p + p\gamma - d)} \left(\sum_{i=1}^m (I_{n_i} + II_{n_i}) + \sum_{1 \leq i < j \leq m} III_{n_i n_j} \right).$$

Applying (3.2) first, followed by Tonelli's theorem, then Hölder's and Young's inequality, leads to

$$\begin{aligned} & \sum_{n_1, \dots, n_m \in \mathbb{Z}} e^{(\sum_{i=1}^m n_i)(\theta + p + p\gamma - d)} \sum_{i=1}^m I_{n_i} \\ &= \sum_{n_1, \dots, n_m \in \mathbb{Z}} e^{(\sum_{i=1}^m n_i)(\theta + p)} \sum_{i=1}^m \mathbb{E} \left[\int_0^T \|f^{(n_i)}(t, e^{n_i \cdot})\|_{H_p^\gamma}^p \prod_{\substack{j=1 \\ j \neq i}}^m \|\Delta u^{(n_j)}(t, e^{n_j \cdot})\|_{H_p^\gamma}^p dt \right] \\ &\leq N \mathbb{E} \left[\int_0^T \left(\sum_{n \in \mathbb{Z}} e^{n(\theta + p)} \|f^{(n)}(t, e^{n \cdot})\|_{H_p^\gamma}^p \right) \left(\sum_{n \in \mathbb{Z}} e^{n(\theta + p)} \|\Delta u^{(n)}(t, e^{n \cdot})\|_{H_p^\gamma}^p \right)^{m-1} dt \right] \\ &\leq N(\varepsilon) \mathbb{E} \left[\int_0^T \left(\sum_{n \in \mathbb{Z}} e^{n(\theta + p)} \|f^{(n)}(t, e^{n \cdot})\|_{H_p^\gamma}^p \right)^{\frac{q}{p}} dt \right] + \varepsilon \mathbb{E} \left[\int_0^T \left(\sum_{n \in \mathbb{Z}} e^{n(\theta + p)} \|\Delta u^{(n)}(t, e^{n \cdot})\|_{H_p^\gamma}^p \right)^{\frac{q}{p}} dt \right]. \end{aligned}$$

Using the definition of $f^{(n)}$ and arguing as at the beginning of the proof, we get

$$\begin{aligned} \sum_{n \in \mathbb{Z}} e^{n(\theta + p)} \|f^{(n)}(t, e^{n \cdot})\|_{H_p^\gamma}^p &\leq N \left(\|u_x(t)\|_{H_{p,\theta}^{\gamma}(G)}^p + \|u(t)\|_{H_{p,\theta-p}^{\gamma}(G)}^p + \|f(t)\|_{H_{p,\theta+p}^{\gamma}(G)}^p \right) \\ &\leq N \left(\|u(t)\|_{H_{p,\theta-p}^{\gamma+1}(G)}^p + \|f(t)\|_{H_{p,\theta+p}^{\gamma}(G)}^p \right). \end{aligned}$$

Moreover,

$$\begin{aligned}
\sum_{n \in \mathbb{Z}} e^{n(\theta+p)} \|\Delta u^{(n)}(t, e^{n \cdot})\|_{H_p^\gamma}^p &\leq \sum_{n \in \mathbb{Z}} e^{n(\theta+p)} \|(\Delta \zeta_{-n} u(t))(e^{n \cdot})\|_{H_p^\gamma}^p \\
&\quad + \sum_{n \in \mathbb{Z}} e^{n(\theta+p)} \|(\zeta_{-n} u_x(t))(e^{n \cdot})\|_{H_p^\gamma}^p + \sum_{n \in \mathbb{Z}} e^{n(\theta+p)} \|(\zeta_{-n} \Delta u(t))(e^{n \cdot})\|_{H_p^\gamma}^p \\
&\leq N \left(\|u(t)\|_{H_{p, \theta-p}^\gamma(G)}^p + \|u_x(t)\|_{H_{p, \theta}^\gamma(G)}^p + \|\Delta u\|_{H_{p, \theta+p}^\gamma(G)}^p \right) \\
&\leq N \left(\|u(t)\|_{H_{p, \theta-p}^{\gamma+1}(G)}^p + \|\Delta u\|_{H_{p, \theta+p}^\gamma(G)}^p \right).
\end{aligned}$$

Combining the last three estimates, we obtain for any $\varepsilon > 0$ a constant $N(\varepsilon) \in (0, \infty)$, such that

$$\sum_{n_1, \dots, n_m \in \mathbb{Z}} e^{(\sum_{i=1}^m n_i)(\theta+p+p\gamma-d)} \sum_{i=1}^m I_{n_i} \leq \varepsilon \|\Delta u\|_{\mathbb{H}_{p, \theta+p}^{\gamma, q}(G, T)}^q + N(\varepsilon) \left(\|f\|_{\mathbb{H}_{p, \theta+p}^{\gamma, q}(G, T)}^q + \|u\|_{\mathbb{H}_{p, \theta-p}^{\gamma+1, q}(G, T)}^q \right).$$

Using similar arguments we obtain

$$\begin{aligned}
\sum_{n_1, \dots, n_m \in \mathbb{Z}} e^{(\sum_{i=1}^m n_i)(\theta+p+p\gamma-d)} &\left(\sum_{i=1}^m \mathbb{I}_{n_i} + \sum_{1 \leq i < j \leq m} \mathbb{III}_{n_i n_j} \right) \\
&\leq \varepsilon \|\Delta u\|_{\mathbb{H}_{p, \theta+p}^{\gamma, q}(G, T)}^q + N(\varepsilon) \left(\|g\|_{\mathbb{H}_{p, \theta}^{\gamma+1, q}(G, T; \ell_2)}^q + \|u\|_{\mathbb{H}_{p, \theta-p}^{\gamma+1, q}(G, T)}^q \right),
\end{aligned}$$

which finishes the proof. \square

Iterating this result has the following consequence.

Corollary 3.9. *Let $\gamma \geq 1$, $p \in [2, \infty)$ and $q = mp$ for some $m \in \mathbb{N}$. Furthermore, assume that $f \in \mathbb{H}_{p, \theta+p}^{\gamma-2, q}(G, T)$, $g \in \mathbb{H}_{p, \theta}^{\gamma-1, q}(G, T; \ell_2)$, and that $u \in \mathbb{H}_{p, \theta-p}^{0, q}(G, T)$ satisfies Eq. (3.1) with $u_0 = 0$. Then $u \in \mathfrak{H}_{p, \theta}^{\gamma, q}(G, T)$, and*

$$\|u\|_{\mathfrak{H}_{p, \theta}^{\gamma, q}(G, T)}^q \leq N \left(\|u\|_{\mathbb{H}_{p, \theta-p}^{0, q}(G, T)}^q + \|f\|_{\mathbb{H}_{p, \theta+p}^{\gamma-2, q}(G, T)}^q + \|g\|_{\mathbb{H}_{p, \theta}^{\gamma-1, q}(G, T; \ell_2)}^q \right),$$

where the constant $N \in (0, \infty)$ does not depend on u , f and g .

Remark 3.10. *An extension of the results above to the case where the coefficients depend on the space variable $x \in G$ can be proved along the lines of [24, 26]. Also, the symmetry of a^{ij} can be dropped. To keep the expositions at a reasonable level, we do not discuss these cases.*

4 Solvability of SPDEs within $\mathfrak{H}_{p, \theta}^{\gamma, q}(\mathcal{O}, T)$

In this section we prove existence and uniqueness of solutions to equations of type (3.1) on bounded Lipschitz domains $\mathcal{O} \subseteq \mathbb{R}^d$ in the spaces $\mathfrak{H}_{p, \theta}^{\gamma, q}(\mathcal{O}, T)$. We are mainly interested in the case $q > p$. The main ingredient will be Corollary 3.9 which allows us to lift up the regularity of the solution

once we have established a certain low $L_q(L_p)$ -regularity and if f and the g^k 's have high $L_q(L_p)$ -regularity. In this section we restrict ourselves to equations of type (3.1) with $\sigma \equiv 0$ and vanishing initial condition, i.e., we consider the problem

$$du = (a^{ij}u_{x^i x^j} + f)dt + g^k dw_t^k, \quad u(0, \cdot) = 0. \quad (4.1)$$

We expect, however, that the lifting argument in Corollary 3.9 can be used to derive similar results for general equations of type (3.1). We establish existence of solutions with low $L_q(L_p)$ -regularity in two different ways which correspond to two different restrictions in our assumptions. First, in Subsection 4.1 we consider Lipschitz domains with sufficiently small Lipschitz constants. Here we use an $L_q(L_p)$ -regularity result for deterministic PDEs and basic estimates for stochastic integrals in UMD Banach spaces to derive a result for general integrability parameters $q \geq p \geq 2$. Then, in Subsection 4.2 we consider the case of general bounded Lipschitz domains. Applying techniques from the semigroup approach to stochastic evolution equations in Banach spaces, we are able to prove existence and uniqueness of solutions of the stochastic heat equation in $\mathfrak{H}_{p,d}^{\gamma,q}(\mathcal{O}, T)$ for integrability parameters $p \in [2, p_0)$ and $q \geq p$.

4.1 A result for domains with small Lipschitz constant

We need the following result concerning existence and uniqueness of solutions to SPDEs of the form (3.1) in $\mathfrak{H}_{p,\theta}^\gamma(\mathcal{O}, T) = \mathfrak{H}_{p,\theta}^{\gamma,p}(\mathcal{O}, T)$, i.e., for the case $p = q$. It is taken from [26], see Theorem 2.12 and Remark 2.13 therein. Note that it also holds under weaker assumptions on the parameters and for more general equations than stated here.

Theorem 4.1. *Let \mathcal{O} be a bounded Lipschitz domain in \mathbb{R}^d and $\gamma \in \mathbb{R}$. For $i, j \in \{1, \dots, d\}$ and $k \in \mathbb{N}$, let a^{ij} , σ^{ik} be given coefficients satisfying Assumption 3.6.*

(i) *For $p \in [2, \infty)$, there exists a constant $\kappa_0 \in (0, 1)$, depending only on d, p, δ_0, K and \mathcal{O} , such that for any $\theta \in (d + p - 2 - \kappa_0, d + p - 2 + \kappa_0)$, $f \in \mathbb{H}_{p,\theta+p}^\gamma(\mathcal{O}, T)$, $g \in \mathbb{H}_{p,\theta}^{\gamma+1}(\mathcal{O}, T; \ell_2)$ and $u_0 \in U_{p,\theta}^{\gamma+2}(\mathcal{O})$, Eq. (3.1) has a unique solution u in the class $\mathfrak{H}_{p,\theta}^{\gamma+2}(\mathcal{O}, T)$. For this solution*

$$\|u\|_{\mathfrak{H}_{p,\theta}^{\gamma+2}(\mathcal{O}, T)}^p \leq N \left(\|f\|_{\mathbb{H}_{p,\theta+p}^\gamma(\mathcal{O}, T)}^p + \|g\|_{\mathbb{H}_{p,\theta}^{\gamma+1}(\mathcal{O}, T; \ell_2)}^p + \|u_0\|_{U_{p,\theta}^{\gamma+2}(\mathcal{O})}^p \right), \quad (4.2)$$

where the constant N depends only on $d, p, \gamma, \theta, \delta_0, K, T$ and \mathcal{O} .

(ii) *There exists $p_0 > 2$, such that the following statement holds: if $p \in [2, p_0)$, then there exists a constant $\kappa_1 \in (0, 1)$, depending only on d, p, δ_0, K and \mathcal{O} , such that for any $\theta \in (d - \kappa_1, d + \kappa_1)$, $f \in \mathbb{H}_{p,\theta+p}^\gamma(\mathcal{O}, T)$, $g \in \mathbb{H}_{p,\theta}^{\gamma+1}(\mathcal{O}, T; \ell_2)$ and $u_0 \in U_{p,\theta}^{\gamma+2}(\mathcal{O})$, Eq. (3.1) has a unique solution u in the class $\mathfrak{H}_{p,\theta}^{\gamma+2}(\mathcal{O}, T)$. For this solution, estimate (4.2) holds.*

Here is the main result of this subsection.

Theorem 4.2. *Let \mathcal{O} be a bounded Lipschitz domain in \mathbb{R}^d and let $\gamma \geq 0$. For given coefficients a^{ij} , $i, j \in \{1, \dots, d\}$, let Assumption 3.6 be satisfied with $\sigma \equiv 0$. Then, for $2 \leq p \leq q < \infty$, there exists a constant $c = c(d, p, q, \delta_0, K)$, such that, if the Lipschitz constant K_0 in Definition 2.5 satisfies $K_0 \leq c$, the following holds: For*

$$f \in \mathbb{H}_{p,d+p}^{\gamma,q}(\mathcal{O}, T) \cap \mathbb{H}_{p,d}^{0,q}(\mathcal{O}, T) \quad \text{and} \quad g \in \mathbb{H}_{p,d}^{\gamma+1,q}(\mathcal{O}, T; \ell_2) \cap \mathbb{H}_{p,d-p}^{1,q}(\mathcal{O}, T; \ell_2),$$

Eq. (4.1) has a unique solution $u \in \mathfrak{H}_{p,d}^{\gamma+2,q}(\mathcal{O}, T)$. Moreover, there exists a constant $N \in (0, \infty)$, which does not depend on u , f and g , such that

$$\begin{aligned} \|u\|_{\mathfrak{H}_{p,d}^{\gamma+2,q}(\mathcal{O}, T)} &\leq N \left(\|f\|_{\mathbb{H}_{p,d+p}^{\gamma,q}(\mathcal{O}, T)} + \|f\|_{\mathbb{H}_{p,d}^{0,q}(\mathcal{O}, T)} \right. \\ &\quad \left. + \|g\|_{\mathbb{H}_{p,d}^{\gamma+1,q}(\mathcal{O}, T; \ell_2)} + \|g\|_{\mathbb{H}_{p,d-p}^{1,q}(\mathcal{O}, T; \ell_2)} \right). \end{aligned} \quad (4.3)$$

Remark 4.3. We note that every bounded C^1 -domain $\mathcal{O} \subseteq \mathbb{R}^d$ is a Lipschitz domain where the Lipschitz constant K_0 in Definition 2.5 can be chosen arbitrarily small. Therefore, the assertion of Theorem 4.2 holds for any bounded C^1 -domain.

Proof of Theorem 4.2. Since $\mathfrak{H}_{p,d}^{\gamma+2,q}(\mathcal{O}, T) \hookrightarrow \mathfrak{H}_{2,d}^{\gamma+2}(\mathcal{O}, T)$, the uniqueness follows from Theorem 4.1. Also by Theorem 4.1, there exists a solution $u \in \mathfrak{H}_{2,d}^{\gamma+2}(\mathcal{O}, T)$. We only need to show that $u \in \mathfrak{H}_{p,d}^{\gamma+2,q}(\mathcal{O}, T)$ and that it satisfies (4.3). For all $\varphi \in C_0^\infty(\mathcal{O})$, with probability one,

$$(u(t), \varphi) = \int_0^t (a^{ij}(s)u_{x^i x^j}(s) + f(s), \varphi) ds + \sum_{k=1}^{\infty} \int_0^t (g^k(s), \varphi) dw_s^k, \quad t \in [0, T].$$

Let us define

$$w(t) := \sum_{k=1}^{\infty} \int_0^t g^k(s) dw_s^k$$

as an infinite sum of, say $H_{2,d-2}^1(\mathcal{O})$ -valued stochastic integrals. This sum converges in the space $\mathbb{H}_{2,d-2}^1(\mathcal{O}, T)$ due to Itô's isometry and since $g \in \mathbb{H}_{p,d-p}^{1,q}(\mathcal{O}, T; \ell_2) \hookrightarrow \mathbb{H}_{2,d-2}^1(\mathcal{O}, T; \ell_2)$. We fix a continuous modification of the $H_{2,d-2}^1(\mathcal{O})$ -valued process $(w(t))_{t \in [0, T]}$ from now on, which is well-known to exist. For all $\varphi \in C_0^\infty(\mathcal{O})$, with probability one,

$$\sum_{k=1}^{\infty} \int_0^t (g^k(s), \varphi) dw_s^k = (w(t), \varphi), \quad t \in [0, T].$$

Therefore, setting $\bar{u} := u - w$, we know that for all $\varphi \in C_0^\infty(\mathcal{O})$, with probability one,

$$(\bar{u}(t), \varphi) = \int_0^t (a^{ij}(s)\bar{u}_{x^i x^j}(s) + f(s) + a^{ij}(s)w_{x^i x^j}(s), \varphi) ds, \quad t \in [0, T]. \quad (4.4)$$

It follows that \bar{u} is the unique solution in $\mathfrak{H}_{2,d}^1(\mathcal{O}, T)$ to

$$d\bar{u} = (a^{ij}\bar{u}_{x^i x^j} + f + a^{ij}w_{x^i x^j})dt, \quad \bar{u}(0) = 0 \quad (4.5)$$

We are going to consider (4.5) ω -wise and apply an $L_q(L_p)$ -regularity result for deterministic PDEs from [16]. To this end, we have to check that in the present situation our notion of a solution fits to the one described therein. Since $\bar{u}, w \in \mathbb{H}_{2,d-2}^1(\mathcal{O}, T)$ and $f \in \mathbb{H}_{p,d}^{0,q}(\mathcal{O}, T) \hookrightarrow \mathbb{H}_{2,d}^0(\mathcal{O}, T)$, we know that, for almost every $\omega \in \Omega$, the mappings $t \mapsto \bar{u}(\omega, t, \cdot)$ and $t \mapsto w(\omega, t, \cdot)$ belong to $L_2([0, T]; H_{2,d-2}^1(\mathcal{O}))$ and $t \mapsto f(\omega, t, \cdot)$ belongs to $L_2([0, T]; L_2(\mathcal{O}))$. In particular, by Lemma 2.1, $\bar{u}_{x^i x^j}(\omega)$ and $w_{x^i x^j}(\omega)$ belong to $L_2([0, T]; H_{2,d+2}^{-1}(\mathcal{O}))$ for all $i, j = 1, \dots, d$, so that

$$\int_0^t \{a^{ij}(\omega, s)\bar{u}_{x^i x^j}(\omega, s) + f(\omega, s) + a^{ij}(\omega, s)w_{x^i x^j}(\omega, s)\} ds =: \int_0^t \Phi(\omega, s) ds$$

exists as an $H_{2,d+2}^{-1}(\mathcal{O})$ -valued Bochner integral. This and (4.4) imply that, for almost all $\omega \in \Omega$,

$$(\bar{u}(\omega, t), \varphi_k) = \left(\int_0^t \Phi(\omega, s) ds, \varphi_k \right), \quad k \in \mathbb{N}, \quad t \in [0, T],$$

where $\{\varphi_k\}_{k \in \mathbb{N}} \subseteq C_0^\infty(\mathcal{O})$ is supposed to be dense in $H_{2,d-2}^1(\mathcal{O})$. As a consequence,

$$\bar{u}(\omega, t) = \int_0^t \Phi(\omega, s) ds, \quad t \in [0, T].$$

Standard arguments lead to

$$\int_0^T \phi'(t) \bar{u}(\omega, t) dt = - \int_0^T \phi(t) \Phi(\omega, t) dt + \phi(T) \bar{u}(\omega, T), \quad \phi \in C_0^\infty(\mathbb{R}),$$

where the integrals are $H_{2,d+2}^{-1}(\mathcal{O})$ -valued Bochner integrals. We obtain

$$\begin{aligned} & \int_0^T \int_{\mathcal{O}} \bar{u}(\omega) \frac{\partial}{\partial t} h \, dx \, dt \\ &= \int_0^T \int_{\mathcal{O}} (a^{ij}(\omega) \bar{u}_{x_j}(\omega) h_{x_i} - f(\omega) h + a^{ij}(\omega) w_{x_j}(\omega) h_{x_i}) \, dx \, dt + \int_{\mathcal{O}} \bar{u}(\omega, T) h(T) \, dx \end{aligned} \quad (4.6)$$

for all test-functions $h = \phi \otimes \varphi$, $\phi \in C_0^\infty(\mathbb{R})$, $\varphi \in C_0^\infty(\mathcal{O})$. Using approximation arguments one can verify that (4.6) even holds for all test-functions h which belong to the space $\mathcal{H}_{2,2}^1((0, T) \times \mathcal{O})$ considered in [16] and which vanish on $(0, T) \times \partial\mathcal{O}$ in the sense that $h(t) \in \dot{W}_2^1(\mathcal{O}) = H_{2,d-2}^1(\mathcal{O})$ for almost all $t \in (0, T)$. Moreover, for almost all $\omega \in \Omega$, $\bar{u}(\omega)$ belongs to the space $\dot{\mathcal{H}}_{2,2}^1((0, T) \times \mathcal{O})$ as defined in [16] and it vanishes on $(0, T) \times \partial\mathcal{O}$. Thus, for almost all $\omega \in \Omega$, $\bar{u}(\omega)$ is the unique solution in $\dot{\mathcal{H}}_{2,2}^1((0, T) \times \mathcal{O})$ to

$$\begin{cases} -\frac{\partial}{\partial t} v + a^{ij}(\omega) v_{x_i x_j} = \operatorname{div}(a^{1j}(\omega) w_{x_j}(\omega), \dots, a^{dj}(\omega) w_{x_j}(\omega))^T - f(\omega) & \text{in } (0, T) \times \mathcal{O} \\ v = 0 & \text{on } (0, T) \times \partial\mathcal{O} \end{cases}$$

in the sense of [16]. Now we can apply [16, Theorem 8.1] and use the fact that the coefficients a^{ij} are uniformly bounded due to Assumption 3.6 to obtain

$$\|D\bar{u}(\omega)\|_{L_q([0,T];L_p(\mathcal{O}))} \leq N (\|Dw(\omega)\|_{L_q([0,T];L_p(\mathcal{O}))} + \|f(\omega)\|_{L_q([0,T];L_p(\mathcal{O}))}) \quad (4.7)$$

for almost all $\omega \in \Omega$, where the constant N does not depend on ω . We remark that the assumption on the Lipschitz constant K_0 comes into play at this point: Theorem 8.1 in [16] implies that there exists a constant $c = c(d, p, q, \delta_0, K)$ such that, if $K_0 \leq c$, then estimate (4.7) holds. Integration w.r.t. \mathbb{P} and Hardy's inequality yield

$$\begin{aligned} \|u\|_{\mathbb{H}_{p,d-p}^{0,q}(\mathcal{O},T)} &\leq N \|Du\|_{\mathbb{H}_{p,d}^{0,q}(\mathcal{O},T)} \\ &\leq N (\|D\bar{u}\|_{\mathbb{H}_{p,d}^{0,q}(\mathcal{O},T)} + \|Dw\|_{\mathbb{H}_{p,d}^{0,q}(\mathcal{O},T)}) \\ &\leq N (\|f\|_{\mathbb{H}_{p,d}^{0,q}(\mathcal{O},T)} + \|Dw\|_{\mathbb{H}_{p,d}^{0,q}(\mathcal{O},T)}). \end{aligned} \quad (4.8)$$

The term $\|Dw\|_{\mathbb{H}_{p,d}^{0,q}(\mathcal{O},T)}$ can be estimated with the help of an inequality for stochastic integrals in UMD Banach spaces with type 2 taken from [46]. To this end, let $\gamma(\ell_2, H_{p,d-p}^1(\mathcal{O}))$ denote the Banach space of γ -radonifying operators from ℓ_2 to $H_{p,d-p}^1(\mathcal{O})$, see [45] for a survey on this class of operators. Furthermore, let $\{\mathbf{e}_k\}_{k \in \mathbb{N}}$ be the standard orthonormal basis of the Hilbert space ℓ_2 . Then, the stochastic process

$$b : \Omega \times [0, T] \rightarrow \gamma(\ell_2, H_{p,d-p}^1(\mathcal{O})),$$

given by

$$b(\omega, t)\mathbf{a} := \sum_{k=1}^{\infty} \langle \mathbf{a}, \mathbf{e}_k \rangle_{\ell_2} g^k(\omega, t), \quad \mathbf{a} \in \ell_2,$$

is well-defined, and

$$b \in L_q(\Omega \times [0, T], \mathcal{P}, \mathbb{P} \otimes dt; \gamma(\ell_2, H_{p,d-p}^1(\mathcal{O}))).$$

Moreover,

$$\|b\|_{L_q(\Omega \times [0, T]; \gamma(\ell_2, H_{p,d-p}^1(\mathcal{O})))} \sim \|g\|_{\mathbb{H}_{p,d-p}^{1,q}(\mathcal{O}, T; \ell_2)}, \quad (4.9)$$

see [4, Remark 3.7]. We remark that $H_{p,d-p}^1(\mathcal{O}) = \mathring{W}_p^1(\mathcal{O})$ is a UMD Banach spaces with type 2, since it is a closed subspace of the classical Sobolev space $W_p^1(\mathcal{O})$. Therefore, by [46, Corollary 3.10], $(b(t))_{t \in [0, T]}$ is L_q -stochastically integrable w.r.t. the ℓ_2 -cylindrical Brownian motion $(W_{\ell_2}(t))_{t \in [0, T]}$ given by

$$W_{\ell_2}(t)\mathbf{a} := \sum_{k=1}^{\infty} \langle \mathbf{a}, \mathbf{e}_k \rangle_{\ell_2} w_t^k, \quad \mathbf{a} \in \ell_2.$$

By [46, Corollary 3.9], for every $t \in [0, T]$,

$$\int_0^t b(s) dW_{\ell_2}(s) = \sum_{k=1}^{\infty} \int_0^t g^k(s) dw_s^k = w(t) \quad \text{a.s.}, \quad (4.10)$$

where the series converges in $L_q(\Omega; H_{p,d-p}^1(\mathcal{O}))$. Thus, we can apply [46, Corollary 3.10] and obtain

$$\|w\|_{\mathbb{H}_{p,d-p}^{1,q}(\mathcal{O}, T)}^q \leq N \|b\|_{L_q(\Omega \times [0, T]; \gamma(\ell_2, H_{p,d-p}^1(\mathcal{O})))}^q \leq N \|g\|_{\mathbb{H}_{p,d-p}^{1,g}(\mathcal{O}, T; \ell_2)}^q,$$

where we used (4.9) for the last estimate. Now let e_1, \dots, e_d be the standard orthonormal basis of \mathbb{R}^d . Note that for every $i \in \{1, \dots, d\}$, D^{e_i} is a linear and bounded operator from $H_{p,d-p}^1(\mathcal{O})$ to the UMD space $L_p(\mathcal{O})$, cf. Lemma 2.1(iii) and (iv). Using this and similar arguments as above we obtain

$$D^{e_i} w(t) = \int_0^t D^{e_i} \circ b(s) dW_{\ell_2}(s),$$

with $D^{e_i} \circ b$ denoting the (point-wise) composition of the operators D^{e_i} and b , and

$$\begin{aligned}
\|Dw\|_{\mathbb{H}_{p,d}^{0,q}(\mathcal{O},T)} &= \sum_{i=1}^d \|D^{e_i}w\|_{\mathbb{H}_{p,d}^{0,q}(\mathcal{O},T)} \\
&= \sum_{i=1}^d \left\| \int_0^{(\cdot)} D^{e_i} \circ b(s) dW_{\ell_2}(s) \right\|_{L_q(\Omega \times [0,T]; L_p(\mathcal{O}))} \\
&\leq N \sum_{i=1}^d \|D^{e_i} \circ b\|_{L_q(\Omega \times [0,T]; \gamma(\ell_2; L_p(\mathcal{O})))} \\
&\leq N \|b\|_{L_q(\Omega \times [0,T]; \gamma(\ell_2; H_{p,d-p}^1(\mathcal{O})))} \\
&\leq N \|g\|_{\mathbb{H}_{p,d-p}^{1,q}(\mathcal{O},T; \ell_2)}.
\end{aligned} \tag{4.11}$$

Combining (4.8) and (4.11) and applying Corollary 3.9 finishes the proof if $q = mp$ with $m \in \mathbb{N}$. We can get rid of this restriction by following the lines of [31, Proof of Theorem 2.1, p. 7] and applying Marcinkiewicz's interpolation theorem. \square

4.2 An $L_q(L_p)$ -theory of the heat equation on general bounded Lipschitz domains

In this subsection we present a first $L_q(L_p)$ -theory for the stochastic heat equation

$$du = (\Delta u + f) dt + g^k dw_t^k, \quad u(0, \cdot) = 0, \tag{4.12}$$

on general bounded Lipschitz domains $\mathcal{O} \subseteq \mathbb{R}^d$. We start by presenting the main result of this subsection, which we will prove later on.

Theorem 4.4. *Let \mathcal{O} be a bounded Lipschitz domain in \mathbb{R}^d and let $\gamma \geq 0$. There exists an exponent p_0 with $p_0 > 3$ when $d \geq 3$ and $p_0 > 4$ when $d = 2$ such that for $p \in [2, p_0)$ and $p \leq q < \infty$, Eq. (4.12) has a unique solution $u \in \mathfrak{H}_{p,d}^{\gamma+2,q}(\mathcal{O}, T)$, provided*

$$f \in \mathbb{H}_{p,d+p}^{\gamma,q}(\mathcal{O}, T) \cap \mathbb{H}_{p,d}^{0,q}(\mathcal{O}, T) \quad \text{and} \quad g \in \mathbb{H}_{p,d}^{\gamma+1,q}(\mathcal{O}, T; \ell_2) \cap \mathbb{H}_{p,d-p}^{1,q}(\mathcal{O}, T; \ell_2).$$

Moreover, there exists a constant $N \in (0, \infty)$, which does not depend on f and g , such that

$$\|u\|_{\mathbb{H}_{p,d-p}^{\gamma+2,q}(\mathcal{O},T)}^q \leq N \left(\|f\|_{\mathbb{H}_{p,d}^{0,q}(\mathcal{O},T)}^q + \|f\|_{\mathbb{H}_{p,d+p}^{\gamma,q}(\mathcal{O},T)}^q + \|g\|_{\mathbb{H}_{p,d}^{\gamma+1,q}(\mathcal{O},T; \ell_2)}^q \right). \tag{4.13}$$

For bounded C^1 -domains $G \subseteq \mathbb{R}^d$ this result has been already proven in [25]. Unfortunately, the techniques used there will not work if the boundary is assumed to be just Lipschitz continuous. Therefore, we choose to take another way. We will mainly use the fact that the domain of the square root of the negative weak Dirichlet-Laplacian on $L_p(\mathcal{O})$ coincides with the closure of the test functions in the $L_p(\mathcal{O})$ -Sobolev space of order one, at least for the range of p allowed in our assertion. Before we prove this fact let us get more precise and introduce some notations and definitions.

Let \mathcal{O} be a bounded Lipschitz domain in \mathbb{R}^d . As in [49, Definition 3.1], for arbitrary $p \in (1, \infty)$, we define the weak Dirichlet-Laplacian $\Delta_{p,w}^D$ on $L_p(\mathcal{O})$ as follows:

$$\begin{aligned} D(\Delta_{p,w}^D) &:= \{u \in \mathring{W}_p^1(\mathcal{O}) : \Delta u \in L_p(\mathcal{O})\}, \\ \Delta_{p,w}^D u &:= \Delta u = \delta_{ij} u_{x^i x^j}, \end{aligned}$$

where δ_{ij} denotes the Kronecker symbol. If we fix $p \in (p_0/(p_0 - 1), p_0)$ with $p_0 = 4 + \delta$ when $d = 2$ and $p_0 = 3 + \delta$ when $d \geq 3$ where $\delta > 0$ is taken from [49, Proposition 4.1], then, the unbounded operator $\Delta_{p,w}^D$ generates a strongly continuous, analytic semigroup $\{S_p(t)\}_{t \geq 0}$ of contractions on $L_p(\mathcal{O})$, see [49, Theorem 3.8 and Corollary 4.2]. Thus, $(-\Delta_{p,w}^D)^{1/2}$, the square root of the negative of $\Delta_{p,w}^D$, can be defined as the inverse of the operator

$$(-\Delta_{p,w}^D)^{-1/2} := \pi^{-1/2} \int_0^\infty t^{-1/2} S_p(t) dt : L_p(\mathcal{O}) \rightarrow L_p(\mathcal{O}) \quad (4.14)$$

with domain

$$D((-\Delta_{p,w}^D)^{1/2}) := \text{Range}((-\Delta_{p,w}^D)^{-1/2}),$$

see [41, Chapter 2.6]. Endowed with the norm

$$\|u\|_{D((-\Delta_{p,w}^D)^{1/2})} := \|(-\Delta_{p,w}^D)^{1/2} u\|_{L_p(\mathcal{O})}, \quad u \in D((-\Delta_{p,w}^D)^{1/2}),$$

$D((-\Delta_{p,w}^D)^{1/2})$ becomes a Banach space. Exploiting the fundamental results from [49] and [19], we can prove the following identity, which is crucial if we want to apply the results from [47] in our setting.

Lemma 4.5. *Let \mathcal{O} be a bounded Lipschitz domain in \mathbb{R}^d . There is an exponent p_0 with $p_0 > 4$ when $d = 2$ and $p_0 > 3$ when $d \geq 3$ such that if $p \in [2, p_0)$*

$$D((-\Delta_{p,w}^D)^{1/2}) = \mathring{W}_p^1(\mathcal{O})$$

with equivalent norms.

Proof. We fix $p \in [2, p_0)$ with $p_0 = 4 + \delta$ when $d = 2$ and $p_0 = 3 + \delta$ when $d \geq 3$ where $\delta > 0$ is taken from [49, Proposition 4.1]. We start with the proof of the embedding

$$D((-\Delta_{p,w}^D)^{1/2}) \hookrightarrow \mathring{W}_p^1(\mathcal{O}). \quad (4.15)$$

By [49, Proposition 4.1] the semigroups $\{S_2(t)\}_{t \geq 0}$ and $\{S_p(t)\}_{t \geq 0}$ are consistent, i.e., for all $t \geq 0$,

$$S_2(t)f = S_p(t)f, \quad f \in L_p(\mathcal{O}).$$

Using (4.14), this leads to

$$(-\Delta_{p,w}^D)^{-1/2} f = (-\Delta_{2,w}^D)^{-1/2} f, \quad f \in L_p(\mathcal{O}),$$

which implies the boundedness of the operator

$$(-\Delta_{p,w}^D)^{-1/2} : L_p(\mathcal{O}) \rightarrow \mathring{W}_p^1(\mathcal{O}),$$

since the restriction of $(-\Delta_{2,w}^D)^{-1/2}$ to $L_p(\mathcal{O})$ is bounded from $L_p(\mathcal{O})$ to $\dot{W}_p^1(\mathcal{O})$, see [19, Theorem 7.5(a)]. Thus

$$\|(-\Delta_{p,w}^D)^{-1/2}f\|_{\dot{W}_p^1(\mathcal{O})} \leq N \|f\|_{L_p(\mathcal{O})}, \quad f \in L_p(\mathcal{O}).$$

Consequently, for any $g \in \text{Range}((-\Delta_{p,w}^D)^{-1/2}) = D((-\Delta_{p,w}^D)^{1/2})$,

$$\|g\|_{\dot{W}_p^1(\mathcal{O})} \leq N \|(-\Delta_{p,w}^D)^{1/2}g\|_{L_p(\mathcal{O})} = N \|g\|_{D((-\Delta_{p,w}^D)^{1/2})},$$

with a constant N independent of g . Embedding (4.15) follows.

It remains to prove the converse direction, i.e.,

$$\dot{W}_p^1(\mathcal{O}) \hookrightarrow D((-\Delta_{p,w}^D)^{1/2}).$$

To this end, we first notice that the strongly continuous analytic contraction-semigroup $\{S_p(t)\}_{t \geq 0}$ on $L_p(\mathcal{O})$ is positive in the sense of [17, p. 353], see [49, Lemma 4.4]. Therefore, by [22, Corollary 5.2], $(-\Delta_{p,w}^D)$ has a bounded H^∞ -calculus of angle less than $\pi/2$. Consequently, it has bounded imaginary powers. This implies

$$[L_p(\mathcal{O}), D(-\Delta_{p,w}^D)]_{1/2} = D((-\Delta_{p,w}^D)^{1/2}), \quad (4.16)$$

see [43, Theorem 1.15.3]. By Remark 2.6 and [38, Proposition 2.4],

$$\dot{W}_p^1(\mathcal{O}) = H_{p,d-p}^1(\mathcal{O}) = [H_{p,d}^0(\mathcal{O}), H_{p,d-2p}^2(\mathcal{O})]_{1/2}.$$

Also, by the definition of the weighted Sobolev spaces one easily sees that

$$H_{p,d}^0(\mathcal{O}) = L_p(\mathcal{O}) \quad \text{and} \quad H_{p,d-2p}^2(\mathcal{O}) \hookrightarrow D(-\Delta_{p,w}^D).$$

Combining these results, we obtain

$$\dot{W}_p^1(\mathcal{O}) = [H_{p,d}^0(\mathcal{O}), H_{p,d-2p}^2(\mathcal{O})]_{1/2} \hookrightarrow [L_p(\mathcal{O}), D(-\Delta_{p,w}^D)]_{1/2} = D((-\Delta_{p,w}^D)^{1/2}),$$

which finishes the proof. \square

Now we are ready to prove our main result in this subsection.

Proof of Theorem 4.4. As in the proof of Lemma 4.5 we fix $p \in [2, p_0)$ with $p_0 = 4 + \delta$ when $d = 2$ and $p_0 = 3 + \delta$ when $d \geq 3$ where $\delta > 0$ is taken from [49, Proposition 4.1]. We start with the case $\gamma = 0$. I.e., we assume that $f \in \mathbb{H}_{p,d}^{0,q}(\mathcal{O}, T)$ and $g \in \mathbb{H}_{p,d-p}^{1,q}(\mathcal{O}, T; \ell_2)$. Recall the corresponding notations $(b(t))_{t \in [0, T]}$, $(w(t))_{t \in [0, T]}$ and W_{ℓ_2} introduced in the proof of Theorem 4.2 and their properties. By Lemma 4.5 and Eq. (4.16) we have

$$[L_p(\mathcal{O}), D(-\Delta_{p,w}^D)]_{1/2} = H_{p,d-p}^1(\mathcal{O}).$$

Also, $D(\Delta_{p,w}^D) \hookrightarrow L_p(\mathcal{O})$ densely, since $C_0^\infty(\mathcal{O})$ is contained in $D(\Delta_{p,w}^D)$. Therefore, we can apply [47, Theorem 4.5(ii)] and obtain the existence of a stochastic process

$$u \in L_q(\Omega \times [0, T], \mathcal{P}, \mathbb{P} \otimes dt; D(-\Delta_{p,w}^D)) \quad (4.17)$$

solving the stochastic evolution equation

$$\begin{cases} du(t) - \Delta_{p,w}^D u(t) dt = f(t) dt + b(t) dW_{\ell_2}(t), & t \in [0, T] \\ u(0) = 0 \end{cases}$$

in the sense of [47, Definition 4.2] with $X_0 := L_p(\mathcal{O})$. Moreover, there exists a version \tilde{u} of u , such that the following equality is fulfilled in $L_p(\mathcal{O})$ a.s. for all $t \in [0, T]$ at once:

$$\tilde{u}(t) = \int_0^t \Delta \tilde{u}(s) ds + \int_0^t f(s) ds + \int_0^t b(s) dW_{\ell_2}(s).$$

We can fix a continuous versions of the stochastic process $(w(t))_{t \in [0, T]}$, so that by (4.10) a.s.

$$\tilde{u}(t) = \int_0^t \Delta \tilde{u}(s) ds + \int_0^t f(s) ds + w(t) \quad \text{for all } t \in [0, T] \text{ in } L_p(\mathcal{O}).$$

Therefore, by Lemma 4.5 and (4.17), $u \in \mathfrak{H}_{p,d}^{1,q}(\mathcal{O}, T)$ and solves Eq. (4.12) in the sense of Definition 3.4. Since $\mathfrak{H}_{p,d}^{1,q}(\mathcal{O}, T) \hookrightarrow \mathfrak{H}_{2,d}^{1,2}(\mathcal{O}, T)$, the uniqueness follows from [26, Theorem 2.12]. Thus, in order to finish the proof of the basic case $\gamma = 0$, we show the a priori estimate

$$\|u\|_{\mathbb{H}_{p,d-p}^{1,q}(\mathcal{O}, T)}^q \leq N \left(\|f\|_{\mathbb{H}_{p,d}^{0,q}(\mathcal{O}, T)}^q + \|g\|_{\mathbb{H}_{p,d}^{1,q}(\mathcal{O}, T; \ell_2)}^q \right), \quad (4.18)$$

which implies estimate (4.13) for $\gamma = 0$. To this end we will use the fact that the stochastic process $V : [0, T] \times \Omega \rightarrow L_p(\mathcal{O})$ defined as

$$V(t) := \int_0^t S_p(t-s) f(s) ds + \int_0^t S_p(t-s) b(s) dW_{\ell_2}(s), \quad t \in [0, T],$$

is a version of u , see [47, Proposition 4.4]. Since $-\Delta_{p,w}^D$ has the (deterministic) maximal regularity property (see [49, Proposition 6.1]) and $0 \in \rho(\Delta_{p,w}^D)$, we obtain

$$\mathbb{E} \left[\left\| t \mapsto \int_0^t S_p(t-s) f(s) ds \right\|_{L_q(0, T; H_{p,d-p}^1(\mathcal{O}))}^q \right] \leq N \|f\|_{\mathbb{H}_{p,d}^{0,q}(\mathcal{O}, T)}^q, \quad (4.19)$$

where we used again Lemma 4.5 and [35, Theorem 9.7]. Simultaneously, notice that $-\Delta_{p,w}^D$ and g (respectively b) fulfil the assumptions of [48, Theorem 1.1]; we have already checked them in our explanations above. Thus, applying this result, we obtain

$$\mathbb{E} \left[\left\| t \mapsto \int_0^t S_p(t-s) b(s) dW_{\ell_2}(s) \right\|_{L_q(0, T; H_{p,d-p}^1(\mathcal{O}))}^q \right] \leq N \|g\|_{\mathbb{H}_{p,d}^{0,q}(\mathcal{O}, T; \ell_2)}^q. \quad (4.20)$$

The constants in (4.19) and (4.20) do not depend on f and g . Therefore, using the last two estimates we obtain the existence of a constant N , independent of f or g , such that

$$\|V\|_{\mathbb{H}_{p,d-p}^{1,q}(\mathcal{O}, T)}^q \leq N \left(\|f\|_{\mathbb{H}_{p,d}^{0,q}(\mathcal{O}, T)}^q + \|g\|_{\mathbb{H}_{p,d}^{0,q}(\mathcal{O}, T; \ell_2)}^q \right).$$

Since V is just a version of the solution u , Eq. (4.18) follows.

In order to prove the assertion for arbitrary $\gamma \geq 0$ and $2 \leq p \leq q < \infty$ we can argue as we have done at the end of the proof of Theorem 4.2. \square

5 Hölder-Sobolev regularity of elements of $\mathfrak{H}_{p,\theta}^{\gamma,q}(\mathcal{O}, T)$ and implications for SPDEs

In this section we analyse the temporal Hölder regularity of functions in $\mathfrak{H}_{p,\theta}^{\gamma,q}(\mathcal{O}, T)$, where \mathcal{O} is a bounded Lipschitz domain in \mathbb{R}^d . Our main interest lies on the case $q \neq p$. As an application, we obtain Hölder-Sobolev regularity for the solutions to SPDEs presented in Section 4. In combination with the Sobolev type embeddings for the spaces $H_{p,\theta}^{\gamma}(\mathcal{O})$ from the Section 2, we also obtain assertions concerning the Hölder regularity in time and space for elements of $\mathfrak{H}_{p,\theta}^{\gamma,q}(\mathcal{O}, T)$ (Corollary 5.2). Here is the main result of this section.

Theorem 5.1. *Let \mathcal{O} be a bounded Lipschitz domain in \mathbb{R}^d . Let $2 \leq p \leq q < \infty$, $\gamma \in \mathbb{N}$, $\theta \in \mathbb{R}$, and $u \in \mathfrak{H}_{p,\theta}^{\gamma,q}(\mathcal{O}, T)$. Moreover, let*

$$2/q < \bar{\beta} < \beta \leq 1.$$

Then there exists a constant N , which does not depend on T and u , such that

$$\begin{aligned} & \mathbb{E}[\psi^{\beta-1}u]_{C^{\bar{\beta}/2-1/q}([0,T];H_{p,\theta}^{\gamma-\beta}(\mathcal{O}))}^q \\ & \leq NT^{(\beta-\bar{\beta})q/2} \left(\|u\|_{\mathbb{H}_{p,\theta-p}^{\gamma,q}(\mathcal{O},T)}^q + \|\mathbb{D}u\|_{\mathbb{H}_{p,\theta+p}^{\gamma-2,q}(\mathcal{O},T)}^q + \|\mathbb{S}u\|_{\mathbb{H}_{p,\theta}^{\gamma-1,q}(\mathcal{O},T;\ell_2)}^q \right), \end{aligned} \quad (5.1)$$

and

$$\begin{aligned} & \mathbb{E}\|\psi^{\beta-1}u\|_{C^{\bar{\beta}/2-1/q}([0,T];H_{p,\theta}^{\gamma-\beta}(\mathcal{O}))}^q \leq NT^{(\beta-\bar{\beta})q/2} \left(\mathbb{E}\|\psi^{\beta-1}u(0, \cdot)\|_{H_{p,\theta}^{\gamma-\beta}(\mathcal{O})}^q + \right. \\ & \left. \|u\|_{\mathbb{H}_{p,\theta-p}^{\gamma,q}(\mathcal{O},T)}^q + \|\mathbb{D}u\|_{\mathbb{H}_{p,\theta+p}^{\gamma-2,q}(\mathcal{O},T)}^q + \|\mathbb{S}u\|_{\mathbb{H}_{p,\theta}^{\gamma-1,q}(\mathcal{O},T;\ell_2)}^q \right). \end{aligned} \quad (5.2)$$

Theorem 5.1 and Lemma 2.1(ii) yield the following so-called interior Schauder estimates of functions in $\mathfrak{H}_{p,\theta}^{\gamma,q}(\mathcal{O}, T)$.

Corollary 5.2. *Given the setting of Theorem 5.1, let $u \in \mathfrak{H}_{p,\theta}^{\gamma,q}(\mathcal{O}, T)$ and $\gamma - \beta - d/p = k + \varepsilon$ where $k \in \mathbb{N}_0$ and $\varepsilon \in (0, 1]$. Then for $\nu := \beta - 1 + \theta/p$ and multi-indices $i, j \in \mathbb{N}_0^d$ such that $|i| \leq k$ and $|j| = k$, we have*

$$\begin{aligned} & \mathbb{E} \left[\sup_{t \neq s} |t - s|^{-(\bar{\beta}q/2-1)} \left(|\psi^{\nu+|i|} D^i(u(t, \cdot) - u(s, \cdot))|_{C(\mathcal{O})}^q + |\psi^{\nu+|j|+\varepsilon} D^j(u(t, \cdot) - u(s, \cdot))|_{C^\varepsilon(\mathcal{O})}^q \right) \right] \\ & \leq N \left(\|u\|_{\mathbb{H}_{p,\theta-p}^{\gamma,q}(\mathcal{O},T)}^q + \|\mathbb{D}u\|_{\mathbb{H}_{p,\theta+p}^{\gamma-2,q}(\mathcal{O},T)}^q + \|\mathbb{S}u\|_{\mathbb{H}_{p,\theta}^{\gamma-1,q}(\mathcal{O},T;\ell_2)}^q \right), \end{aligned}$$

where the constant $N \in (0, \infty)$ does not depend on u . In particular, if $u_0 = 0$, $\gamma \geq 1$, $\theta \leq d$ and $r_0 := 1 - 2/q - d/p > 0$, then for any $r \in (0, r_0)$

$$\begin{aligned} & \mathbb{E} \left[\sup_{t \leq T} \sup_{x, y \in \mathcal{O}} \frac{|u(t, x) - u(t, y)|^q}{|x - y|^{rq}} \right] < \infty, \\ & \mathbb{E} \left[\sup_{x \in \mathcal{O}} \sup_{t, s \leq T} \frac{|u(t, x) - u(s, x)|^q}{|t - s|^{rq/2}} \right] < \infty, \end{aligned}$$

see [32, Remark 4.8] for details concerning the last implication.

Combining Theorem 5.1 with the results of Section 4, we immediately obtain the following result on the Hölder-Sobolev regularity of solutions of SPDEs.

Theorem 5.3. *Let \mathcal{O} be a bounded Lipschitz domain in \mathbb{R}^d . Let $2 \leq p \leq q < \infty$, $\gamma \in \mathbb{N}_0$, $\theta \in \mathbb{R}$ and*

$$2/q < \bar{\beta} < \beta \leq 1.$$

(i) *Given the setting of Theorem 4.4, the solution $u \in \mathfrak{H}_{p,d}^{\gamma+2,q}(\mathcal{O}, T)$ of Eq. (4.12) fulfils*

$$\mathbb{E} \|\psi^{\beta-1} u\|_{C^{\bar{\beta}/2-1/q}([0,T]; H_{p,d}^{\gamma+2-\beta}(\mathcal{O}))}^q \leq N \left(\|f\|_{\mathbb{H}_{p,d}^{0,q}(\mathcal{O}, T)}^q + \|f\|_{\mathbb{H}_{p,d+p}^{\gamma,q}(\mathcal{O}, T)}^q + \|g\|_{\mathbb{H}_{p,d}^{\gamma+1,q}(\mathcal{O}, T; \ell_2)}^q \right),$$

where the constant $N \in (0, \infty)$ does not depend on u , f and g .

(ii) *Given the setting of Theorem 4.2, the solution $u \in \mathfrak{H}_{p,d}^{\gamma+2,q}(\mathcal{O}, T)$ of Eq. (4.1) fulfils*

$$\begin{aligned} \mathbb{E} \|\psi^{\beta-1} u\|_{C^{\bar{\beta}/2-1/q}([0,T]; H_{p,d}^{\gamma+2-\beta}(\mathcal{O}))}^q &\leq N \left(\|f\|_{\mathbb{H}_{p,d+p}^{\gamma,q}(\mathcal{O}, T)}^q + \|f\|_{\mathbb{H}_{p,d}^{0,q}(\mathcal{O}, T)}^q \right. \\ &\quad \left. + \|g\|_{\mathbb{H}_{p,d}^{\gamma+1,q}(\mathcal{O}, T; \ell_2)}^q + \|g\|_{\mathbb{H}_{p,d-p}^{1,q}(\mathcal{O}, T; \ell_2)}^q \right), \end{aligned}$$

where the constant $N \in (0, \infty)$ does not depend on u , f and g .

For the case that the summability parameters in time and space coincide, i.e., $q = p$, a result similar to Theorem 5.1 has been proven in [26], see Theorem 2.9 therein. The proof in [26] is straightforward and relies on [32, Corollary 4.12], which is a variant of Theorem 5.1 on the whole space \mathbb{R}^d . However, we are explicitly interested in the case $q > p$ since it allows a wider range of parameters $\bar{\beta}$ and β , and therefore leads to better regularity results. Unfortunately, the proof technique used in [26, Proposition 2.9] does not work any more in this case. Therefore, we take a different path: We use [32, Proposition 4.1], which covers the assertion of Theorem 5.1 with \mathbb{R}_+^d instead of \mathcal{O} , and the Lipschitz character of $\partial\mathcal{O}$ to derive Theorem 5.1 via a boundary flattening argument. To this end, we need the following two lemmas whose proofs are postponed to the appendix.

Lemma 5.4. *Let $G^{(1)}, G^{(2)}$ be domains in \mathbb{R}^d with non-empty boundaries, and let $\phi : G^{(1)} \rightarrow G^{(2)}$ be a bijective map, such that ϕ and ϕ^{-1} are Lipschitz continuous. Furthermore, assume that there exists a constant $N \in (0, \infty)$, such that*

$$\frac{1}{N} \rho_{G^{(1)}}(\phi^{-1}(y)) \leq \rho_{G^{(2)}}(y) \leq N \rho_{G^{(1)}}(\phi^{-1}(y)) \text{ for all } y \in G^{(2)},$$

and that the (a.e. existing) Jacobians $J\phi$ and $J\phi^{-1}$ fulfil

$$|\text{Det } J\phi| = 1 \text{ and } |\text{Det } J\phi^{-1}| = 1 \quad (\text{a.e.}).$$

Then, for any $\gamma \in [-1, 1]$, there exists a constant $N = N(d, \gamma, p, \theta, \phi) \in (0, \infty)$, which does not depend on u , such that

$$\frac{1}{N} \|u\|_{H_{p,\theta}^{\gamma}(G^{(1)})} \leq \|u \circ \phi^{-1}\|_{H_{p,\theta}^{\gamma}(G^{(2)})} \leq N \|u\|_{H_{p,\theta}^{\gamma}(G^{(1)})}$$

in the sense that, if one of the norms exists, so does the other one and the above inequality holds.

Lemma 5.5. *Let $G^{(1)}, G^{(2)}$ be bounded domains in \mathbb{R}^d and let $\phi : G^{(1)} \rightarrow G^{(2)}$ satisfy the assumptions of Lemma 5.4. Furthermore, let $u \in \mathfrak{H}_{p,\bar{\theta}}^{1,q}(G^{(1)}, T)$ with $2 \leq p \leq q < \infty$. Then $u \circ \phi^{-1} \in \mathfrak{H}_{p,\bar{\theta}}^{1,q}(G^{(2)}, T)$ with deterministic part $\mathbb{D}(u \circ \phi^{-1}) = \mathbb{D}u \circ \phi^{-1}$ and stochastic part $\mathbb{S}(u \circ \phi^{-1}) = \mathbb{S}u \circ \phi^{-1}$. In particular, for any $\varphi \in C_0^\infty(G^{(2)})$, with probability one, the equality*

$$(u(t, \cdot) \circ \phi^{-1}, \varphi) = (u(0, \cdot) \circ \phi^{-1}, \varphi) + \int_0^t ((\mathbb{D}u)(s, \cdot) \circ \phi^{-1}, \varphi) ds + \sum_{k=1}^{\infty} \int_0^t ((\mathbb{S}^k u)(s, \cdot) \circ \phi^{-1}, \varphi) dw_s^k \quad (5.3)$$

holds for all $t \in [0, T]$.

Now we are able to prove our main result in this section.

Proof of Theorem 5.1. Let us simplify notation and write $f := \mathbb{D}u$ and $g := \mathbb{S}u$ throughout the proof. We will show that (5.1) is true by induction over $\gamma \in \mathbb{N}$; estimate (5.2) can be proved analogously.

We start with the case $\gamma = 1$. Fix $x_0 \in \partial\mathcal{O}$ and choose $r > 0$ small enough, e.g., $r := r_0(10K_0)^{-1}$ with r_0 and $K_0 > 1$ from Definition 2.5. Let us assume for a moment that the supports (in the sense of distributions) of u, f and g are contained in $B_r(x_0)$ for each t and ω . With μ_0 from Definition 2.5, we introduce the function

$$\begin{aligned} \phi : G^{(1)} := \mathcal{O} \cap B_{r_0}(x_0) &\longrightarrow G^{(2)} := \phi(\mathcal{O} \cap B_{r_0}(x_0)) \subseteq \mathbb{R}_+^d \\ x = (x^1, x') &\longmapsto (x^1 - \mu_0(x'), x'), \end{aligned}$$

which fulfils all the assumptions of Lemma 5.4. Note that, since r has been chosen sufficiently small, one has $\rho_{\mathcal{O}}(x) = \rho_{G^{(1)}}(x)$ for all $x \in \mathcal{O} \cap B_r(x_0)$, so that one can easily show that the equivalence

$$\|v\|_{H_{\bar{p},\bar{\theta}}^\nu(\mathcal{O})} \sim \|v\|_{H_{\bar{p},\bar{\theta}}^\nu(G^{(1)})}, \quad v \in \mathcal{D}'(\mathcal{O}), \text{ supp } v \subseteq B_r(x_0),$$

holds for all $\nu, \bar{\theta} \in \mathbb{R}$ and $\bar{p} > 1$. Together with Lemma 5.4 we obtain for any $\nu \in [-1, 1]$,

$$\|v\|_{H_{\bar{p},\bar{\theta}}^\nu(\mathcal{O})} \sim \|v \circ \phi^{-1}\|_{H_{\bar{p},\bar{\theta}}^\nu(G^{(2)})}, \quad v \in \mathcal{D}'(\mathcal{O}), \text{ supp } v \subseteq B_r(x_0).$$

Thus, denoting $\bar{u} := u \circ \phi^{-1}$, $\bar{f} := f \circ \phi^{-1}$ and $\bar{g} := g \circ \phi^{-1}$, by Lemma 5.5 we know that on $G^{(2)}$ we have $d\bar{u} = \bar{f}dt + \bar{g}^k dw_t^k$ in the sense of distributions. Furthermore, since $\rho_{G^{(2)}}(y) = \rho_{\mathbb{R}_+^d}(y)$ for all $y \in \phi(\mathcal{O} \cap B_r(x_0))$, the equivalence

$$\|v \circ \phi^{-1}\|_{H_{\bar{p},\bar{\theta}}^\nu(G^{(2)})} \sim \|v \circ \phi^{-1}\|_{H_{\bar{p},\bar{\theta}}^\nu(\mathbb{R}_+^d)}, \quad v \in \mathcal{D}'(\mathcal{O}), \text{ supp } v \subseteq B_r(x_0),$$

holds for any $\nu \in [-1, 1]$, where we identify $v \circ \phi^{-1}$ with its extension to \mathbb{R}_+^d by zero. Therefore, by making slight abuse of notation and writing \bar{u}, \bar{f} and \bar{g} for the extension by zero on \mathbb{R}_+^d of \bar{u}, \bar{f} and \bar{g} respectively, we have

$$\bar{u} \in \mathbb{H}_{p,\bar{\theta}-p}^{1,q}(\mathbb{R}_+^d, T), \quad \bar{u}(0) \in U_{p,\bar{\theta}}^{1,q}(\mathbb{R}_+^d), \quad \bar{f} \in \mathbb{H}_{p,\bar{\theta}+p}^{-1,q}(\mathbb{R}_+^d, T), \quad \bar{g} \in \mathbb{H}_{p,\bar{\theta}}^{0,q}(\mathbb{R}_+^d, T; \ell_2),$$

and $d\bar{u} = \bar{f}dt + \bar{g}^k dw_t^k$ is fulfilled on \mathbb{R}_+^d in the sense of distributions. Thus, we can apply [32, Theorem 4.1] and use the equivalences above to get estimate (5.1) in the following way:

$$\begin{aligned} \mathbb{E}[u]_{C^{\bar{\beta}/2-1/q}([0,T];H_{p,\theta+p(\beta-1)}^{1-\beta}(\mathcal{O}))}^q &\leq N \mathbb{E}[\bar{u}]_{C^{\bar{\beta}/2-1/q}([0,T];H_{p,\theta+p(\beta-1)}^{1-\beta}(\mathbb{R}_+^d))}^q \\ &\leq NT^{(\beta-\bar{\beta})q/2} \left(\|\bar{u}\|_{\mathbb{H}_{p,\theta-p}^{1,q}(\mathbb{R}_+^d,T)}^q + \|\bar{f}\|_{\mathbb{H}_{p,\theta+p}^{-1,q}(\mathbb{R}_+^d,T)}^q + \|\bar{g}\|_{\mathbb{H}_{p,\theta}^{0,q}(\mathbb{R}_+^d,T;\ell_2)}^q \right) \\ &\leq NT^{(\beta-\bar{\beta})q/2} \left(\|u\|_{\mathbb{H}_{p,\theta-p}^{1,q}(\mathcal{O},T)}^q + \|f\|_{\mathbb{H}_{p,\theta+p}^{-1,q}(\mathcal{O},T)}^q + \|g\|_{\mathbb{H}_{p,\theta}^{0,q}(\mathcal{O},T;\ell_2)}^q \right). \end{aligned}$$

Now let us give up the assumption on the supports of u , f and g . Let $\xi_0, \xi_1, \dots, \xi_m$, be a partition of unity of \mathcal{O} , such that $\xi_0 \in C_0^\infty(\mathcal{O})$, and, for $i = 1, \dots, m$, $\xi_i \in C_0^\infty(B_r(x_i))$ with $x_i \in \partial\mathcal{O}$. Obviously, $d(\xi_i u) = \xi_i f dt + \xi_i g_t^k dw_t^k$ for $i = 0, \dots, m$. Since

$$\mathbb{E}[\psi^{\beta-1}u]_{C^{\bar{\beta}/2-1/q}([0,T];H_{p,\theta}^{1-\beta}(\mathcal{O}))}^q \leq N(m, q) \sum_{i=0}^m \mathbb{E}[\psi^{\beta-1}(\xi_i u)]_{C^{\bar{\beta}/2-1/q}([0,T];H_{p,\theta}^{1-\beta}(\mathcal{O}))}^q,$$

we just have to estimate $\mathbb{E}[\psi^{\beta-1}\xi_i u]_{C^{\bar{\beta}/2-1/q}([0,T];H_{p,\theta}^{1-\beta}(\mathcal{O}))}^q$ for each $i \in \{0, \dots, m\}$. For $i \geq 1$ one gets the required estimate as before, using the fact that $C_0^\infty(\mathcal{O})$ -functions are pointwise multipliers in all spaces $H_{\bar{p},\bar{\theta}}^\nu(\mathcal{O})$, $\nu, \bar{\theta} \in \mathbb{R}$, $\bar{p} > 1$, see, e.g., [38, Theorem 3.1]. The case $i = 0$ can be treated as follows: Since ξ_0 has compact support in \mathcal{O} , for all $\nu, \bar{\theta} \in \mathbb{R}$ and $\bar{p} > 1$, we have

$$\|v\xi_0\|_{H_{\bar{p},\bar{\theta}}^\nu(\mathcal{O})} \sim \|v\xi_0\|_{H_{\bar{p}}^\nu(\mathbb{R}^d)}, \quad v \in \mathcal{D}'(\mathcal{O}), \quad (5.4)$$

and consequently

$$\mathbb{E}[\psi^{\beta-1}\xi_0 u]_{C^{\bar{\beta}/2-1/q}([0,T];H_{p,\theta}^{1-\beta}(\mathcal{O}))}^q \sim \mathbb{E}[\xi_0 u]_{C^{\bar{\beta}/2-1/q}([0,T];H_p^{1-\beta}(\mathbb{R}^d))}^q.$$

By [32, Theorem 4.11], a further application of (5.4) and the fact that $C_0^\infty(\mathcal{O})$ -functions are pointwise multipliers in all spaces $H_{\bar{p},\bar{\theta}}^\nu(\mathcal{O})$, we obtain

$$\begin{aligned} &\mathbb{E}[\xi_0 u]_{C^{\bar{\beta}/2-1/q}([0,T];H_p^{1-\beta}(\mathbb{R}^d))}^q \\ &\leq NT^{(\beta-\bar{\beta})q/2} \left(\|\xi_0 u\|_{\mathbb{H}_p^{1,q}(\mathbb{R}^d,T)}^q + \|\xi_0 f\|_{\mathbb{H}_p^{-1,q}(\mathbb{R}^d,T)}^q + \|\xi_0 g\|_{\mathbb{H}_p^{0,q}(\mathbb{R}^d,T;\ell_2)}^q \right) \\ &\leq NT^{(\beta-\bar{\beta})q/2} \left(\|\psi^{-1}\xi_0 u\|_{\mathbb{H}_{p,\theta}^{1,q}(\mathcal{O},T)}^q + \|\psi\xi_0 f\|_{\mathbb{H}_{p,\theta}^{-1,q}(\mathcal{O},T)}^q + \|\xi_0 g\|_{\mathbb{H}_{p,\theta}^{0,q}(\mathcal{O},T;\ell_2)}^q \right) \\ &\leq NT^{(\beta-\bar{\beta})q/2} \left(\|\psi^{-1}u\|_{\mathbb{H}_{p,\theta}^{1,q}(\mathcal{O},T)}^q + \|\psi f\|_{\mathbb{H}_{p,\theta}^{-1,q}(\mathcal{O},T)}^q + \|g\|_{\mathbb{H}_{p,\theta}^{0,q}(\mathcal{O},T;\ell_2)}^q \right). \end{aligned}$$

This finishes the proof of estimate (5.1) for the case $\gamma = 1$.

Next, let us move to the inductive step and assume that the assertion is true for some $\gamma = n \in \mathbb{N}$. Fix $u \in \mathfrak{H}_{p,\theta}^{n+1,q}(\mathcal{O}, T)$. Then $v := \psi u_x \in \mathfrak{H}_{p,\theta}^{n,q}(\mathcal{O}, T)$ and $dv = \psi f_x dt + \psi g_x^k dw_t^k$ (component-wise). Also, by Lemma 2.1 (iii) and (iv),

$$\begin{aligned} \mathbb{E}[\psi^{\beta-1}u]_{C^{\bar{\beta}/2-1/q}([0,T];H_{p,\theta}^{n+1-\beta}(\mathcal{O}))}^q &\leq N \left(\mathbb{E}[\psi^{\beta-1}u]_{C^{\bar{\beta}/2-1/q}([0,T];H_{p,\theta}^{n-\beta}(\mathcal{O}))}^q + \right. \\ &\quad \left. \mathbb{E}[\psi^{\beta-1}v]_{C^{\bar{\beta}/2-1/q}([0,T];H_{p,\theta}^{n-\beta}(\mathcal{O}))}^q \right). \end{aligned}$$

Using the induction hypothesis and applying Lemma 2.1(iii) and (iv) once more, we see that the induction goes through. \square

6 Besov spaces and their relationship to weighted Sobolev spaces

We turn our attention to the scale of Besov spaces

$$B_{\tau,\tau}^\alpha(\mathcal{O}), \quad \frac{1}{\tau} = \frac{\alpha}{d} + \frac{1}{p}, \quad \alpha > 0, \quad (*)$$

where $p \geq 2$ is fixed and $\mathcal{O} \subseteq \mathbb{R}^d$ is a bounded Lipschitz domain. As pointed out in the introduction, our motivation for considering this scale is its close connection to nonlinear approximation theory. The main result of this section, Theorem 6.9, is a general embedding of the weighted Sobolev spaces

$$H_{p,d-\nu p}^\gamma(\mathcal{O}), \quad \gamma, \nu > 0, \quad (6.1)$$

into Besov spaces of the scale (*). In the previous sections we have seen that the stochastic parabolic weighted Sobolev spaces $\mathfrak{H}_{p,\theta}^{\gamma,q}(\mathcal{O}, T)$ are suitable for the analysis of SPDEs of the type (3.1). The elements of these spaces are stochastic processes with values in the spaces $H_{p,\theta-p}^\gamma(\mathcal{O}) = H_{p,d-\nu p}^\gamma(\mathcal{O})$ with $\nu = 1 + (d - \theta)/p$. Thus, by proving the embedding mentioned above, we show that the regularity analysis for SPDEs of type (3.1) in terms of the scale (*) can be traced back to the analysis of such equations in terms of the spaces $\mathfrak{H}_{p,\theta}^{\gamma,q}(\mathcal{O}, T)$.

The outline of this section is as follows. In Subsection 6.1 we give a definition of Besov spaces and describe their characterization in terms of wavelets. Based on this characterization and some auxiliary results, we investigate the relation of the scales (6.1) and (*) in Subsection 6.2. Here we will also see that, for the relevant range of parameters γ and ν , the spaces $H_{p,d-\nu p}^\gamma(\mathcal{O})$ act like Besov spaces $B_{p,p}^{\gamma \wedge \nu}(\mathcal{O})$ with zero trace on the boundary (Remark 6.7).

6.1 Besov spaces: Definition and wavelet decomposition.

Our standard reference concerning Besov spaces and wavelets is the monograph [5]. Throughout this subsection, let $G \subseteq \mathbb{R}^d$ be an arbitrary domain.

For a function $f : G \rightarrow \mathbb{R}$ and a natural number $n \in \mathbb{N}$ let

$$\Delta_h^n f(x) := \prod_{i=0}^n \mathbb{1}_G(x + ih) \cdot \sum_{j=0}^n \binom{n}{j} (-1)^{n-j} f(x + jh)$$

be the n -th difference of f with step $h \in \mathbb{R}^d$. For $p \in (0, \infty)$, the n -th order L_p -modulus of smoothness of f is given by

$$\omega^n(t, f, G)_p := \omega^n(t, f)_p := \sup_{|h| < t} \|\Delta_h^n f\|_{L_p(G)}, \quad t > 0.$$

One definition of Besov spaces that fits in our purpose is the following:

Definition 6.1. *Let $s, p, q \in (0, \infty)$ and $n \in \mathbb{N}$ with $n > s$. Then $B_{p,q}^s(G)$ is the collection of all functions $f \in L_p(G)$ such that*

$$|f|_{B_{p,q}^s(G)} := \left(\int_0^\infty \left[t^{-s} \omega^n(t, f)_p \right]^q \frac{dt}{t} \right)^{1/q} < \infty.$$

These classes are equipped with a (quasi-)norm by taking

$$\|f\|_{B_{p,q}^s(G)} := \|f\|_{L_p(G)} + |f|_{B_{p,q}^s(G)}.$$

Remark 6.2. For a more general definition of Besov spaces, including the cases where $p, q = \infty$ and $s < 0$ see, e.g., [44].

We want to describe $B_{p,q}^s(\mathbb{R}^d)$ by means of wavelet expansions. To this end let φ be a scaling function of tensor product type on \mathbb{R}^d and let ψ_i , $i = 1, \dots, 2^d - 1$, be corresponding multivariate mother wavelets such that, for a given $r \in \mathbb{N}$ and some $M > 0$, the following locality, smoothness and vanishing moment conditions hold: for all $i = 1, \dots, 2^d - 1$,

$$\text{supp } \varphi, \text{supp } \psi_i \subset [-M, M]^d, \quad (6.2)$$

$$\varphi, \psi_i \in C^r(\mathbb{R}^d), \quad (6.3)$$

$$\int x^\alpha \psi_i(x) dx = 0 \quad \text{for all } \alpha \in \mathbb{N}^d \text{ with } |\alpha| \leq r. \quad (6.4)$$

We assume that

$$\{\varphi_k, \psi_{i,j,k} : (i, j, k) \in \{1, \dots, 2^d - 1\} \times \mathbb{N}_0 \times \mathbb{Z}^d\}$$

is a Riesz basis of $L_2(\mathbb{R}^d)$, where we used the abbreviations for dyadic shifts and dilations of the scaling function and the corresponding wavelets

$$\varphi_k(x) := \varphi(x - k), \quad x \in \mathbb{R}^d, \quad \text{for } k \in \mathbb{Z}^d, \text{ and} \quad (6.5)$$

$$\psi_{i,j,k}(x) := 2^{jd/2} \psi_i(2^j x - k), \quad x \in \mathbb{R}^d, \quad \text{for } (i, j, k) \in \{1, \dots, 2^d - 1\} \times \mathbb{N} \times \mathbb{Z}^d. \quad (6.6)$$

Further, we assume that there exists a dual Riesz basis satisfying the same requirements. More precisely, there exist functions $\tilde{\varphi}$ and $\tilde{\psi}_i$, $i = 1, \dots, 2^d - 1$, such that conditions (6.2), (6.3) and (6.4) hold if φ and ψ are replaced by $\tilde{\varphi}$ and $\tilde{\psi}_i$, and such that the biorthogonality relations

$$\langle \tilde{\varphi}_k, \psi_{i,j,k} \rangle = \langle \tilde{\psi}_{i,j,k}, \varphi_k \rangle = 0, \quad \langle \tilde{\varphi}_k, \varphi_l \rangle = \delta_{k,l}, \quad \langle \tilde{\psi}_{i,j,k}, \psi_{u,v,l} \rangle = \delta_{i,u} \delta_{j,v} \delta_{k,l},$$

are fulfilled. Here we use analogous abbreviations to (6.5) and (6.6) for the dyadic shifts and dilations of $\tilde{\varphi}$ and $\tilde{\psi}_i$, and $\delta_{k,l}$ denotes the Kronecker symbol. We refer to [5, Chapter 2] for the construction of biorthogonal wavelet bases, see also [12] and [6]. To keep notation simple, we will write

$$\psi_{i,j,k,p} := 2^{jd(1/p-1/2)} \psi_{i,j,k} \quad \text{and} \quad \tilde{\psi}_{i,j,k,p'} := 2^{jd(1/p'-1/2)} \tilde{\psi}_{i,j,k},$$

for the L_p -normalized wavelets and the correspondingly modified duals, with $p' := p/(p-1)$ if $p \in (0, \infty)$, $p \neq 1$, and $p' := \infty$, $1/p' := 0$ if $p = 1$.

The following theorem shows how Besov spaces can be described by decay properties of the wavelet coefficients, if the parameters fulfil certain conditions.

Theorem 6.3. *Let $p, q \in (0, \infty)$ and $s > \max\{0, d(1/p - 1)\}$. Choose $r \in \mathbb{N}$ such that $r > s$ and construct a biorthogonal wavelet Riesz basis as described above. Then a locally integrable function $f : \mathbb{R}^d \rightarrow \mathbb{R}$ is in the Besov space $B_{p,q}^s(\mathbb{R}^d)$ if, and only if,*

$$f = \sum_{k \in \mathbb{Z}^d} \langle f, \tilde{\varphi}_k \rangle \varphi_k + \sum_{i=1}^{2^d-1} \sum_{j \in \mathbb{N}} \sum_{k \in \mathbb{Z}^d} \langle f, \tilde{\psi}_{i,j,k,p'} \rangle \psi_{i,j,k,p} \quad (6.7)$$

(convergence in $\mathcal{D}'(\mathbb{R}^d)$) with

$$\left(\sum_{k \in \mathbb{Z}^d} |\langle f, \tilde{\varphi}_k \rangle|^p \right)^{1/p} + \left(\sum_{i=1}^{2^d-1} \sum_{j \in \mathbb{N}_0} 2^{jsq} \left(\sum_{k \in \mathbb{Z}^d} |\langle f, \tilde{\psi}_{i,j,k,p'} \rangle|^p \right)^{1/q} \right)^{1/q} < \infty, \quad (6.8)$$

and (6.8) is an equivalent (quasi-)norm for $B_{p,q}^s(\mathbb{R}^d)$.

Remark 6.4. A proof of this theorem for the case $p \geq 1$ can be found in [40, §10 of Chapter 6]. For the general case see for example [36] or [5, Theorem 3.7.7]. Of course, if (6.8) holds then the infinite sum in (6.7) converges also in $B_{p,q}^s(\mathbb{R}^d)$. If $s > \max\{0, d(1/p - 1)\}$ we have the embedding $B_{p,q}^s(\mathbb{R}^d) \subset L_{\bar{s}}(\mathbb{R}^d)$ for some $\bar{s} > 1$, see, e.g. [5, Corollary 3.7.1].

A simple computation gives us the following characterization of Besov spaces from the scale (*) on \mathbb{R}^d .

Corollary 6.5. Let $p \in (1, \infty)$, $\alpha > 0$ and $\tau \in \mathbb{R}$ such that $1/\tau = \alpha/d + 1/p$. Choose $r \in \mathbb{N}$ such that $r > \alpha$ and construct a biorthogonal wavelet Riesz basis as described above. Then a locally integrable function $f : \mathbb{R}^d \rightarrow \mathbb{R}$ is in the Besov space $B_{\tau,\tau}^\alpha(\mathbb{R}^d)$ if and only if

$$f = \sum_{k \in \mathbb{Z}^d} \langle f, \tilde{\varphi}_k \rangle \varphi_k + \sum_{i=1}^{2^d-1} \sum_{j \in \mathbb{N}_0} \sum_{k \in \mathbb{Z}^d} \langle f, \tilde{\psi}_{i,j,k,p'} \rangle \psi_{i,j,k,p} \quad (6.9)$$

(convergence in $\mathcal{D}'(\mathbb{R}^d)$) with

$$\left(\sum_{k \in \mathbb{Z}^d} |\langle f, \tilde{\varphi}_k \rangle|^\tau \right)^{1/\tau} + \left(\sum_{i=1}^{2^d-1} \sum_{j \in \mathbb{N}_0} \sum_{k \in \mathbb{Z}^d} |\langle f, \tilde{\psi}_{i,j,k,p'} \rangle|^\tau \right)^{1/\tau} < \infty, \quad (6.10)$$

and (6.10) is an equivalent (quasi-)norm for $B_{\tau,\tau}^\alpha(\mathbb{R}^d)$.

6.2 From weighted Sobolev spaces to Besov spaces

In this subsection we will prove two embeddings of weighted Sobolev spaces into Besov spaces. We first focus on the case where the integrability parameter $p \in [2, \infty)$ of the weighted Sobolev spaces and the Besov spaces under consideration coincide, see Lemma 6.6. This will pave the way for proving a general embedding of weighted Sobolev spaces into the Besov spaces from the scale (*) in Theorem 6.9. Remember that in this article $\mathcal{O} \subseteq \mathbb{R}^d$ always denotes a bounded Lipschitz domain.

Lemma 6.6. Let \mathcal{O} be a bounded Lipschitz domain in \mathbb{R}^d . Let $\gamma, \nu \in (0, \infty)$ and $p \in [2, \infty)$. Then the following embedding holds:

$$H_{p,d-\nu p}^\gamma(\mathcal{O}) \hookrightarrow B_{p,p}^{\gamma \wedge \nu}(\mathcal{O}). \quad (6.11)$$

Proof. We start the proof by considering the case where $\gamma = \nu$, i.e., we prove that for $\gamma > 0$ and $p \geq 2$ we have

$$H_{p,d-\gamma p}^\gamma(\mathcal{O}) \hookrightarrow B_{p,p}^\gamma(\mathcal{O}). \quad (6.12)$$

It is well-known, see [35, Theorem 9.7], that for $k \in \mathbb{N}_0$,

$$H_{p,d-kp}^k(\mathcal{O}) = \mathring{W}_p^k(\mathcal{O}), \quad (6.13)$$

where $\mathring{W}_p^k(\mathcal{O})$ denotes the completion of $C_0^\infty(\mathcal{O})$ in the classical $L_p(\mathcal{O})$ -Sobolev space $W_p^k(\mathcal{O})$. We use the convention $W_p^0(\mathcal{O}) := L_p(\mathcal{O})$. Since $p \geq 2$ we have $W_p^k(\mathcal{O}) \hookrightarrow B_{p,p}^k(\mathcal{O})$, see, e.g., [43, Remark 2.3.3/4 and Theorem 4.6.1(b)] together with [15]. We therefore obtain (6.12) for $k \in \mathbb{N}_0$. In the case of fractional $\gamma \in (0, \infty) \setminus \mathbb{N}$ we argue as follows. Let $\gamma = k + \eta$ with $k \in \mathbb{N}_0$ and $\eta \in (0, 1)$. By [38, Proposition 2.4],

$$H_{p,d-(k+\eta)p}^{k+\eta}(\mathcal{O}) = [H_{p,d-kp}^k(\mathcal{O}), H_{p,d-(k+1)p}^{k+1}(\mathcal{O})]_\eta.$$

Using (6.13) we have

$$H_{p,d-(k+\eta)p}^{k+\eta}(\mathcal{O}) = [\mathring{W}_p^k(\mathcal{O}), \mathring{W}_p^{k+1}(\mathcal{O})]_\eta \hookrightarrow [W_p^k(\mathcal{O}), W_p^{k+1}(\mathcal{O})]_\eta.$$

For any $k \in \mathbb{N}$, it is well-known that the Sobolev space $W_p^k(\mathcal{O})$ coincides with the Triebel-Lizorkin spaces $F_{p,2}^k(\mathcal{O})$, see, e.g., [44, Theorem 1.122]. Moreover, we have $L_p(\mathcal{O}) \hookrightarrow F_{p,2}^0(\mathcal{O})$, see [44, (1.2) together with Definition 1.95]. Thus,

$$H_{p,d-(k+\eta)p}^{k+\eta}(\mathcal{O}) \hookrightarrow [F_{p,2}^k(\mathcal{O}), F_{p,2}^{k+1}(\mathcal{O})]_\eta.$$

The fact that Triebel-Lizorkin spaces constitute a scale of complex interpolation spaces, see, e.g., [44, Corollary 1.111], leads to

$$H_{p,d-(k+\eta)p}^{k+\eta}(\mathcal{O}) \hookrightarrow F_{p,2}^{k+\eta}(\mathcal{O}).$$

For $p \geq 2$, it is well known that $F_{p,2}^s(\mathcal{O}) \hookrightarrow B_{p,p}^s(\mathcal{O})$ for any $s \in \mathbb{R}$, see, e.g., (1.299) in [44] together with [15]. Therefore,

$$H_{p,d-(k+\eta)p}^{k+\eta}(\mathcal{O}) \hookrightarrow B_{p,p}^{k+\eta}(\mathcal{O}),$$

and (6.12) is proved for general $\gamma > 0$. The embedding (6.11) for $\gamma \neq \nu$ follows now by using standard arguments. Indeed, since $\gamma \geq \gamma \wedge \nu$ we have

$$H_{p,d-\nu p}^\gamma(\mathcal{O}) \hookrightarrow H_{p,d-\nu p}^{\gamma \wedge \nu}(\mathcal{O}),$$

see [38], the line after Definition 2.1. Furthermore, $d - \nu p \leq d - (\gamma \wedge \nu)p$ implies

$$H_{p,d-\nu p}^{\gamma \wedge \nu}(\mathcal{O}) \hookrightarrow H_{p,d-(\gamma \wedge \nu)p}^{\gamma \wedge \nu}(\mathcal{O}),$$

see [38, Corollary 4.2]. A combination of these two embeddings with (6.12) finally gives (6.11). \square

Remark 6.7. *Since $\mathcal{O} \subseteq \mathbb{R}^d$ is assumed to be a bounded Lipschitz domain, we know by [21, Chapter VIII, Theorem 2] that for $1/p < s$ the operator Tr , initially defined to $C^\infty(\overline{\mathcal{O}})$ as the restriction on $\partial\mathcal{O}$, extends to a bounded linear operator from $B_{p,p}^s(\mathcal{O})$ to $B_{p,p}^{s-1/p}(\partial\mathcal{O})$. In this case we denote by $B_{p,p,0}^s(\mathcal{O})$ the subspace of $B_{p,p}^s(\mathcal{O})$ with zero boundary trace, i.e.,*

$$B_{p,p,0}^s(\mathcal{O}) := \{u \in B_{p,p}^s(\mathcal{O}) : \text{Tr } u = 0\}, \quad \frac{1}{p} < s < 1 + \frac{1}{p}.$$

By [19, Theorem 3.12] these spaces coincide with the closure of $C_0^\infty(\mathcal{O})$ in $B_{p,p}^s(\mathcal{O})$, i.e.,

$$\mathring{B}_{p,p}^s(\mathcal{O}) := \overline{C_0^\infty(\mathcal{O})}^{\|\cdot\|_{B_{p,p}^s(\mathcal{O})}} = B_{p,p,0}^s(\mathcal{O}) \quad \text{for} \quad \frac{1}{p} < s < 1 + \frac{1}{p}.$$

Thus, if $1/p < \gamma \wedge \nu < 1 + 1/p$, Lemma 6.6 together with Lemma 2.1(i), lead to

$$H_{p,d-\nu p}^\gamma(\mathcal{O}) \hookrightarrow \mathring{B}_{p,p}^{\gamma \wedge \nu}(\mathcal{O}) = B_{p,p,0}^{\gamma \wedge \nu}(\mathcal{O}) = \{u \in B_{p,p}^{\gamma \wedge \nu}(\mathcal{O}) : \text{Tr } u = 0\}.$$

In Section 4 we considered SPDEs in the setting of [26]. The solutions to these equations are stochastic processes taking values in $H_{p,d-\nu p}^\gamma(\mathcal{O})$ with $\nu := 1 + \frac{d-\theta}{p}$, where the value of θ never leaves the range $d-1 < \theta < d+p-1$, compare also [34]. This condition is equivalent to $1/p < \nu < 1 + 1/p$ with ν as introduced before. Hence, if $\gamma > 1/p$ we deal with solutions with zero boundary condition, in the sense that the well defined linear and continuous boundary trace Tr equals zero.

In the second part of this subsection we investigate the relationship between weighted Sobolev spaces and the Besov spaces from the scale $(*)$. In [10], the scale $(*)$ is used to analyse the regularity of harmonic functions on a bounded Lipschitz domain $\mathcal{O} \subseteq \mathbb{R}^d$. Denoting by $\Theta(\mathcal{O})$ the set of harmonic functions on \mathcal{O} we can formulate the main result therein, [10, Theorem 3.2], as follows:

$$\Theta(\mathcal{O}) \cap B_{p,p}^\nu(\mathcal{O}) \hookrightarrow B_{\tau,\tau}^\alpha(\mathcal{O}), \quad \frac{1}{\tau} = \frac{\alpha}{d} + \frac{1}{p}, \quad \text{for all } 0 < \alpha < \nu \frac{d}{d-1} = \sup_{m \in \mathbb{N}} \min \left\{ m, \nu \frac{d}{d-1} \right\}.$$

One of the main ingredients for the proof of this statement is the fact that harmonic functions contained in $B_{p,p}^\nu(\mathcal{O})$ have finite weighted Sobolev half-norm

$$\|u\|_{H_{p,d-\nu p}^m(\mathcal{O})} := \left(\sum_{\substack{\alpha \in \mathbb{N}_0^d \\ |\alpha|=m}} \int_{\mathcal{O}} |\rho(x)^{|\alpha|} D^\alpha u(x)|^p \rho(x)^{-\nu p} dx \right)^{1/p}$$

for any $m \in \mathbb{N}$, see [10, Theorem 3.1] for details. It turns out that arguing along the lines of [10, Theorem 3.2] one can even show that for $\nu > 0$ and $m \in \mathbb{N}$,

$$H_{p,d-\nu p}^m(\mathcal{O}) \cap B_{p,p}^\nu(\mathcal{O}) \hookrightarrow B_{\tau,\tau}^\alpha(\mathcal{O}), \quad \frac{1}{\tau} = \frac{\alpha}{d} + \frac{1}{p}, \quad \text{for all } 0 < \alpha < \min \left\{ m, \nu \frac{d}{d-1} \right\}.$$

Combining this with Lemma 6.6, we obtain

$$H_{p,d-\nu p}^m(\mathcal{O}) \hookrightarrow B_{\tau,\tau}^\alpha(\mathcal{O}), \quad \frac{1}{\tau} = \frac{\alpha}{d} + \frac{1}{p}, \quad \text{for all } 0 < \alpha < \min \left\{ m, \nu \frac{d}{d-1} \right\}. \quad (6.14)$$

In what follows we give a detailed proof of the extension of (6.14) to arbitrary smoothness parameters $\gamma > 0$ instead of $m \in \mathbb{N}$. To this end, let us fix some notations. We will use a wavelet Riesz-basis

$$\{\varphi_k, \psi_{i,j,k} : (i,j,k) \in \{1, \dots, 2^d - 1\} \times \mathbb{N}_0 \times \mathbb{Z}^d\}$$

of $L_2(\mathbb{R}^d)$ which satisfies the assumptions from Section 6.1 with $\mathcal{O} \subseteq [-M, M]^d$ and r large enough— we will always clarify what we mean by that in the particular theorems. Given $(j,k) \in \mathbb{N}_0 \times \mathbb{Z}^d$, let

$$Q_{j,k} := 2^{-j}k + 2^{-j}[-M, M]^d,$$

so that $\text{supp } \psi_{i,j,k} \subset Q_{j,k}$ for all $i \in \{1, \dots, 2^d - 1\}$ and $\text{supp } \varphi_k \subset Q_{0,k}$ for all $k \in \mathbb{Z}^d$. Remember that the supports of the corresponding dual basis fulfil the same requirements. For our purpose the set of all indices associated with those wavelets that may have common support with the domain \mathcal{O} will play an important role and we denote it by

$$\Lambda := \{(i, j, k) \in \{1, \dots, 2^d - 1\} \times \mathbb{N}_0 \times \mathbb{Z}^d \mid Q_{j,k} \cap \mathcal{O} \neq \emptyset\}.$$

Furthermore, we want to distinguish the indices corresponding to wavelets with support in the interior of the domain from the ones corresponding to wavelets which might have support on the boundary of \mathcal{O} . To this end we write

$$\begin{aligned} \rho_{j,k} &:= \text{dist}(Q_{j,k}, \partial\mathcal{O}) = \inf_{x \in Q_{j,k}} \rho(x), \\ \Lambda_j &:= \{(i, l, k) \in \Lambda : l = j\}, \\ \Lambda_{j,m} &:= \{(i, j, k) \in \Lambda_j : m2^{-j} \leq \rho_{j,k} < (m+1)2^{-j}\}, \\ \Lambda_j^0 &:= \Lambda_j \setminus \Lambda_{j,0}, \\ \Lambda^0 &:= \bigcup_{j \in \mathbb{N}_0} \Lambda_j^0, \end{aligned}$$

where $j, m \in \mathbb{N}_0$ and $k \in \mathbb{Z}^d$. Later we will also use the notation

$$\Gamma := \{k \in \mathbb{Z}^d : Q_{0,k} \cap \mathcal{O} \neq \emptyset\}.$$

The following lemma paves the way for proving (6.14) for arbitrary $\gamma > 0$ instead of just $\gamma = m \in \mathbb{N}$. It establishes an estimate for, roughly speaking, a discretization of a weighted Sobolev norm in terms of the supports of the wavelets in the interior of \mathcal{O} at a fixed scaling level $j \in \mathbb{N}_0$. For better readability, we place the quite technical proof in the appendix. Remember that we write A° for the interior of an arbitrary subset A of \mathbb{R}^d .

Lemma 6.8. *Let \mathcal{O} be a bounded Lipschitz domain in \mathbb{R}^d . Let $p \in [2, \infty)$, $\gamma \in (0, \infty)$ and $\nu \in \mathbb{R}$ with $\gamma \geq \nu$. Furthermore, assume $u \in H_{p,d-\nu p}^\gamma(\mathcal{O})$. Then, for all $j \in \mathbb{N}_0$, the inequality*

$$\sum_{(i,j,k) \in \Lambda_j^0} (\rho_{j,k}^{\gamma-\nu} |u|_{B_{p,p}^\gamma(Q_{j,k}^\circ)})^p \leq N \|u\|_{H_{p,d-\nu p}^\gamma(\mathcal{O})}^p$$

holds, with a constant $N \in (0, \infty)$ which does not depend on j and u .

Now we can prove the main result of this section. We use the convention ‘ $1/0 := \infty$ ’.

Theorem 6.9. *Let \mathcal{O} be a bounded Lipschitz domain in \mathbb{R}^d . Let $p \in [2, \infty)$, and $\gamma, \nu \in (0, \infty)$. Then*

$$H_{p,d-\nu p}^\gamma(\mathcal{O}) \hookrightarrow B_{\tau,\tau}^\alpha(\mathcal{O}), \quad \frac{1}{\tau} = \frac{\alpha}{d} + \frac{1}{p}, \quad \text{for all } 0 < \alpha < \min \left\{ \gamma, \nu \frac{d}{d-1} \right\}.$$

Proof. Let us start with the case $\nu > \gamma$. Then, for any $0 < \alpha < \gamma$, we have

$$H_{p,d-\nu p}^\gamma(\mathcal{O}) \hookrightarrow B_{p,p}^\gamma(\mathcal{O}) \hookrightarrow B_{\tau,\tau}^\alpha(\mathcal{O}), \quad \frac{1}{\tau} = \frac{\alpha}{d} + \frac{1}{p},$$

where we used Lemma 6.6 and standard embeddings for Besov spaces. Therefore, in this case the assertion of the theorem follows immediately. From now on, let us assume that $0 < \nu \leq \gamma$. We fix α and τ as stated in the theorem and choose the wavelet Riesz-basis of $L_2(\mathbb{R})$ from above with $r > \gamma$. We also fix $u \in H_{p,d-\nu p}^\gamma(\mathcal{O})$. Due to Lemma 6.6 we have $u \in B_{p,p}^\nu(\mathcal{O})$. As \mathcal{O} is a Lipschitz domain there exists a linear and bounded extension operator $\mathcal{E} : B_{p,p}^\nu(\mathcal{O}) \rightarrow B_{p,p}^\nu(\mathbb{R}^d)$, i.e., there exists a constant $N > 0$ such that:

$$\mathcal{E}u|_{\mathcal{O}} = u \quad \text{and} \quad \|\mathcal{E}u\|_{B_{p,p}^\nu(\mathbb{R}^d)} \leq N\|u\|_{B_{p,p}^\nu(\mathcal{O})},$$

see, e.g., [42]. In the sequel we will omit the \mathcal{E} in our notation and write u instead of $\mathcal{E}u$. Theorem 6.3 tells us that the following equality holds on the domain \mathcal{O} :

$$u = \sum_{k \in \Gamma} \langle u, \tilde{\varphi}_k \rangle \varphi_k + \sum_{(i,j,k) \in \Lambda} \langle u, \tilde{\psi}_{i,j,k,p'} \rangle \psi_{i,j,k,p},$$

where the sums converge unconditionally in $B_{p,p}^\nu(\mathbb{R}^d)$. Furthermore, cf. Corollary 6.5, we have

$$\|u\|_{B_{p,p}^{\alpha,\tau}(\mathcal{O})}^\tau \leq N \left(\sum_{k \in \Gamma} |\langle u, \tilde{\varphi}_k \rangle|^\tau + \sum_{(i,j,k) \in \Lambda} |\langle u, \tilde{\psi}_{i,j,k,p'} \rangle|^\tau \right),$$

see also [15]. Hence, by Lemma 6.6, it is enough to prove that

$$\sum_{k \in \Gamma} |\langle u, \tilde{\varphi}_k \rangle|^\tau \leq N \|u\|_{B_{p,p}^\nu(\mathcal{O})}^\tau \quad (6.15)$$

and

$$\sum_{(i,j,k) \in \Lambda} |\langle u, \tilde{\psi}_{i,j,k,p'} \rangle|^\tau \leq N \left(\|u\|_{H_{p,d-\nu p}^\gamma(\mathcal{O})} + \|u\|_{B_{p,p}^\nu(\mathcal{O})} \right)^\tau. \quad (6.16)$$

We start with (6.15). The index set Γ introduced above is finite because of the boundedness of \mathcal{O} , so that we can use Jensen's inequality to obtain

$$\sum_{k \in \Gamma} |\langle u, \tilde{\varphi}_k \rangle|^\tau \leq N \left(\left(\sum_{k \in \Gamma} |\langle u, \tilde{\varphi}_k \rangle|^p \right)^{1/p} \right)^\tau \leq N \|u\|_{B_{p,p}^\nu(\mathcal{O})}^\tau.$$

In the last step we have used Theorem 6.3 and the boundedness of the extension operator.

Now let us focus on inequality (6.16). To this end, we use the notations from above and split the expression on the left hand side of (6.16) into

$$\sum_{(i,j,k) \in \Lambda^0} |\langle u, \tilde{\psi}_{i,j,k,p'} \rangle|^\tau + \sum_{(i,j,k) \in \Lambda \setminus \Lambda^0} |\langle u, \tilde{\psi}_{i,j,k,p'} \rangle|^\tau =: I + \mathbf{II} \quad (6.17)$$

and estimate each term separately.

Let us begin with I . Fix $(i, j, k) \in \Lambda^0$. As a consequence of Lemma 6.8, we know that $u|_{Q_{j,k}^\circ} \in B_{p,p}^\gamma(Q_{j,k}^\circ)$. By a Whitney-type inequality, also known as the Deny-Lions lemma, see, e.g., [14, Theorem 3.5], there exists a polynomial $P_{j,k}$ of total degree less than γ , and a constant N , which does not depend on j or k , such that

$$\|u - P_{j,k}\|_{L_p(Q_{j,k})} \leq N 2^{-j\gamma} |u|_{B_{p,p}^\gamma(Q_{j,k}^\circ)}.$$

Since $\tilde{\psi}_{i,j,k,p'}$ is orthogonal to every polynomial of total degree less than γ , we have

$$\begin{aligned} |\langle u, \tilde{\psi}_{i,j,k,p'} \rangle| &= |\langle u - P_{j,k}, \tilde{\psi}_{i,j,k,p'} \rangle| \\ &\leq \|u - P_{j,k}\|_{L_p(Q_{j,k})} \|\tilde{\psi}_{i,j,k,p'}\|_{L_{p'}(Q_{j,k})} \\ &\leq N 2^{-j\gamma} |u|_{B_{p,p}^\gamma(Q_{j,k}^\circ)} \\ &\leq N 2^{-j\gamma} \rho_{j,k}^{\nu-\gamma} \rho_{j,k}^{\gamma-\nu} |u|_{B_{p,p}^\gamma(Q_{j,k}^\circ)}. \end{aligned}$$

Fix $j \in \mathbb{N}_0$. Summing over all indices $(i, j, k) \in \Lambda_j^0$ and applying Hölder's inequality with exponents $p/\tau > 1$ and $p/(p-\tau)$ one finds

$$\begin{aligned} \sum_{(i,j,k) \in \Lambda_j^0} |\langle u, \tilde{\psi}_{i,j,k,p'} \rangle|^\tau &\leq N \sum_{(i,j,k) \in \Lambda_j^0} 2^{-j\gamma\tau} \rho_{j,k}^{(\nu-\gamma)\tau} \rho_{j,k}^{(\gamma-\nu)\tau} |u|_{B_{p,p}^\gamma(Q_{j,k}^\circ)}^\tau \\ &\leq N \left(\sum_{(i,j,k) \in \Lambda_j^0} \left(\rho_{j,k}^{\gamma-\nu} |u|_{B_{p,p}^\gamma(Q_{j,k}^\circ)} \right)^p \right)^{\frac{\tau}{p}} \left(\sum_{(i,j,k) \in \Lambda_j^0} 2^{-j\frac{\gamma\tau p}{p-\tau}} \rho_{j,k}^{\frac{(\nu-\gamma)\tau p}{p-\tau}} \right)^{\frac{p-\tau}{p}}. \end{aligned}$$

Now we use Lemma 6.8 to obtain

$$\sum_{(i,j,k) \in \Lambda_j^0} |\langle u, \tilde{\psi}_{i,j,k,p'} \rangle|^\tau \leq N \|u\|_{H_{p,d-\nu p}^\gamma(\mathcal{O})}^\tau \left(\sum_{(i,j,k) \in \Lambda_j^0} 2^{-j\frac{\gamma\tau p}{p-\tau}} \rho_{j,k}^{\frac{(\nu-\gamma)\tau p}{p-\tau}} \right)^{\frac{p-\tau}{p}}, \quad (6.18)$$

with a constant N , which does not depend on the level j . In order to estimate the sum on the right hand side we use the Lipschitz character of the domain \mathcal{O} which implies that

$$|\Lambda_{j,m}| \leq N 2^{j(d-1)} \quad \text{for all } j, m \in \mathbb{N}_0. \quad (6.19)$$

Moreover, the boundedness of \mathcal{O} yields $\Lambda_{j,m} = \emptyset$ for all $j, m \in \mathbb{N}_0$ with $m \geq N2^j$, where the constant N does not depend on j or m . Consequently,

$$\begin{aligned} \left(\sum_{(i,j,k) \in \Lambda_j^0} 2^{-j\frac{\gamma\tau p}{p-\tau}} \rho_{j,k}^{\frac{(\nu-\gamma)\tau p}{p-\tau}} \right)^{\frac{p-\tau}{p}} &\leq \left(\sum_{m=1}^{N2^j} \sum_{(i,j,k) \in \Lambda_{j,m}} 2^{-j\frac{\gamma\tau p}{p-\tau}} \rho_{j,k}^{\frac{(\nu-\gamma)\tau p}{p-\tau}} \right)^{\frac{p-\tau}{p}} \\ &\leq N \left(\sum_{m=1}^{N2^j} 2^{j(d-1)} 2^{-j\frac{\gamma\tau p}{p-\tau}} (m 2^{-j})^{\frac{(\nu-\gamma)\tau p}{p-\tau}} \right)^{\frac{p-\tau}{p}} \\ &\leq N \left(2^{j(d-1-\frac{\nu p \tau}{p-\tau})} + 2^{j(d-\frac{\gamma p \tau}{p-\tau})} \right)^{\frac{p-\tau}{p}}. \end{aligned} \quad (6.20)$$

Now, let us sum over all $j \in \mathbb{N}_0$. Inequalities (6.20) and (6.18) imply

$$\sum_{(i,j,k) \in \Lambda^0} |\langle u, \tilde{\psi}_{i,j,k,p'} \rangle|^\tau \leq N \sum_{j \in \mathbb{N}_0} \left(2^{j(d-1-\frac{\nu p \tau}{p-\tau})} + 2^{j(d-\frac{\gamma p \tau}{p-\tau})} \right)^{\frac{p-\tau}{p}} \|u\|_{H_{p,d-\nu p}^\gamma(\mathcal{O})}^\tau.$$

Obviously, the sums on the right hand side converge if, and only if, $\alpha \in \left(0, \gamma \wedge \nu \frac{d}{d-1}\right)$. Finally,

$$\sum_{(i,j,k) \in \Lambda^0} |\langle u, \tilde{\psi}_{i,j,k,p'} \rangle|^\tau \leq N \|u\|_{H_{p,d-\nu p}^\gamma(\mathcal{O})}^\tau.$$

Now we estimate the second term II in (6.17). First we fix $j \in \mathbb{N}_0$ and use Hölder's inequality and (6.19) to obtain

$$\sum_{(i,j,k) \in \Lambda_{j,0}} |\langle u, \tilde{\psi}_{i,j,k,p'} \rangle|^\tau \leq N 2^{j(d-1)\frac{p-\tau}{p}} \left(\sum_{(i,j,k) \in \Lambda_{j,0}} |\langle u, \tilde{\psi}_{i,j,k,p'} \rangle|^p \right)^{\frac{\tau}{p}}.$$

Summing over all $j \in \mathbb{N}_0$ and using Hölder's inequality again yields

$$\begin{aligned} \sum_{(i,j,k) \in \Lambda \setminus \Lambda^0} |\langle u, \tilde{\psi}_{i,j,k,p'} \rangle|^\tau &= \sum_{j \in \mathbb{N}_0} \left[\sum_{(i,j,k) \in \Lambda_{j,0}} |\langle u, \tilde{\psi}_{i,j,k,p'} \rangle|^\tau \right] \\ &\leq N \sum_{j \in \mathbb{N}_0} \left[2^{j(d-1)\frac{p-\tau}{p}} \left(\sum_{(i,j,k) \in \Lambda_{j,0}} |\langle u, \tilde{\psi}_{i,j,k,p'} \rangle|^p \right)^{\frac{\tau}{p}} \right] \\ &\leq N \left(\sum_{j \in \mathbb{N}_0} 2^{j\left(\frac{(d-1)(p-\tau)}{p} - \nu\tau\right)\frac{p}{p-\tau}} \right)^{\frac{p-\tau}{p}} \left(\sum_{j \in \mathbb{N}_0} \sum_{(i,j,k) \in \Lambda_{j,0}} 2^{j\nu p} |\langle u, \tilde{\psi}_{i,j,k,p'} \rangle|^p \right)^{\frac{\tau}{p}}. \end{aligned}$$

Using Theorem 6.3 and the boundedness of the extension operator, we obtain

$$\sum_{(i,j,k) \in \Lambda \setminus \Lambda^0} |\langle u, \tilde{\psi}_{i,j,k,p'} \rangle|^\tau \leq N \|u\|_{B_{p,p}^\nu(\mathcal{O})}^\tau \left(\sum_{j \in \mathbb{N}_0} 2^{j\left(\frac{(d-1)(p-\tau)}{p} - \nu\tau\right)\frac{p}{p-\tau}} \right)^{\frac{p-\tau}{p}}.$$

The series on the right hand side converges if, and only if, $\alpha \in \left(0, \nu \frac{d}{d-1}\right)$. We thus have

$$\sum_{(i,j,k) \in \Lambda \setminus \Lambda^0} |\langle u, \tilde{\psi}_{i,j,k,p'} \rangle|^\tau \leq N \|u\|_{B_{p,p}^\nu(\mathcal{O})}^\tau \leq N \|u\|_{H_{p,d-\nu p}^\gamma(\mathcal{O})}^\tau. \quad \square$$

7 Hölder–Besov regularity for elements of $\mathfrak{H}_{p,\theta}^{\gamma,q}(\mathcal{O}, T)$ and implications for SPDEs

In this section, we state and prove our second main result concerning the time-space regularity of the solutions to SPDEs of the form (1.1) on bounded Lipschitz domains. We use the scale $(*)$ to measure the regularity in space, whereas the time-regularity will be measured in terms of Hölder norms. Since the stochastic parabolic weighted Sobolev spaces $\mathfrak{H}_{p,\theta}^{\gamma,q}(\mathcal{O}, T)$ are the right spaces to construct a solvability theory for SPDEs, we will first formulate our results in terms of these spaces. As a consequence, each result about existence of solutions to SPDEs in these spaces automatically encodes a statement about the Hölder-Besov regularity of the solution. The corresponding results for the solutions in the different settings from Section 4 will be presented here in detail. We will use the following short notation

$$L_q(\Omega_T; B_{\tau,\tau}^\alpha(\mathcal{O})) := L_q(\Omega \times [0, T], \mathcal{P}, \mathbb{P} \otimes dt; B_{\tau,\tau}^\alpha(\mathcal{O}))$$

Let us first clarify for which range of $\alpha > 0$ a stochastic process $u \in \mathfrak{H}_{p,\theta}^{\gamma,q}(\mathcal{O}, T)$ takes values in $B_{\tau,\tau}^\alpha(\mathcal{O})$, $1/\tau = \alpha/d + 1/p$.

Theorem 7.1. *Let \mathcal{O} be a bounded Lipschitz domain in \mathbb{R}^d . Let $\gamma + 2 \in (0, \infty)$, $p, q \in [2, \infty)$, $\theta \in \mathbb{R}$, and $u \in \mathbb{H}_{p, \theta-p}^{\gamma+2, q}(\mathcal{O}, T)$. Then,*

$$u \in L_q(\Omega_T; B_{\tau, \tau}^\alpha(\mathcal{O})), \quad \frac{1}{\tau} = \frac{\alpha}{d} + \frac{1}{p}, \quad \text{for all } 0 < \alpha < \min \left\{ \gamma + 2, \left(1 + \frac{d-\theta}{p}\right) \frac{d}{d-1} \right\}. \quad (7.1)$$

Moreover, for α fulfilling (7.1), there exists a constant N which does not depend on u , such that

$$\mathbb{E} \left[\int_0^T \|u(t, \cdot)\|_{B_{\tau, \tau}^\alpha(\mathcal{O})}^q dt \right] \leq N \|u\|_{\mathbb{H}_{p, \theta-p}^{\gamma+2, q}(\mathcal{O}, T)}^q.$$

Proof. This is a direct consequence of Theorem 6.9. \square

Combining this assertion with the results from Section 4 we obtain the following spatial regularity results for SPDEs.

Theorem 7.2. *Let \mathcal{O} be a bounded Lipschitz domain in \mathbb{R}^d .*

(i) *Given the setting of Theorem 4.1, the solution $u \in \mathfrak{H}_{p, \theta}^{\gamma+2, p}(\mathcal{O}, T)$ of Eq. (3.1) fulfils (7.1). Moreover, for any α in (7.1), there exists a constant N , which does not depend on u , f , g and u_0 , such that*

$$\mathbb{E} \left[\int_0^T \|u(t, \cdot)\|_{B_{\tau, \tau}^\alpha(\mathcal{O})}^p dt \right] \leq N \left(\|f\|_{\mathbb{H}_{p, \theta+p}^{\gamma, p}(\mathcal{O}, T)}^p + \|g\|_{\mathbb{H}_{\theta, p}^{\gamma+1, p}(\mathcal{O}, T; \ell_2)}^p + \|u_0\|_{U_{p, \theta}^{\gamma, p}(\mathcal{O})}^p \right).$$

(ii) *Given the setting of Theorem 4.4, the solution $u \in \mathfrak{H}_{p, d}^{\gamma+2, q}(\mathcal{O}, T)$ of Eq. (4.12) fulfils (7.1) with $\theta = d$. Moreover, for any α in (7.1), there exists a constant N , which does not depend on u , f , and g , such that*

$$\mathbb{E} \left[\int_0^T \|u(t, \cdot)\|_{B_{\tau, \tau}^\alpha(\mathcal{O})}^q dt \right] \leq N \left(\|f\|_{\mathbb{H}_{p, d}^{0, q}(\mathcal{O}, T)}^q + \|f\|_{\mathbb{H}_{p, d+p}^{\gamma, q}(\mathcal{O}, T)}^q + \|g\|_{\mathbb{H}_{d, p}^{\gamma+1, p}(\mathcal{O}, T; \ell_2)}^q \right).$$

(iii) *Given the setting of Theorem 4.2, the solution $u \in \mathfrak{H}_{p, d}^{\gamma+2, q}(\mathcal{O}, T)$ of Eq. (4.1) fulfils (7.1) with $\theta = d$. Moreover, for any α in (7.1), there exists a constant N , which does not depend on u , f , and g , such that*

$$\begin{aligned} \mathbb{E} \left[\int_0^T \|u(t, \cdot)\|_{B_{\tau, \tau}^\alpha(\mathcal{O})}^q dt \right] &\leq N \left(\|f\|_{\mathbb{H}_{p, d+p}^{\gamma, q}(\mathcal{O}, T)}^q + \|f\|_{\mathbb{H}_{p, d}^{0, q}(\mathcal{O}, T)}^q \right. \\ &\quad \left. + \|g\|_{\mathbb{H}_{p, d}^{\gamma+1, q}(\mathcal{O}, T; \ell_2)}^q + \|g\|_{\mathbb{H}_{p, d-p}^{1, q}(\mathcal{O}, T; \ell_2)}^q \right). \end{aligned}$$

Proof. The assertions are immediate consequences of Theorem 7.1 and the corresponding existence results from Section 4. \square

Remark 7.3. *A result similar to Theorem 7.2(i) has been proved in [4, Theorem 3.1, see also Theorem B.3]. There are three major improvements in Theorem 7.2(i) compared to [4, Theorem 3.1]. Firstly, we have no restriction on $\gamma \in (0, \infty)$, whereas in [4] only integer $\gamma \in \mathbb{N}_0$ are considered. Secondly, we obtain L_p -integrability in time of the $B_{\tau, \tau}^\alpha(\mathcal{O})$ -valued process for arbitrary $p \in [2, \infty)$. With the techniques used in [4] just L_τ -integrability in time can be established. Thirdly, we do not need the extra assumption $u \in L_p([0, T] \times \Omega; B_{p, p}^s(\mathcal{O}))$ for some $s > 0$. It suffices that $u \in \mathbb{H}_{p, \theta-p}^{\gamma+2, p}(\mathcal{O}, T)$. Note that, obviously, this improvements also hold for the solutions of more general equations as considered in [4, Theorem B.3].*

Here is the main result of this section. It concerns the Hölder-Besov regularity of processes in $\mathfrak{H}_{p,\theta}^{\gamma,q}(\mathcal{O}, T)$.

Theorem 7.4. *Let \mathcal{O} be a bounded Lipschitz domain in \mathbb{R}^d . Let $2 \leq p \leq q < \infty$, $\gamma + 2 \in \mathbb{N}$, $\theta \in \mathbb{R}$, and $u \in \mathfrak{H}_{p,\theta}^{\gamma+2,q}(\mathcal{O}, T)$. Moreover, let*

$$\frac{2}{q} < \bar{\beta} < \min \left\{ 1, 1 + \frac{d - \theta}{p} \right\}.$$

Then, for all α and τ with

$$\frac{1}{\tau} = \frac{\alpha}{d} + \frac{1}{p} \quad \text{and} \quad 0 < \alpha < \min \left\{ \gamma + 2 - \bar{\beta}, \left(1 + \frac{d - \theta}{p} - \bar{\beta} \right) \frac{d}{d - 1} \right\}, \quad (7.2)$$

we have

$$\mathbb{E}[u]_{\mathcal{C}^{\bar{\beta}/2-1/q}([0,T]; B_{\tau,\tau}^{\alpha}(\mathcal{O}))}^q \leq N(T) \left(\|u\|_{\mathbb{H}_{p,\theta-p}^{\gamma+2,q}(\mathcal{O},T)}^q + \|\mathbb{D}u\|_{\mathbb{H}_{p,\theta+p}^{\gamma,q}(\mathcal{O},T)}^q + \|\mathbb{S}u\|_{\mathbb{H}_{p,\theta}^{\gamma+1,q}(\mathcal{O},T;\ell_2)}^q \right),$$

and

$$\mathbb{E}\|u\|_{\mathcal{C}^{\bar{\beta}/2-1/q}([0,T]; B_{\tau,\tau}^{\alpha}(\mathcal{O}))}^q \leq N(T) \|u\|_{\mathfrak{H}_{p,\theta}^{\gamma+2,q}(\mathcal{O},T)}^q.$$

The constants $N(T)$ are given by $N(T) = N \sup_{\beta \in [\bar{\beta}, 1]} \{T^{(\beta - \bar{\beta})q/2}\}$, with N from (5.1) and (5.2) respectively.

Proof. The assertion is an immediate consequence of Theorem 5.1 and Theorem 6.9. \square

We obtain the following implications concerning the path regularity of the solutions of SPDEs from Section 4.

Theorem 7.5. *Let \mathcal{O} be a bounded Lipschitz domain in \mathbb{R}^d .*

(i) *Given the setting of Theorem 4.4 with $\gamma \in \mathbb{N}_0$, let $u \in \mathfrak{H}_{p,d}^{\gamma+2,q}(\mathcal{O}, T)$ be the solution of Eq. (4.12). Assume furthermore that $2/q < \bar{\beta} < 1$, and that α and τ fulfil (7.2). Then,*

$$\mathbb{E}\|u\|_{\mathcal{C}^{\bar{\beta}/2-1/q}([0,T]; B_{\tau,\tau}^{\alpha}(\mathcal{O}))}^q \leq N \left(\|f\|_{\mathbb{H}_{p,d+p}^{\gamma,q}(\mathcal{O},T)}^q + \|f\|_{\mathbb{H}_{p,d}^{0,q}(\mathcal{O},T)}^q + \|g\|_{\mathbb{H}_{p,d}^{\gamma+1,q}(\mathcal{O},T;\ell_2)}^q \right),$$

where the constant N does not depend on u , f and g .

(ii) *Given the setting of Theorem 4.2 with $\gamma \in \mathbb{N}_0$, let $u \in \mathfrak{H}_{p,d}^{\gamma+2,q}(\mathcal{O}, T)$ be the solution of Eq. (4.1). Assume furthermore that $2/q < \bar{\beta} < 1$, and that α and τ fulfil (7.2). Then,*

$$\begin{aligned} \mathbb{E}\|u\|_{\mathcal{C}^{\bar{\beta}/2-1/q}([0,T]; B_{\tau,\tau}^{\alpha}(\mathcal{O}))}^q &\leq N \left(\|f\|_{\mathbb{H}_{p,d+p}^{\gamma,q}(\mathcal{O},T)}^q + \|f\|_{\mathbb{H}_{p,d}^{0,q}(\mathcal{O},T)}^q \right. \\ &\quad \left. + \|g\|_{\mathbb{H}_{p,d}^{\gamma+1,q}(\mathcal{O},T;\ell_2)}^q + \|g\|_{\mathbb{H}_{p,d-p}^{1,q}(\mathcal{O},T;\ell_2)}^q \right), \end{aligned}$$

where the constant N does not depend on u , f and g .

Proof. The assertions are immediate consequences of Theorem 7.4 and the corresponding existence results from Section 4. \square

A Appendix

Proof of Lemma 5.4. We consider consecutively the cases $\gamma = 0, 1, -1$. For fractional $\gamma \in (-1, 1)$, the statement follows then by using interpolation arguments, see [38, Proposition 2.4]. Furthermore, we resume ourselves to the proof of the right inequality in the assertion of the Lemma, i.e., that there exists a constant $N = N(d, \gamma, p, \theta, \phi) \in (0, \infty)$, such that for any $u \in H_{p,\theta}^\gamma(G^{(1)})$ the following inequality holds:

$$\|u \circ \phi^{-1}\|_{H_{p,\theta}^\gamma(G^{(2)})} \leq N \|u\|_{H_{p,\theta}^\gamma(G^{(1)})}.$$

The left inequality can be proven analogously. For $\gamma = 0$, the assertion follows immediately by using the assumptions of the Lemma and the change of variables formula for bi-Lipschitz transformations, see, e.g., [18, Theorem 3]. Let us go on and look at the case $\gamma = 1$. Because of the density of the test functions $C_0^\infty(G^{(1)})$ in $H_{p,\theta}^1(G^{(1)})$, it suffices to prove the asserted inequality for $u \in C_0^\infty(G^{(1)})$. In this case, because of the assumed Lipschitz-continuity of ϕ^{-1} , the classical partial derivatives of $u \circ \phi^{-1}$ exist a.e. and

$$\left| \frac{\partial}{\partial x^j} (u \circ \phi^{-1}) \right| = \left| \sum_{i=1}^d \left(\frac{\partial}{\partial x^i} u \right) \circ \phi^{-1} \frac{\partial}{\partial x^j} \phi_i^{-1} \right| \leq N \sum_{i=1}^d \left| \left(\frac{\partial}{\partial x^i} u \right) \circ \phi^{-1} \right| \quad (\text{a.e.}).$$

Thus, using e.g. [39, Section 1.1.3, Theorem 2] we can conclude that these a.e. existing classical derivatives coincide with the weak derivatives, so that

$$\begin{aligned} & \|u \circ \phi^{-1}\|_{H_{p,\theta}^1(G^{(2)})}^p \\ & \leq N \left(\int_{G^{(2)}} |(u \circ \phi^{-1})(y)|^p \rho_{G^{(2)}}(y)^{\theta-d} dy + \sum_{j=1}^d \int_{G^{(2)}} \left| \frac{\partial}{\partial x^j} (u \circ \phi^{-1})(y) \right|^p \rho_{G^{(2)}}(y)^{p+\theta-d} dy \right) \\ & \leq N \left(\int_{G^{(2)}} |(u \circ \phi^{-1})(y)|^p \rho_{G^{(2)}}(y)^{\theta-d} dy + \int_{G^{(2)}} \sum_{i=1}^d \left| \left(\frac{\partial}{\partial x^i} u \right) (\phi^{-1}(y)) \right|^p \rho_{G^{(2)}}(y)^{p+\theta-d} dy \right) \\ & \leq N \left(\int_{G^{(1)}} |u(x)|^p \rho_{G^{(1)}}(x)^{\theta-d} dx + \sum_{i=1}^d \int_{G^{(1)}} \left| \frac{\partial}{\partial x^i} u(x) \right|^p \rho_{G^{(1)}}(x)^{p+\theta-d} dx \right) \\ & \leq N \|u\|_{H_{p,\theta}^1(G^{(1)})}^p. \end{aligned}$$

Finally, for $\gamma = -1$, we can use the fact that $H_{p',\theta'}^1(G)$ is the dual space of $H_{p,\theta}^{-1}(G)$, if $1/p + 1/p' = 1$ and $\theta/p + \theta'/p' = d$, see [38, Proposition 2.4], and fall back to the cases we have already proven. \square

Proof of Lemma 5.5. We set $f := \mathbb{D}u$ and $g := \mathbb{S}u$. Since $u \in \mathfrak{H}_{p,\theta}^{1,q}(G^{(1)}, T)$, Lemma 5.4 guarantees that $f \circ \phi^{-1} \in \mathbb{H}_{p,\theta+p}^{-1,q}(G^{(2)}, T)$, $g \circ \phi^{-1} \in \mathbb{H}_{p,\theta}^{0,q}(G^{(2)}, T; \ell_2)$ and $u(0, \cdot) \circ \phi^{-1} \in U_{p,\theta}^{1,q}(G^{(2)})$. Therefore, we only have to show that for all $\varphi \in C_0^\infty(G^{(2)})$, with probability one, the equality

$$(u(t, \cdot), \varphi \circ \phi) = (u(0, \cdot), \varphi \circ \phi) + \int_0^t (f(s, \cdot), \varphi \circ \phi) ds + \sum_{k=1}^\infty \int_0^t (g^k(s, \cdot), \varphi \circ \phi) dw_s^k \quad (\text{A.1})$$

holds for all $t \in [0, T]$. Thus, let us fix $\varphi \in C_0^\infty(G^{(2)})$. We consider first the case $p > 2$. By Lemma 5.4, $\varphi \circ \phi \in H_{\bar{p}, \bar{\theta}-\bar{p}}^1(G^{(1)})$ for any $\bar{p} \in (1, \infty)$ and $\bar{\theta} \in \mathbb{R}$, hence also for $\bar{p} := 2p/(p-2)$ and

$\bar{\theta} := 2\theta'(p-1)/(p-2) - dp/(p-2)$, where $\theta/p + \theta'/p' = d$ with $1/p + 1/p' = 1$. Moreover, by Lemma 2.1(i) we can choose a sequence $\bar{\varphi}_n \subseteq C_0^\infty(G^{(1)})$ approximating $\varphi \circ \phi$ in $H_{\bar{p}, \bar{\theta} - \bar{p}}^1(G^{(1)})$. Furthermore, a consequence of our assumptions is, that for all $n \in \mathbb{N}$, with probability one, the equality

$$(u(t, \cdot), \bar{\varphi}_n) = (u(0, \cdot), \bar{\varphi}_n) + \int_0^t (f(s, \cdot), \bar{\varphi}_n) ds + \sum_{k=1}^{\infty} \int_0^t (g^k(s, \cdot), \bar{\varphi}_n) dw_s^k \quad (\text{A.2})$$

holds for all $t \in [0, T]$. Thus, if we can show that each side of (A.2) converges in $L_2(\Omega; C([0, T]))$ to the respective side of (A.1), the assertion follows. To this end, let us fix an arbitrary $\bar{v} \in H_{\bar{p}, \bar{\theta} - \bar{p}}^1(G^{(1)})$. A standard estimate yields

$$\mathbb{E} \left[\sup_{t \in [0, T]} \left| \int_0^t (f(s, \cdot), \bar{v}) ds \right|^2 \right] \leq N \|f\|_{\mathbb{H}_{p, \theta + p}^{-1, q}(G^{(1)}, T)}^2 \|\bar{v}\|_{H_{p', \theta' - p'}^1(G^{(1)})}^2.$$

Also, using Doob's inequality, Itô's isometry and Hölder's inequality we get

$$\mathbb{E} \left[\sup_{t \in [0, T]} \left| \sum_{k \in \mathbb{N}} \int_0^t (g^k(s, \cdot), \bar{v}) dw_s^k \right|^2 \right] \leq N \|g\|_{\mathbb{H}_{p, \theta}^{0, q}(G^{(1)}, T; \ell_2)}^2 \|\bar{v}\|_{L_{\bar{p}, \bar{\theta}}^2(G^{(1)})}^2. \quad (\text{A.3})$$

Furthermore, an application of [26, Proposition 2.9] and the fact that $q \geq p$ lead to

$$\begin{aligned} \mathbb{E} \left[\sup_{t \in [0, T]} |(u(t, \cdot), \bar{v})|^2 \right] &\leq \left(\mathbb{E} \left[\sup_{t \in [0, T]} \|u(t, \cdot)\|_{L_{p, \theta}(G^{(1)})}^p \right] \right)^{2/p} \|\bar{v}\|_{L_{p', \theta'}^2(G^{(1)})}^2 \\ &\leq N \|u\|_{\mathfrak{H}_{p, \theta}^{1, p}(G^{(1)}, T)}^2 \|\bar{v}\|_{L_{p', \theta'}^2(G^{(1)})}^2 \\ &\leq N \|u\|_{\mathfrak{H}_{p, \theta}^{1, q}(G^{(1)}, T)}^2 \|\bar{v}\|_{L_{p', \theta'}^2(G^{(1)})}^2. \end{aligned}$$

Hence, since $H_{\bar{p}, \bar{\theta} - \bar{p}}^1(G^{(1)})$ is continuously embedded in $H_{p', \theta' - p'}^1(G^{(1)}) \cap L_{p', \theta'}(G^{(1)}) \cap L_{\bar{p}, \bar{\theta}}(G^{(1)})$, the assertion follows for the case $p > 2$. The same arguments can be used to prove the case $p = 2$: Just replace \bar{p} by 2, and $\bar{\theta}$ by $\theta' = 2d - \theta$ and use the estimate

$$\mathbb{E} \left[\sup_{t \in [0, T]} \left| \sum_{k \in \mathbb{N}} \int_0^t (g^k(s, \cdot), \bar{v}) dw_s^k \right|^2 \right] \leq N \|g\|_{\mathbb{H}_{2, \theta}^{0, q}(G^{(1)}, T; \ell_2)}^2 \|\bar{v}\|_{L_{2, \theta'}^2(G^{(1)})}^2$$

instead of (A.3). □

Proof of Lemma 6.8. Let us fix $j \in \mathbb{N}_0$. We use the notations introduced before Lemma 6.8. Remember that $M > 0$ has been chosen in such a way that $\mathcal{O} \subseteq [-M, M]^d$. Let us fix $k_1 \geq 1$ such that

$$2 + 2M\sqrt{d} < 2^{k_1}, \quad (\text{A.4})$$

and construct a sequence $\{\xi_n : n \in \mathbb{Z}\} \subseteq C_0^\infty(\mathcal{O})$ as in Remark 2.4(ii). In order to prove the assertion we are going to show the estimates

$$\sum_{(i, j, k) \in \Lambda_j^0} \left(\rho_{j, k}^{\gamma - \nu} |u|_{B_{p, p}^\gamma(Q_{j, k}^\circ)} \right)^p \leq N \sum_{n \in \mathbb{N}_0} 2^{-(j-n)(\gamma - \nu)p} |\xi_{j-n} u|_{B_{p, p}^\gamma(\mathbb{R}^d)}^p, \quad (\text{A.5})$$

and

$$|\xi_{j-n}u|_{B_{p,p}^\gamma(\mathbb{R}^d)}^p \leq N 2^{-(j-n)(d-\gamma p)} \|\xi_{j-n}(2^{-(j-n)} \cdot)u(2^{-(j-n)} \cdot)\|_{H_p^\gamma(\mathbb{R}^d)}^p, \quad (\text{A.6})$$

where the constant N does not depend on j and n . This will prove the assertion since, assuming that (A.5) and (A.6) are true, their combination gives

$$\begin{aligned} \sum_{(i,j,k) \in \Lambda_j^0} (\rho_{j,k}^{\gamma-\nu} |u|_{B_{p,p}^\gamma(Q_{j,k}^\circ)})^p &\leq N \sum_{n \in \mathbb{N}_0} 2^{-(j-n)(d-\nu p)} \|\xi_{j-n}(2^{-(j-n)} \cdot)u(2^{-(j-n)} \cdot)\|_{H_p^\gamma(\mathbb{R}^d)}^p \\ &\leq N \sum_{n \in \mathbb{Z}} 2^{n(d-\nu p)} \|\xi_{-n}(2^n \cdot)u(2^n \cdot)\|_{H_p^\gamma(\mathbb{R}^d)}^p \\ &\leq N \|u\|_{H_{p,d-\nu p}^\gamma(\mathcal{O})}^p. \end{aligned}$$

In the last step we used Remark 2.4(ii) and Lemma 2.3.

Let us first verify inequality (A.6). To this end, let r be the smallest integer strictly greater than γ . For the sake of clarity we use here the notation $\Delta_h^r[f]$ for the r -th difference of a function $f : \mathbb{R}^d \rightarrow \mathbb{R}$ with step $h \in \mathbb{R}^d$, whereas $\Delta_h^r[f](x)$ denotes the value of this r -th difference at a point $x \in \mathbb{R}^d$, compare Subsection 6.1. Writing out the Besov semi-norm and applying the transformation formula for integrals we see that

$$\begin{aligned} |\xi_{j-n}u|_{B_{p,p}^\gamma(\mathbb{R}^d)}^p &= \int_0^\infty t^{-\gamma p} \sup_{|h|<t} \|\Delta_h^r[\xi_{j-n}u]\|_{L_p(\mathbb{R}^d)}^p \frac{dt}{t} \\ &= 2^{-(j-n)d} \int_0^\infty t^{-\gamma p} \sup_{|h|<t} \left\{ \int_{\mathbb{R}^d} |\Delta_h^r[\xi_{j-n}u](2^{-(j-n)}x)|^p dx \right\} \frac{dt}{t} \\ &= 2^{-(j-n)d} \int_0^\infty t^{-\gamma p} \sup_{|h|<2^{j-n}t} \left\{ \int_{\mathbb{R}^d} |\Delta_h^r[\xi_{j-n}(2^{-(j-n)} \cdot)u(2^{-(j-n)} \cdot)](x)|^p dx \right\} \frac{dt}{t}. \end{aligned}$$

A further application of the transformation formula for integrals yields

$$\begin{aligned} |\xi_{j-n}u|_{B_{p,p}^\gamma(\mathbb{R}^d)}^p &= 2^{-(j-n)d} 2^{(j-n)\gamma p} \int_0^\infty t^{-\gamma p} \sup_{|h|<t} \left\| \Delta_h^r \left[\xi_{j-n}(2^{-(j-n)} \cdot)u(2^{-(j-n)} \cdot) \right] \right\|_{L_p(\mathbb{R}^d)}^p \frac{dt}{t} \\ &= 2^{-(j-n)(d-\gamma p)} |\xi_{j-n}(2^{-(j-n)} \cdot)u(2^{-(j-n)} \cdot)|_{B_{p,p}^\gamma(\mathbb{R}^d)}^p, \end{aligned}$$

which implies (A.6) since the space $H_p^\gamma(\mathbb{R}^d)$ of Bessel potentials is continuously embedded in the Besov space $B_{p,p}^\gamma(\mathbb{R}^d)$, see [43, Theorem 2.3.2(d) combined with Theorem 2.3.3(a)].

It remains to prove inequality (A.5). Recall that the index i referring to the different types of wavelets on a cube $Q_{j,k}$ ranges from 1 to $2^d - 1$. Since Λ_j^0 consists of those indices $(i, j, k) \in \Lambda_j$ with $2^{-j} \leq \rho_{j,k}$, we have

$$\sum_{(i,j,k) \in \Lambda_j^0} (\rho_{j,k}^{\gamma-s} |u|_{B_{p,p}^\gamma(Q_{j,k})})^p = (2^d - 1) \sum_{k \in \Lambda_j^*} (\rho_{j,k}^{\gamma-s} |u|_{B_{p,p}^\gamma(Q_{j,k})})^p, \quad (\text{A.7})$$

where we used the notation

$$\Lambda_j^* := \{k \in \mathbb{Z}^d : (i, j, k) \in \Lambda_j^0\}.$$

Now we get the required estimate in four steps.

Step 1. We first show that the cubes supporting the wavelets fit into the stripes where the cut-off functions (ξ_n) are identical to one. More precisely, we claim that the proper choice of k_1 , see (A.4), leads to the fact that, for any $k \in \Lambda_j^*$, there exists a non-negative integer $n \in \mathbb{N}_0$ such that

$$Q_{j,k} \subseteq S_{j-n} := \rho^{-1}(2^{-(j-n)}[2^{-k_1}, 2^{k_1}]).$$

We denote

$$S_{j,n}^* := \{k \in \Lambda_j^* : Q_{j,k} \in S_{j-n}\}, \quad n \in \mathbb{N}_0.$$

To prove this, we first note that, since $k_1 \geq 1$ fulfils (A.4),

$$\bigcup_{k \in \Lambda_j^*} Q_{j,k} \subseteq \bigcup_{n \in \mathbb{N}_0} S_{j-n} = \bigcup_{n=0}^j S_{j-n}.$$

Fix $k \in \Lambda_j^*$ and let n^* be the smallest non-negative integer such that $Q_{j,k} \cap S_{j-n^*} \neq \emptyset$, i.e.,

$$n^* := \inf \{n \in \mathbb{N} : Q_{j,k} \cap S_{j-n} \neq \emptyset\} \leq j.$$

Then, there are two possibilities: On the one hand, $Q_{j,k}$ might be contained completely in S_{j-n^*} , i.e., $Q_{j,k} \subseteq S_{j-n^*}$. Then we are done. On the other hand, it might happen that $Q_{j,k}$ is not completely contained in the stripe S_{j-n^*} . In this case, we claim that $Q_{j,k} \subseteq S_{j-(n^*+1)}$, i.e.,

$$\rho(x) \in [2^{-j+n^*+1}2^{-k_1}, 2^{-j+n^*+1}2^{k_1}] \text{ for all } x \in Q_{j,k}.$$

Let us therefore fix $x \in Q_{j,k}$. Then, since the length of the diagonal of $Q_{j,k}$ is $2^{-j}2M\sqrt{d}$, we have

$$\rho(x) \leq \rho_{j,k} + 2^{-j}2M\sqrt{d}.$$

Also, $\rho_{j,k} \leq 2^{-j+n^*}2^{k_1}$ since $Q_{j,k} \cap S_{j-n^*} \neq \emptyset$. Hence,

$$\rho(x) \leq 2^{-j+n^*}2^{k_1} + 2^{-j}2M\sqrt{d}.$$

Since $2M\sqrt{d} \leq 2^{k_1}$, we conclude that

$$\rho(x) \leq 2^{-j+n^*+1}2^{k_1} \left(\frac{1}{2} + \frac{2M\sqrt{d}}{2^{n^*+1}2^{k_1}} \right) \leq 2^{-j+n^*+1}2^{k_1} \left(\frac{1}{2} + \frac{1}{2^{n^*+1}} \right) \leq 2^{-j+n^*+1}2^{k_1}.$$

It remains to show that $\rho(x) \geq 2^{-j+n^*+1}2^{-k_1}$. We argue as follows: Since $Q_{j,k}$ is not completely contained in S_{j-n^*} , there exists a point $x_0 \in Q_{j,k}$ such that $\rho(x_0) > 2^{-j+n^*}2^{k_1}$. Therefore, since the length of the diagonal of $Q_{j,k}$ is $2^{-j}2M\sqrt{d}$, we have

$$\rho(x) > 2^{-j+n^*}2^{k_1} - 2M\sqrt{d}2^{-j} \geq 2^{-j+n^*+1}2^{-k_1} \left(\frac{2^{2k_1}}{2} - \frac{2M\sqrt{d}2^{k_1}}{2^{n^*+1}} \right) \geq 2^{-j+n^*+1}2^{-k_1}.$$

In the last step we used (A.4).

Step 2. We rearrange the cubes supporting the wavelets in classes containing only cubes with disjoint interiors. More precisely, let e_1, \dots, e_d be the canonical orthonormal basis in \mathbb{R}^d . Since $Q_{j,k} = 2^{-j}(k + [-M, M]^d)$ it is clear that, for all $k \in \mathbb{Z}$ and $l \in \{1, \dots, d\}$,

$$Q_{j,k}^\circ \cap (2^{-j}(2M)e_l + Q_{j,k}^\circ) = \emptyset.$$

Consequently, setting $\{a_m : m = 1, \dots, (2M)^d\} := \{0, \dots, 2M - 1\}^d$ and denoting

$$R_{j,m} := \{Q_{j,k} : k \in a_m + 2M\mathbb{Z}^d\} \quad \text{for } m = 1, \dots, (2M)^d,$$

we cover the whole range of cubes, i.e.,

$$\bigcup_{k \in \mathbb{Z}^d} Q_{j,k} = \bigcup_{m=1}^{(2M)^d} R_{j,m},$$

and for any fixed $m \in \{1, \dots, (2M)^d\}$, if $Q_{j,k}, Q_{j,\ell} \in R_{j,m}$ with $k \neq \ell$, then $Q_{j,k}^\circ \cap Q_{j,\ell}^\circ = \emptyset$. In the sequel, we write

$$R_{j,m}^* := \{k \in \Lambda_j^* : Q_{j,k} \in R_{j,m}\}, \quad m \in \{0, \dots, (2M)^d - 1\}.$$

Step 3. Let us fix $k \in \Lambda_j^*$ and concentrate on the Besov semi-norm of the restriction of u to the corresponding cube $Q_{j,k}^\circ$. Using the Peetre K -functional

$$K_r(t, u, Q_{j,k}^\circ)_p := \inf_{g \in W_p^r(Q_{j,k}^\circ)} \left\{ \|u - g\|_{L_p(Q_{j,k})} + t |g|_{W_p^r(Q_{j,k}^\circ)} \right\}$$

and applying [20, Lemma 1] leads to

$$\begin{aligned} |u|_{B_{p,p}^\gamma(Q_{j,k}^\circ)}^p &= \int_0^\infty t^{-\gamma p} \omega^r(t, u, Q_{j,k}^\circ)_p^p \frac{dt}{t} \\ &\leq N \int_0^\infty t^{-\gamma p} K_r(t^r, u, Q_{j,k}^\circ)_p^p \frac{dt}{t} \\ &= N \int_0^\infty t^{-\gamma p} \inf_{g \in W_p^r(Q_{j,k}^\circ)} \left\{ \|u - g\|_{L_p(Q_{j,k})} + t^r |g|_{W_p^r(Q_{j,k}^\circ)} \right\}^p \frac{dt}{t} \\ &\leq N \int_0^\infty t^{-\gamma p} \inf_{g \in W_p^r(Q_{j,k}^\circ)} \left\{ \|u - g\|_{L_p(Q_{j,k})}^p + t^{rp} |g|_{W_p^r(Q_{j,k}^\circ)}^p \right\} \frac{dt}{t}, \end{aligned}$$

where the constant N depends only on r , d and p . (Recall that r is the smallest integer strictly greater than γ .)

Step 4. Now we collect the fruits of our work and approximate the right hand side of (A.7). Because of the first step and since $k_1 \geq 1$ it is easy to see that

$$\Lambda_j^* = \bigcup_{n=0}^j \bigcup_{m=1}^{(2M)^d} S_{j,n}^* \cap R_{j,m}^*,$$

where

$$S_{j,n}^* := \{k \in \Lambda_j^* : Q_{j,k} \in S_{j-n}\}, \quad n \in \mathbb{N}_0.$$

Thus,

$$\sum_{k \in \Lambda_j^*} \left(\rho_{j,k}^{\gamma-\nu} |u|_{B_{p,p}^\gamma(Q_{j,k}^\circ)} \right)^p \leq \sum_{n=0}^j \sum_{m=1}^{(2M)^d} \sum_{k \in S_{j,n}^* \cap R_{j,m}^*} \left(\rho_{j,k}^{(\gamma-\nu)p} |u|_{B_{p,p}^\gamma(Q_{j,k}^\circ)}^p \right). \quad (\text{A.8})$$

Let us fix $n \in \mathbb{N}_0$ such that $S_{j,n}^* \neq \emptyset$ as well as $m \in \{1, \dots, (2M)^d\}$. Then, $\rho_{j,k} \leq 2^{k_1} 2^{-(j-n)}$ for $k \in S_{j,n}^*$, and using the third step we obtain

$$\begin{aligned} & \sum_{k \in S_{j,n}^* \cap R_{j,m}^*} \left(\rho_{j,k}^{(\gamma-\nu)p} |u|_{B_{p,p}^\gamma(Q_{j,k}^\circ)}^p \right) \\ & \leq N \sum_{k \in S_{j,n}^* \cap R_{j,m}^*} \left(2^{-(j-n)(\gamma-\nu)p} \int_0^\infty t^{-\gamma p} \inf_{g \in W_p^r(Q_{j,k}^\circ)} \left\{ \|u - g\|_{L_p(Q_{j,k}^\circ)}^p + t^{rp} |g|_{W_p^r(Q_{j,k}^\circ)}^p \right\} \frac{dt}{t} \right) \\ & \leq N 2^{-(j-n)(\gamma-\nu)p} \sum_{k \in S_{j,n}^* \cap R_{j,m}^*} \int_0^\infty t^{-\gamma p} \inf_{g \in W_p^r(\mathcal{O})} \left\{ \|u - g\|_{L_p(Q_{j,k}^\circ)}^p + t^{rp} |g|_{W_p^r(Q_{j,k}^\circ)}^p \right\} \frac{dt}{t}. \end{aligned}$$

Furthermore, since $\xi_{j-n} = 1$ on $Q_{j,k}$ for any $k \in S_{j,n}^*$ and since $Q_{j,k}^\circ \cap Q_{j,\ell}^\circ = \emptyset$ for $k, \ell \in R_{j,m}^*$ if $k \neq \ell$, we can continue our estimate as follows:

$$\begin{aligned} & \sum_{k \in S_{j,n}^* \cap R_{j,m}^*} \left(\rho_{j,k}^{(\gamma-\nu)p} |u|_{B_{p,p}^\gamma(Q_{j,k}^\circ)}^p \right) \\ & \leq N 2^{-(j-n)(\gamma-\nu)p} \int_0^\infty t^{-\gamma p} \inf_{g \in W_p^r(\mathcal{O})} \left\{ \sum_{k \in S_{j,n}^* \cap R_{j,m}^*} \left(\|\xi_{j-n} u - g\|_{L_p(Q_{j,k}^\circ)}^p + t^{rp} |g|_{W_p^r(Q_{j,k}^\circ)}^p \right) \right\} \frac{dt}{t} \\ & \leq N 2^{-(j-n)(\gamma-\nu)p} \int_0^\infty t^{-\gamma p} \inf_{g \in W_p^r(\mathcal{O})} \left\{ \|\xi_{j-n} u - g\|_{L_p(\mathcal{O})}^p + t^{rp} |g|_{W_p^r(\mathcal{O})}^p \right\} \frac{dt}{t} \\ & \leq N 2^{-(j-n)(\gamma-\nu)p} \int_0^\infty t^{-\gamma p} K_r(t^r, \xi_{j-n} u, \mathcal{O})_p^p \frac{dt}{t}. \end{aligned}$$

By [20, Theorem 1], we know that there exists a constant N , depending only on r , p and \mathcal{O} , such that

$$K_r(t^r, \xi_{j-n} u, \mathcal{O})_p \leq N \omega^r(t, \xi_{j-n} u, \mathcal{O})_p.$$

Putting everything together, we have shown that there exists a constant N which does not depend on j , n or m such that

$$\begin{aligned} \sum_{k \in S_{j,n}^* \cap R_{j,m}^*} \left(\rho_{j,k}^{(\gamma-\nu)p} |u|_{B_{p,p}^\gamma(Q_{j,k}^\circ)}^p \right) & \leq N 2^{-(j-n)(\gamma-\nu)p} \int_0^\infty t^{-\gamma p} \omega^r(t, \xi_{j-n} u, \mathcal{O})_p^p \frac{dt}{t} \\ & = N 2^{-(j-n)(\gamma-\nu)p} |u|_{B_{p,p}^\gamma(\mathcal{O})}^p. \end{aligned}$$

Hence, (A.5) follows after inserting this estimate into (A.8) and using (A.7). \square

Acknowledgements. The second and the third author are thankful for the hospitality of the other authors during their stay at TU Dresden. The first and last author are deeply grateful for the hospitality of the other authors during their stay at the 2nd NIMS Summer School in Probability Theory in Daejeon and during their stay at Korea University.

References

- [1] Hugo Aimar and Ivana Gómez, *Parabolic Besov Regularity for the Heat Equation*, Constr. Approx. **36** (2012), no. 1, 145–159.
- [2] Petru A. Cioica, Stephan Dahlke, Nicolas Döhring, Ulrich Friedrich, Felix Lindner, Thorsten Raasch, Klaus Ritter, and René L. Schilling, *On the convergence analysis of Rothe's method*, DFG-SPP 1324 Preprint **124** (2012). URL: <http://www.dfg-spp1324.de>, last accessed: 2012/12/03.
- [3] Petru A. Cioica, Stephan Dahlke, Nicolas Döhring, Stefan Kinzel, Felix Lindner, Thorsten Raasch, Klaus Ritter, and René L. Schilling, *Adaptive wavelet methods for the stochastic Poisson equation*, BIT Numer Math **52** (2012), no. 3, 589–614.
- [4] Petru A. Cioica, Stephan Dahlke, Stefan Kinzel, Felix Lindner, Thorsten Raasch, Klaus Ritter, and René L. Schilling, *Spatial Besov regularity for stochastic partial differential equations on Lipschitz domains*, Studia Math. **207** (2011), no. 3, 197–234.
- [5] Albert Cohen, *Numerical analysis of wavelet methods*, 1st ed., Studies in mathematics and its applications, vol. 32, Elsevier, Amsterdam, 2003.
- [6] Albert Cohen, Ingrid Daubechies, and Jean-Christophe Feauveau, *Biorthogonal bases of compactly supported wavelets*, Comm. Pure Appl. Math. **45** (1992), no. 5, 485–560.
- [7] Stephan Dahlke, *Besov Regularity for Second Order Elliptic Boundary Value Problems with Variable Coefficients*, Manuscripta Math. **95** (1998), no. 1, 59–77.
- [8] ———, *Besov Regularity for Elliptic Boundary Value Problems in Polygonal Domains*, Appl. Math. Lett. **12** (1999), no. 6, 31–36.
- [9] ———, *Besov regularity of edge singularities for the Poisson equation in polyhedral domains*, Numer. Linear Algebra Appl. **9** (2002), no. 6-7, 457–466.
- [10] Stephan Dahlke and Ronald A. DeVore, *Besov regularity for elliptic boundary value problems*, Commun. Partial Differ. Equations **22** (1997), no. 1-2, 1–16.
- [11] Stephan Dahlke and Winfried Sickel, *Besov Regularity for the Poisson Equation in Smooth and Polyhedral Cones*, Sobolev Spaces in Mathematics II, Applications to Partial Differential Equations (Vladimir G. Maz'ya, ed.), International Mathematical Series 9, Springer, jointly published with Tamara Rozhkovskaya Publisher, Novosibirsk, 2008.
- [12] Ingrid Daubechies, *Ten Lectures on Wavelets*, CBMS-NSF regional conference series in applied mathematics, vol. 61, Society for Industrial and Applied Mathematics, Philadelphia PA, 1992.
- [13] Ronald A. DeVore, *Nonlinear approximation*, Acta Numer. **7** (1998), 51–150.
- [14] Ronald A. DeVore and Robert C. Sharpley, *Maximal functions measuring smoothness*, Memoirs of the American Mathematical Society, vol. 293, AMS, Providence RI, 1984.
- [15] Sophie Dispa, *Intrinsic characterizations of Besov spaces on Lipschitz domains*, Math. Nachr. **260** (2003), no. 1, 21–33.

- [16] Hongjie Dong and Doyoon Kim, *L_p solvability of divergence type parabolic and elliptic systems with partially BMO coefficients*, Calculus of Variations and Partial Differential Equations **40** (2011), no. 3&4, 357–389.
- [17] Klaus-Jochen Engel and Rainer Nagel, *One-Parameter Semigroups for Linear Evolution Equations*, Graduate Texts in Mathematics, vol. 194, Springer, New York, 2000.
- [18] Piotr Hajlasz, *Change of variables formula under minimal assumptions*, Colloq. Math. **64** (1993), no. 1, 93–101.
- [19] David Jerison and Carlos E. Kenig, *The Inhomogeneous Dirichlet Problem in Lipschitz Domains*, J. Funct. Anal. **130** (1995), no. 1, 161–219.
- [20] Hans Johnen and Karl Scherer, *On the equivalence of the K -functional and moduli of continuity and some applications*, Constructive Theory of Functions of Several Variables (Walter Schempp and Karl Zeller, eds.), Lecture Notes in Mathematics, vol. 571, Springer, 1977, pp. 119–140.
- [21] Alf Jonsson and Hans Wallin, *Function Spaces on Subsets of \mathbb{R}^n* , Mathematical Reports, vol. 2, pt. 1, Harwood Academic Publishers, Chur, 1984.
- [22] Nigel J. Kalton and Lutz Weis, *The H^∞ -calculus and sums of closed operators*, Math. Ann. **321** (2001), no. 2, 319–345.
- [23] Kyeong-Hun Kim, *On stochastic partial differential equations with variable coefficients in C^1 domains*, Stochastic Processes Appl. **112** (2004), no. 2, 261–283.
- [24] ———, *An L_p -Theory of SPDEs on Lipschitz Domains*, Potential Anal. **29** (2008), no. 3, 303–326.
- [25] ———, *Sobolev space theory of SPDEs with continuous or measurable leading coefficients*, Stochastic Processes Appl. **119** (2009), no. 1, 1644.
- [26] ———, *A weighted Sobolev space theory of parabolic stochastic PDEs on non-smooth domains*, Preprint (2011), arXiv:1109.4727.
- [27] Kyeong-Hun Kim and Nicolai V. Krylov, *On The Sobolev Space Theory of Parabolic and Elliptic Equations in C^1 Domains*, SIAM J. Math. Anal. **36** (2004), no. 2, 618–642.
- [28] Nicolai V. Krylov, *A W_2^n -theory of the Dirichlet problem for SPDEs in general smooth domains*, Probab. Theory Relat. Fields **98** (1994), no. 3, 389–421.
- [29] ———, *An analytic approach to SPDEs*, Stochastic Partial Differential Equations: Six Perspectives (René A. Carmona and Boris L. Rozovskii, eds.), Math. Surv. Monogr., vol. 64, AMS, Providence RI, 1999, pp. 185–242.
- [30] ———, *Some Properties of Weighted Sobolev Spaces in \mathbb{R}_+^d* , Ann. Sc. Norm. Super. Pisa, Cl. Sci., IV. Ser. **28** (1999), no. 4, 675–693.
- [31] ———, *SPDEs in $L_q((0, \tau], L_p)$ spaces*, Electron. J. Probab. **5** (2000), no. 13, 1–29.

- [32] ———, *Some Properties of Traces for Stochastic and Deterministic Parabolic Weighted Sobolev Spaces*, J. Funct. Anal. **183** (2001), no. 1, 1–41.
- [33] Nicolai V. Krylov and Sergey V. Lototsky, *A Sobolev Space Theory of SPDE with Constant Coefficients on a Half Line*, SIAM J. Math. Anal. **30** (1999), no. 2, 298–325.
- [34] ———, *A Sobolev Space Theory of SPDEs with Constant Coefficients in a Half Space*, SIAM J. Math. Anal. **31** (1999), no. 1, 19–33.
- [35] Alois Kufner, *Weighted Sobolev Spaces*, Teubner-Texte zur Mathematik, vol. 31, BSB B. G. Teubner Verlagsgesellschaft, Leipzig, 1980.
- [36] George C. Kyriazis, *Wavelet Coefficients Measuring Smoothness in $H_p(\mathbb{R}^d)$* , Appl. Comput. Harmon. Anal. **3** (1996), no. 2, 100–119.
- [37] Felix Lindner, *Approximation and Regularity of Stochastic PDEs*, Shaker, 2011, Doctoral Thesis.
- [38] Sergey V. Lototsky, *Sobolev Spaces with Weights in Domains and Boundary Value Problems for Degenerate Elliptic Equations*, Methods Appl. Anal. **7** (2000), no. 1, 195–204.
- [39] Vladimir G. Maz’ya, *Sobolev Spaces. With applications to elliptic partial differential equations. Translated from the Russian by T.O. Shaposhnikova*, 2 ed., Grundlehren der mathematischen Wissenschaften, vol. 342, Springer, 2011.
- [40] Yves Meyer, *Wavelets and Operators. Translated by D.H. Salinger*, Cambridge Studies in Advanced Mathematics, vol. 37, Cambridge University Press, 1992.
- [41] Amnon Pazy, *Semigroups of Linear Operators and Applications to Partial Differential Equations*, Applied Mathematical Sciences, vol. 44, Springer, New York, 1983.
- [42] Vyacheslav S. Rychkov, *On Restrictions and Extensions of the Besov and Triebel-Lizorkin Spaces with Respect to Lipschitz Domains*, J. London Math. Soc. **60** (1999), no. 1, 237–257.
- [43] Hans Triebel, *Interpolation theory, function spaces, differential operators*, 2nd ed., Barth, 1995.
- [44] ———, *Theory of Function Spaces III*, Monographs in Mathematics, vol. 100, Birkhäuser, Basel-Boston-Berlin, 2006.
- [45] Jan van Neerven, *γ -Radonifying Operators – A Survey*, The AMSI-ANU Workshop on Spectral Theory and Harmonic Analysis (Andrew Hassel, Alan McIntosh, and Robert Taggart, eds.), Proceedings of the Centre for Mathematics and its Applications, vol. 44, 2010, pp. 1–62.
- [46] Jan van Neerven, Mark Christiaan Veraar, and Lutz Weis, *Stochastic integration in UMD Banach spaces*, Ann. Probab. **35** (2007), no. 4, 1438–1478.
- [47] ———, *Maximal L^p -Regularity for Stochastic Evolution Equations*, SIAM J. Math. Anal. **44** (2012), no. 3, 1372–1414.
- [48] ———, *Stochastic maximal L^p -regularity*, Ann. Probab. **40** (2012), no. 2, 788–812.
- [49] Ian Wood, *Maximal L_p -regularity for the Laplacian on Lipschitz domains*, Math. Z. **255** (2007), no. 4, 855–875.

Affiliations & Postal Addresses

Petru A. Cioica
Philipps-Universität Marburg
FB Mathematik und Informatik, AG Numerik/Optimierung
Hans-Meerwein-Strasse
35032 Marburg, Germany
Phone: 00 49 6421 28 25 481
E-mail: cioica@mathematik.uni-marburg.de

Kyeong-Hun Kim
Korea University
Department of Mathematics
Seoul, South Korea, 136-701
Phone: 00 82 2 3290 3071
E-mail: kyeonghun@korea.ac.kr

Kijung Lee
Ajou University
Department of Mathematics
Suwon, South Korea, 443-749
Phone: 00 82 31 219 1936
E-mail: kijung@ajou.ac.kr

Felix Lindner
TU Dresden
FR Mathematik, Institut für Mathematische Stochastik
01062 Dresden, Germany
Phone: 00 49 351 463 32 437
E-mail: felix.lindner@tu-dresden.de

Preprint Series DFG-SPP 1324

<http://www.dfg-spp1324.de>

Reports

- [1] R. Ramlau, G. Teschke, and M. Zhariy. A Compressive Landweber Iteration for Solving Ill-Posed Inverse Problems. Preprint 1, DFG-SPP 1324, September 2008.
- [2] G. Plonka. The Easy Path Wavelet Transform: A New Adaptive Wavelet Transform for Sparse Representation of Two-dimensional Data. Preprint 2, DFG-SPP 1324, September 2008.
- [3] E. Novak and H. Woźniakowski. Optimal Order of Convergence and (In-) Tractability of Multivariate Approximation of Smooth Functions. Preprint 3, DFG-SPP 1324, October 2008.
- [4] M. Espig, L. Grasedyck, and W. Hackbusch. Black Box Low Tensor Rank Approximation Using Fibre-Crosses. Preprint 4, DFG-SPP 1324, October 2008.
- [5] T. Bonesky, S. Dahlke, P. Maass, and T. Raasch. Adaptive Wavelet Methods and Sparsity Reconstruction for Inverse Heat Conduction Problems. Preprint 5, DFG-SPP 1324, January 2009.
- [6] E. Novak and H. Woźniakowski. Approximation of Infinitely Differentiable Multivariate Functions Is Intractable. Preprint 6, DFG-SPP 1324, January 2009.
- [7] J. Ma and G. Plonka. A Review of Curvelets and Recent Applications. Preprint 7, DFG-SPP 1324, February 2009.
- [8] L. Denis, D. A. Lorenz, and D. Tiede. Greedy Solution of Ill-Posed Problems: Error Bounds and Exact Inversion. Preprint 8, DFG-SPP 1324, April 2009.
- [9] U. Friedrich. A Two Parameter Generalization of Lions' Nonoverlapping Domain Decomposition Method for Linear Elliptic PDEs. Preprint 9, DFG-SPP 1324, April 2009.
- [10] K. Bredies and D. A. Lorenz. Minimization of Non-smooth, Non-convex Functionals by Iterative Thresholding. Preprint 10, DFG-SPP 1324, April 2009.
- [11] K. Bredies and D. A. Lorenz. Regularization with Non-convex Separable Constraints. Preprint 11, DFG-SPP 1324, April 2009.

- [12] M. Döhler, S. Kunis, and D. Potts. Nonequispaced Hyperbolic Cross Fast Fourier Transform. Preprint 12, DFG-SPP 1324, April 2009.
- [13] C. Bender. Dual Pricing of Multi-Exercise Options under Volume Constraints. Preprint 13, DFG-SPP 1324, April 2009.
- [14] T. Müller-Gronbach and K. Ritter. Variable Subspace Sampling and Multi-level Algorithms. Preprint 14, DFG-SPP 1324, May 2009.
- [15] G. Plonka, S. Tenorth, and A. Iske. Optimally Sparse Image Representation by the Easy Path Wavelet Transform. Preprint 15, DFG-SPP 1324, May 2009.
- [16] S. Dahlke, E. Novak, and W. Sickel. Optimal Approximation of Elliptic Problems by Linear and Nonlinear Mappings IV: Errors in L_2 and Other Norms. Preprint 16, DFG-SPP 1324, June 2009.
- [17] B. Jin, T. Khan, P. Maass, and M. Pidcock. Function Spaces and Optimal Currents in Impedance Tomography. Preprint 17, DFG-SPP 1324, June 2009.
- [18] G. Plonka and J. Ma. Curvelet-Wavelet Regularized Split Bregman Iteration for Compressed Sensing. Preprint 18, DFG-SPP 1324, June 2009.
- [19] G. Teschke and C. Borries. Accelerated Projected Steepest Descent Method for Nonlinear Inverse Problems with Sparsity Constraints. Preprint 19, DFG-SPP 1324, July 2009.
- [20] L. Grasedyck. Hierarchical Singular Value Decomposition of Tensors. Preprint 20, DFG-SPP 1324, July 2009.
- [21] D. Rudolf. Error Bounds for Computing the Expectation by Markov Chain Monte Carlo. Preprint 21, DFG-SPP 1324, July 2009.
- [22] M. Hansen and W. Sickel. Best m-term Approximation and Lizorkin-Triebel Spaces. Preprint 22, DFG-SPP 1324, August 2009.
- [23] F.J. Hickernell, T. Müller-Gronbach, B. Niu, and K. Ritter. Multi-level Monte Carlo Algorithms for Infinite-dimensional Integration on \mathbb{R}^N . Preprint 23, DFG-SPP 1324, August 2009.
- [24] S. Dereich and F. Heidenreich. A Multilevel Monte Carlo Algorithm for Lévy Driven Stochastic Differential Equations. Preprint 24, DFG-SPP 1324, August 2009.
- [25] S. Dahlke, M. Fornasier, and T. Raasch. Multilevel Preconditioning for Adaptive Sparse Optimization. Preprint 25, DFG-SPP 1324, August 2009.

- [26] S. Dereich. Multilevel Monte Carlo Algorithms for Lévy-driven SDEs with Gaussian Correction. Preprint 26, DFG-SPP 1324, August 2009.
- [27] G. Plonka, S. Tenorth, and D. Roşca. A New Hybrid Method for Image Approximation using the Easy Path Wavelet Transform. Preprint 27, DFG-SPP 1324, October 2009.
- [28] O. Koch and C. Lubich. Dynamical Low-rank Approximation of Tensors. Preprint 28, DFG-SPP 1324, November 2009.
- [29] E. Faou, V. Gradinaru, and C. Lubich. Computing Semi-classical Quantum Dynamics with Hagedorn Wavepackets. Preprint 29, DFG-SPP 1324, November 2009.
- [30] D. Conte and C. Lubich. An Error Analysis of the Multi-configuration Time-dependent Hartree Method of Quantum Dynamics. Preprint 30, DFG-SPP 1324, November 2009.
- [31] C. E. Powell and E. Ullmann. Preconditioning Stochastic Galerkin Saddle Point Problems. Preprint 31, DFG-SPP 1324, November 2009.
- [32] O. G. Ernst and E. Ullmann. Stochastic Galerkin Matrices. Preprint 32, DFG-SPP 1324, November 2009.
- [33] F. Lindner and R. L. Schilling. Weak Order for the Discretization of the Stochastic Heat Equation Driven by Impulsive Noise. Preprint 33, DFG-SPP 1324, November 2009.
- [34] L. Kämmerer and S. Kunis. On the Stability of the Hyperbolic Cross Discrete Fourier Transform. Preprint 34, DFG-SPP 1324, December 2009.
- [35] P. Cerejeiras, M. Ferreira, U. Kähler, and G. Teschke. Inversion of the noisy Radon transform on $SO(3)$ by Gabor frames and sparse recovery principles. Preprint 35, DFG-SPP 1324, January 2010.
- [36] T. Jahnke and T. Udrescu. Solving Chemical Master Equations by Adaptive Wavelet Compression. Preprint 36, DFG-SPP 1324, January 2010.
- [37] P. Kittipoom, G. Kutyniok, and W.-Q Lim. Irregular Shearlet Frames: Geometry and Approximation Properties. Preprint 37, DFG-SPP 1324, February 2010.
- [38] G. Kutyniok and W.-Q Lim. Compactly Supported Shearlets are Optimally Sparse. Preprint 38, DFG-SPP 1324, February 2010.

- [39] M. Hansen and W. Sickel. Best m -Term Approximation and Tensor Products of Sobolev and Besov Spaces – the Case of Non-compact Embeddings. Preprint 39, DFG-SPP 1324, March 2010.
- [40] B. Niu, F.J. Hickernell, T. Müller-Gronbach, and K. Ritter. Deterministic Multi-level Algorithms for Infinite-dimensional Integration on $\mathbb{R}^{\mathbb{N}}$. Preprint 40, DFG-SPP 1324, March 2010.
- [41] P. Kittipoom, G. Kutyniok, and W.-Q Lim. Construction of Compactly Supported Shearlet Frames. Preprint 41, DFG-SPP 1324, March 2010.
- [42] C. Bender and J. Steiner. Error Criteria for Numerical Solutions of Backward SDEs. Preprint 42, DFG-SPP 1324, April 2010.
- [43] L. Grasedyck. Polynomial Approximation in Hierarchical Tucker Format by Vector-Tensorization. Preprint 43, DFG-SPP 1324, April 2010.
- [44] M. Hansen und W. Sickel. Best m -Term Approximation and Sobolev-Besov Spaces of Dominating Mixed Smoothness - the Case of Compact Embeddings. Preprint 44, DFG-SPP 1324, April 2010.
- [45] P. Binev, W. Dahmen, and P. Lamby. Fast High-Dimensional Approximation with Sparse Occupancy Trees. Preprint 45, DFG-SPP 1324, May 2010.
- [46] J. Ballani and L. Grasedyck. A Projection Method to Solve Linear Systems in Tensor Format. Preprint 46, DFG-SPP 1324, May 2010.
- [47] P. Binev, A. Cohen, W. Dahmen, R. DeVore, G. Petrova, and P. Wojtaszczyk. Convergence Rates for Greedy Algorithms in Reduced Basis Methods. Preprint 47, DFG-SPP 1324, May 2010.
- [48] S. Kestler and K. Urban. Adaptive Wavelet Methods on Unbounded Domains. Preprint 48, DFG-SPP 1324, June 2010.
- [49] H. Yserentant. The Mixed Regularity of Electronic Wave Functions Multiplied by Explicit Correlation Factors. Preprint 49, DFG-SPP 1324, June 2010.
- [50] H. Yserentant. On the Complexity of the Electronic Schrödinger Equation. Preprint 50, DFG-SPP 1324, June 2010.
- [51] M. Guillemard and A. Iske. Curvature Analysis of Frequency Modulated Manifolds in Dimensionality Reduction. Preprint 51, DFG-SPP 1324, June 2010.
- [52] E. Herrholz and G. Teschke. Compressive Sensing Principles and Iterative Sparse Recovery for Inverse and Ill-Posed Problems. Preprint 52, DFG-SPP 1324, July 2010.

- [53] L. Kämmerer, S. Kunis, and D. Potts. Interpolation Lattices for Hyperbolic Cross Trigonometric Polynomials. Preprint 53, DFG-SPP 1324, July 2010.
- [54] G. Kutyniok and W.-Q Lim. Shearlets on Bounded Domains. Preprint 54, DFG-SPP 1324, July 2010.
- [55] A. Zeiser. Wavelet Approximation in Weighted Sobolev Spaces of Mixed Order with Applications to the Electronic Schrödinger Equation. Preprint 55, DFG-SPP 1324, July 2010.
- [56] G. Kutyniok, J. Lemvig, and W.-Q Lim. Compactly Supported Shearlets. Preprint 56, DFG-SPP 1324, July 2010.
- [57] A. Zeiser. On the Optimality of the Inexact Inverse Iteration Coupled with Adaptive Finite Element Methods. Preprint 57, DFG-SPP 1324, July 2010.
- [58] S. Jokar. Sparse Recovery and Kronecker Products. Preprint 58, DFG-SPP 1324, August 2010.
- [59] T. Aboiyar, E. H. Georgoulis, and A. Iske. Adaptive ADER Methods Using Kernel-Based Polyharmonic Spline WENO Reconstruction. Preprint 59, DFG-SPP 1324, August 2010.
- [60] O. G. Ernst, A. Mugler, H.-J. Starkloff, and E. Ullmann. On the Convergence of Generalized Polynomial Chaos Expansions. Preprint 60, DFG-SPP 1324, August 2010.
- [61] S. Holtz, T. Rohwedder, and R. Schneider. On Manifolds of Tensors of Fixed TT-Rank. Preprint 61, DFG-SPP 1324, September 2010.
- [62] J. Ballani, L. Grasedyck, and M. Kluge. Black Box Approximation of Tensors in Hierarchical Tucker Format. Preprint 62, DFG-SPP 1324, October 2010.
- [63] M. Hansen. On Tensor Products of Quasi-Banach Spaces. Preprint 63, DFG-SPP 1324, October 2010.
- [64] S. Dahlke, G. Steidl, and G. Teschke. Shearlet Coorbit Spaces: Compactly Supported Analyzing Shearlets, Traces and Embeddings. Preprint 64, DFG-SPP 1324, October 2010.
- [65] W. Hackbusch. Tensorisation of Vectors and their Efficient Convolution. Preprint 65, DFG-SPP 1324, November 2010.
- [66] P. A. Cioica, S. Dahlke, S. Kinzel, F. Lindner, T. Raasch, K. Ritter, and R. L. Schilling. Spatial Besov Regularity for Stochastic Partial Differential Equations on Lipschitz Domains. Preprint 66, DFG-SPP 1324, November 2010.

- [67] E. Novak and H. Woźniakowski. On the Power of Function Values for the Approximation Problem in Various Settings. Preprint 67, DFG-SPP 1324, November 2010.
- [68] A. Hinrichs, E. Novak, and H. Woźniakowski. The Curse of Dimensionality for Monotone and Convex Functions of Many Variables. Preprint 68, DFG-SPP 1324, November 2010.
- [69] G. Kutyniok and W.-Q. Lim. Image Separation Using Shearlets. Preprint 69, DFG-SPP 1324, November 2010.
- [70] B. Jin and P. Maass. An Analysis of Electrical Impedance Tomography with Applications to Tikhonov Regularization. Preprint 70, DFG-SPP 1324, December 2010.
- [71] S. Holtz, T. Rohwedder, and R. Schneider. The Alternating Linear Scheme for Tensor Optimisation in the TT Format. Preprint 71, DFG-SPP 1324, December 2010.
- [72] T. Müller-Gronbach and K. Ritter. A Local Refinement Strategy for Constructive Quantization of Scalar SDEs. Preprint 72, DFG-SPP 1324, December 2010.
- [73] T. Rohwedder and R. Schneider. An Analysis for the DIIS Acceleration Method used in Quantum Chemistry Calculations. Preprint 73, DFG-SPP 1324, December 2010.
- [74] C. Bender and J. Steiner. Least-Squares Monte Carlo for Backward SDEs. Preprint 74, DFG-SPP 1324, December 2010.
- [75] C. Bender. Primal and Dual Pricing of Multiple Exercise Options in Continuous Time. Preprint 75, DFG-SPP 1324, December 2010.
- [76] H. Harbrecht, M. Peters, and R. Schneider. On the Low-rank Approximation by the Pivoted Cholesky Decomposition. Preprint 76, DFG-SPP 1324, December 2010.
- [77] P. A. Cioica, S. Dahlke, N. Döhring, S. Kinzel, F. Lindner, T. Raasch, K. Ritter, and R. L. Schilling. Adaptive Wavelet Methods for Elliptic Stochastic Partial Differential Equations. Preprint 77, DFG-SPP 1324, January 2011.
- [78] G. Plonka, S. Tenorth, and A. Iske. Optimal Representation of Piecewise Hölder Smooth Bivariate Functions by the Easy Path Wavelet Transform. Preprint 78, DFG-SPP 1324, January 2011.

- [79] A. Mugler and H.-J. Starkloff. On Elliptic Partial Differential Equations with Random Coefficients. Preprint 79, DFG-SPP 1324, January 2011.
- [80] T. Müller-Gronbach, K. Ritter, and L. Yaroslavtseva. A Derandomization of the Euler Scheme for Scalar Stochastic Differential Equations. Preprint 80, DFG-SPP 1324, January 2011.
- [81] W. Dahmen, C. Huang, C. Schwab, and G. Welper. Adaptive Petrov-Galerkin methods for first order transport equations. Preprint 81, DFG-SPP 1324, January 2011.
- [82] K. Grella and C. Schwab. Sparse Tensor Spherical Harmonics Approximation in Radiative Transfer. Preprint 82, DFG-SPP 1324, January 2011.
- [83] D.A. Lorenz, S. Schiffler, and D. Tiede. Beyond Convergence Rates: Exact Inversion With Tikhonov Regularization With Sparsity Constraints. Preprint 83, DFG-SPP 1324, January 2011.
- [84] S. Dereich, M. Scheutzow, and R. Schottstedt. Constructive quantization: Approximation by empirical measures. Preprint 84, DFG-SPP 1324, January 2011.
- [85] S. Dahlke and W. Sickel. On Besov Regularity of Solutions to Nonlinear Elliptic Partial Differential Equations. Preprint 85, DFG-SPP 1324, January 2011.
- [86] S. Dahlke, U. Friedrich, P. Maass, T. Raasch, and R.A. Ressel. An adaptive wavelet method for parameter identification problems in parabolic partial differential equations. Preprint 86, DFG-SPP 1324, January 2011.
- [87] A. Cohen, W. Dahmen, and G. Welper. Adaptivity and Variational Stabilization for Convection-Diffusion Equations. Preprint 87, DFG-SPP 1324, January 2011.
- [88] T. Jahnke. On Reduced Models for the Chemical Master Equation. Preprint 88, DFG-SPP 1324, January 2011.
- [89] P. Binev, W. Dahmen, R. DeVore, P. Lamby, D. Savu, and R. Sharpley. Compressed Sensing and Electron Microscopy. Preprint 89, DFG-SPP 1324, March 2011.
- [90] P. Binev, F. Blanco-Silva, D. Blom, W. Dahmen, P. Lamby, R. Sharpley, and T. Vogt. High Quality Image Formation by Nonlocal Means Applied to High-Angle Annular Dark Field Scanning Transmission Electron Microscopy (HAADF-STEM). Preprint 90, DFG-SPP 1324, March 2011.
- [91] R. A. Ressel. A Parameter Identification Problem for a Nonlinear Parabolic Differential Equation. Preprint 91, DFG-SPP 1324, May 2011.

- [92] G. Kutyniok. Data Separation by Sparse Representations. Preprint 92, DFG-SPP 1324, May 2011.
- [93] M. A. Davenport, M. F. Duarte, Y. C. Eldar, and G. Kutyniok. Introduction to Compressed Sensing. Preprint 93, DFG-SPP 1324, May 2011.
- [94] H.-C. Kreuzler and H. Yserentant. The Mixed Regularity of Electronic Wave Functions in Fractional Order and Weighted Sobolev Spaces. Preprint 94, DFG-SPP 1324, June 2011.
- [95] E. Ullmann, H. C. Elman, and O. G. Ernst. Efficient Iterative Solvers for Stochastic Galerkin Discretizations of Log-Transformed Random Diffusion Problems. Preprint 95, DFG-SPP 1324, June 2011.
- [96] S. Kunis and I. Melzer. On the Butterfly Sparse Fourier Transform. Preprint 96, DFG-SPP 1324, June 2011.
- [97] T. Rohwedder. The Continuous Coupled Cluster Formulation for the Electronic Schrödinger Equation. Preprint 97, DFG-SPP 1324, June 2011.
- [98] T. Rohwedder and R. Schneider. Error Estimates for the Coupled Cluster Method. Preprint 98, DFG-SPP 1324, June 2011.
- [99] P. A. Cioica and S. Dahlke. Spatial Besov Regularity for Semilinear Stochastic Partial Differential Equations on Bounded Lipschitz Domains. Preprint 99, DFG-SPP 1324, July 2011.
- [100] L. Grasedyck and W. Hackbusch. An Introduction to Hierarchical (H-) Rank and TT-Rank of Tensors with Examples. Preprint 100, DFG-SPP 1324, August 2011.
- [101] N. Chegini, S. Dahlke, U. Friedrich, and R. Stevenson. Piecewise Tensor Product Wavelet Bases by Extensions and Approximation Rates. Preprint 101, DFG-SPP 1324, September 2011.
- [102] S. Dahlke, P. Oswald, and T. Raasch. A Note on Quarkonial Systems and Multi-level Partition of Unity Methods. Preprint 102, DFG-SPP 1324, September 2011.
- [103] A. Uschmajew. Local Convergence of the Alternating Least Squares Algorithm For Canonical Tensor Approximation. Preprint 103, DFG-SPP 1324, September 2011.
- [104] S. Kvaal. Multiconfigurational time-dependent Hartree method for describing particle loss due to absorbing boundary conditions. Preprint 104, DFG-SPP 1324, September 2011.

- [105] M. Guillemand and A. Iske. On Groupoid C^* -Algebras, Persistent Homology and Time-Frequency Analysis. Preprint 105, DFG-SPP 1324, September 2011.
- [106] A. Hinrichs, E. Novak, and H. Woźniakowski. Discontinuous information in the worst case and randomized settings. Preprint 106, DFG-SPP 1324, September 2011.
- [107] M. Espig, W. Hackbusch, A. Litvinenko, H. Matthies, and E. Zander. Efficient Analysis of High Dimensional Data in Tensor Formats. Preprint 107, DFG-SPP 1324, September 2011.
- [108] M. Espig, W. Hackbusch, S. Handschuh, and R. Schneider. Optimization Problems in Contracted Tensor Networks. Preprint 108, DFG-SPP 1324, October 2011.
- [109] S. Dereich, T. Müller-Gronbach, and K. Ritter. On the Complexity of Computing Quadrature Formulas for SDEs. Preprint 109, DFG-SPP 1324, October 2011.
- [110] D. Belomestny. Solving optimal stopping problems by empirical dual optimization and penalization. Preprint 110, DFG-SPP 1324, November 2011.
- [111] D. Belomestny and J. Schoenmakers. Multilevel dual approach for pricing American style derivatives. Preprint 111, DFG-SPP 1324, November 2011.
- [112] T. Rohwedder and A. Uschmajew. Local convergence of alternating schemes for optimization of convex problems in the TT format. Preprint 112, DFG-SPP 1324, December 2011.
- [113] T. Görner, R. Hielscher, and S. Kunis. Efficient and accurate computation of spherical mean values at scattered center points. Preprint 113, DFG-SPP 1324, December 2011.
- [114] Y. Dong, T. Görner, and S. Kunis. An iterative reconstruction scheme for photoacoustic imaging. Preprint 114, DFG-SPP 1324, December 2011.
- [115] L. Kämmerer. Reconstructing hyperbolic cross trigonometric polynomials by sampling along generated sets. Preprint 115, DFG-SPP 1324, February 2012.
- [116] H. Chen and R. Schneider. Numerical analysis of augmented plane waves methods for full-potential electronic structure calculations. Preprint 116, DFG-SPP 1324, February 2012.
- [117] J. Ma, G. Plonka, and M.Y. Hussaini. Compressive Video Sampling with Approximate Message Passing Decoding. Preprint 117, DFG-SPP 1324, February 2012.

- [118] D. Heinen and G. Plonka. Wavelet shrinkage on paths for scattered data denoising. Preprint 118, DFG-SPP 1324, February 2012.
- [119] T. Jahnke and M. Kreim. Error bound for piecewise deterministic processes modeling stochastic reaction systems. Preprint 119, DFG-SPP 1324, March 2012.
- [120] C. Bender and J. Steiner. A-posteriori estimates for backward SDEs. Preprint 120, DFG-SPP 1324, April 2012.
- [121] M. Espig, W. Hackbusch, A. Litvinenkoy, H.G. Matthiesy, and P. Wähnert. Efficient low-rank approximation of the stochastic Galerkin matrix in tensor formats. Preprint 121, DFG-SPP 1324, May 2012.
- [122] O. Bokanowski, J. Garcke, M. Griebel, and I. Klompaker. An adaptive sparse grid semi-Lagrangian scheme for first order Hamilton-Jacobi Bellman equations. Preprint 122, DFG-SPP 1324, June 2012.
- [123] A. Mugler and H.-J. Starkloff. On the convergence of the stochastic Galerkin method for random elliptic partial differential equations. Preprint 123, DFG-SPP 1324, June 2012.
- [124] P.A. Cioica, S. Dahlke, N. Döhring, U. Friedrich, S. Kinzel, F. Lindner, T. Raasch, K. Ritter, and R.L. Schilling. On the convergence analysis of Rothe’s method. Preprint 124, DFG-SPP 1324, July 2012.
- [125] P. Binev, A. Cohen, W. Dahmen, and R. DeVore. Classification Algorithms using Adaptive Partitioning. Preprint 125, DFG-SPP 1324, July 2012.
- [126] C. Lubich, T. Rohwedder, R. Schneider, and B. Vandereycken. Dynamical approximation of hierarchical Tucker and Tensor-Train tensors. Preprint 126, DFG-SPP 1324, July 2012.
- [127] M. Kovács, S. Larsson, and K. Urban. On Wavelet-Galerkin methods for semilinear parabolic equations with additive noise. Preprint 127, DFG-SPP 1324, August 2012.
- [128] M. Bachmayr, H. Chen, and R. Schneider. Numerical analysis of Gaussian approximations in quantum chemistry. Preprint 128, DFG-SPP 1324, August 2012.
- [129] D. Rudolf. Explicit error bounds for Markov chain Monte Carlo. Preprint 129, DFG-SPP 1324, August 2012.
- [130] P.A. Cioica, K.-H. Kim, K. Lee, and F. Lindner. On the $L_q(L_p)$ -regularity and Besov smoothness of stochastic parabolic equations on bounded Lipschitz domains. Preprint 130, DFG-SPP 1324, December 2012.