

DFG-Schwerpunktprogramm 1324

„Extraktion quantifizierbarer Information aus komplexen Systemen“

Adaptive Wavelet Schwarz Methods for the Navier-Stokes Equation

Stephan Dahlke, Dominik Lellek, Shiu Hong Lui, Rob Stevenson

Preprint 140



Edited by

AG Numerik/Optimierung
Fachbereich 12 - Mathematik und Informatik
Philipps-Universität Marburg
Hans-Meerwein-Str.
35032 Marburg

DFG-Schwerpunktprogramm 1324

„Extraktion quantifizierbarer Information aus komplexen Systemen“

Adaptive Wavelet Schwarz Methods for the Navier-Stokes Equation

Stephan Dahlke, Dominik Lellek, Shiu Hong Lui, Rob Stevenson

Preprint 140



The consecutive numbering of the publications is determined by their chronological order.

The aim of this preprint series is to make new research rapidly available for scientific discussion. Therefore, the responsibility for the contents is solely due to the authors. The publications will be distributed by the authors.

Adaptive Wavelet Schwarz Methods for the Navier-Stokes Equation *

Stephan Dahlke, Dominik Lellek, Shiu Hong Lui, Rob Stevenson

May 22, 2013

Abstract

In this paper we are concerned with domain decomposition methods for the stationary incompressible Navier-Stokes equation. We construct an adaptive additive Schwarz method based on discretization by means of a divergence-free wavelet frame. We prove that the method is convergent and asymptotically optimal with respect to the degrees of freedom involved.

1 Introduction

Over the last years, adaptive wavelet methods for both linear and nonlinear partial differential equations have intensively been investigated, see, for instance, [1, 2]. One can often prove that these methods are not only convergent, but also asymptotically optimal. This means that the algorithm converges with the same rate as the best N -term wavelet approximation with respect to the degrees of freedom involved. The techniques used to show these results heavily rely on the properties of the underlying wavelet Riesz basis. This basis can be constructed such that its elements have vanishing moments, are piecewise smooth and characterize function spaces in the sense that weighted sequence norms of wavelet expansion coefficients are equivalent to smoothness norms such as Besov norms. Moreover, it is also possible to construct divergence-free wavelet bases, see [3, 4] which are very useful for the numerical treatment of incompressible flow problems, see [5, 6].

However, on more complicated domains, the design of such a wavelet basis becomes increasingly difficult and the condition numbers become worse. A way to facilitate the construction is to use redundant generating systems, namely wavelet frames, instead of bases, see [7]. To do so, let us assume that we can decompose the domain into overlapping subdomains that are affine images of the unit cube. Then, we can construct wavelet bases on each of the subdomains, which is significantly easier, and simply collect these bases. From this, we obtain a wavelet frame.

Because the construction of the wavelet frame is based on an overlapping domain decomposition, it is natural to combine wavelet methods with domain decomposition solvers such as Schwarz methods. For some early work on domain decomposition methods for nonlinear problems, we refer to [8, 9]. The combination with wavelet methods has proven to be very effective for the numerical solution of linear elliptic problems, see [10], and has recently been generalized to a range of nonlinear problems, see [11]. Based

*This work has been supported by Deutsche Forschungsgemeinschaft (DFG), Grant DA360/12-2.

on an idea from [12], in this paper we are going to extend this approach to the stationary, incompressible Navier-Stokes equation. We are going to show that the method is convergent and asymptotically optimal, at least for sufficiently small Reynolds numbers.

In this work, by $C \lesssim D$ we will mean that C can be bounded by a multiple of D , independently of parameters which C and D may depend on. Obviously, $C \gtrsim D$ is defined as $D \lesssim C$, and $C \approx D$ as $C \lesssim D$ and $C \gtrsim D$.

2 The Navier-Stokes equations in frame coordinates

2.1 Navier-Stokes equations

We are concerned with the incompressible, steady-state, viscous Navier-Stokes equation on a bounded Lipschitz domain $\Omega \subset \mathbb{R}^d$, $d \leq 4$, with Dirichlet boundary conditions

$$\begin{aligned} (u \cdot \nabla)u &= -\nabla p + \frac{1}{\text{Re}}\Delta u + f && \text{on } \Omega, \\ \text{div } u &= 0 && \text{on } \Omega, \\ u &= 0 && \text{on } \partial\Omega, \end{aligned}$$

where u denotes the velocity field of a fluid, p is the pressure term, f is the given inertial force and Re is the Reynolds number that describes the viscosity of the fluid. In addition, we normalize the pressure term p by $\int_{\Omega} p = 0$.

There are basically two general approaches for the numerical treatment of this equation. One common approach is to solve for the velocity u and the pressure p simultaneously. Doing so leads to an indefinite saddle point problem, see, for instance, [13] for an overview in the context of finite element methods or [14] for a wavelet-based method.

An alternative approach is to reformulate the equation using a divergence-free ansatz space

$$V := \{v \in (H_0^1(\Omega))^d, \text{div } v = 0\}.$$

The Leray weak formulation of original problem then reduces to finding a $u \in V$ such that

$$\int_{\Omega} v(u \cdot \nabla)u = -\frac{1}{\text{Re}}a(u, v) + \int_{\Omega} f \cdot v \quad (v \in V), \quad (1)$$

with $a(u, v) := \sum_{k=1}^d \int_{\Omega} \nabla u_k \nabla v_k$ or, equivalently,

$$a(u, v) + \text{Re} \int_{\Omega} v(u \cdot \nabla)u = \text{Re} \int_{\Omega} f \cdot v \quad (v \in V).$$

Note that in this formulation, the pressure term drops out, and we solve for the velocity field u only. For details of the derivation of the weak formulation, see, for instance, [15]. The formulation there coincides with (1) for our case $d \in \{2, 3, 4\}$. In particular, existence of a weak solution u is shown there. To guarantee uniqueness, it has to be assumed that the Reynolds number is sufficiently small or that the data f fulfills a smallness condition. We will equip V with energy norm

$$\|v\| = a(v, v)^{\frac{1}{2}}.$$

In order to write the equation (1) as an infinite system of scalar equations, we need a generating system for the space V . Hence, in the next subsections, we outline the construction of a divergence-free wavelet frame for this space.

2.2 Frames

Recall that a countable collection $\Psi = \{\psi_\lambda : \lambda \in \Lambda\}$ in a Hilbert space V is called a *frame* for V when there exist two positive constants A_Ψ, B_Ψ such that

$$A_\Psi \|f\|_{V'} \leq \|(f(\psi_\lambda))_{\lambda \in \Lambda}\|_{\ell_2(\Lambda)} \leq B_\Psi \|f\|_{V'} \quad (f \in V'). \quad (2)$$

As a consequence of (2), the frame operator

$$F : V' \rightarrow \ell_2(\Lambda) : f \mapsto f(\Psi) := (f(\psi_\lambda))_{\lambda \in \Lambda},$$

and so its adjoint

$$F' : \ell_2(\Lambda) \rightarrow V : \mathbf{c} \mapsto \mathbf{c}^\top \Psi := \sum_{\lambda \in \Lambda} c_\lambda \psi_\lambda,$$

are bounded with norm less than or equal to B_Ψ . The composition $F'F : V' \rightarrow V$ is boundedly invertible with $\|(F'F)^{-1}\|_{V \rightarrow V'} \leq A_\Psi^{-2}$. The collection $\tilde{\Psi} := (F'F)^{-1}\Psi$ is a frame for V' with frame operators

$$\tilde{F} := F(F'F)^{-1}, \quad \tilde{F}' = (F'F)^{-1}F'$$

and frame constants $B_{\tilde{\Psi}}^{-1}, A_{\tilde{\Psi}}^{-1}$. Since $F'\tilde{F} = \text{Id} = \tilde{F}'F$, $\tilde{\Psi}$ is called a dual frame for Ψ , known as the canonical dual frame. We have $\ell_2(\Lambda) = \text{ran } F \oplus^\perp \ker F'$, and $F(F'F)^{-1}F'$ is the orthogonal projector onto $\text{ran } F$. For these facts and further reading on frames, we refer to [16].

The key to the construction of a frame for a space of functions on a domain by means of an overlapping domain decomposition is the following lemma:

Lemma 2.1 *The property of a countable $\Psi \subset V$ being a frame for V with constants A_Ψ, B_Ψ is equivalent to $\overline{\text{span } \Psi} = V$ and*

$$B_\Psi^{-1} \|u\|_V \leq \inf_{\{\mathbf{u} \in \ell_2(\Lambda), \mathbf{u}^\top \Psi = u\}} \|\mathbf{u}\|_{\ell_2(\Lambda)} \leq A_\Psi^{-1} \|u\|_V \quad (u \in V). \quad (3)$$

Proof. If Ψ is a frame, then $(\tilde{\psi}_\lambda(u))_{\lambda \in \Lambda} = \arg \min\{\|\mathbf{u}\|_{\ell_2(\Lambda)} : \mathbf{u} \in \ell_2(\Lambda), \mathbf{u}^\top \Psi = u\}$, and (3) follows from $\tilde{\Psi}$ being a frame with frame constants $B_{\tilde{\Psi}}^{-1}, A_{\tilde{\Psi}}^{-1}$.

Conversely, let (3) be valid. Then from its first inequality, we deduce that $\mathbf{c} \mapsto \mathbf{c}^\top \Psi \in B(\ell_2(\Lambda), V)$, with norm less or equal to B_Ψ . From its second inequality, we infer that for $v \in V$, $\sup_{0 \neq \mathbf{d} \in \ell_2(\Lambda)} \frac{\langle \mathbf{d}^\top \Psi, v \rangle_V}{\|\mathbf{d}\|_{\ell_2(\Lambda)} \|v\|_V} \geq \frac{\|v\|_V}{\inf_{\{\mathbf{d} \in \ell_2(\Lambda), \mathbf{d}^\top \Psi = v\}} \|\mathbf{d}\|_{\ell_2(\Lambda)}} \geq A_\Psi$. Consequently, for given $u \in V$, there exists a unique solution $(\hat{\mathbf{u}}, w) \in \ell_2(\Lambda) \times V$ of the linear problem

$$\begin{cases} \langle \hat{\mathbf{u}}, \mathbf{d} \rangle_{\ell_2(\Lambda)} + \langle \mathbf{d}^\top \Psi, w \rangle_V = 0 & (\mathbf{d} \in \ell_2(\Lambda)), \\ \langle \hat{\mathbf{u}}^\top \Psi, v \rangle_V = \langle u, v \rangle_V & (v \in V), \end{cases} \quad (4)$$

and $\hat{\mathbf{u}} = \arg \min\{\|\mathbf{u}\|_{\ell_2(\Lambda)} : \mathbf{u} \in \ell_2(\Lambda), \mathbf{u}^\top \Psi = u\}$. Defining $\tilde{\psi}_\lambda : u \mapsto \hat{\mathbf{u}}_\lambda \in V'$, (3) means that $\tilde{\Psi}$ is a frame for V' with frame constants $B_{\tilde{\Psi}}^{-1}, A_{\tilde{\Psi}}^{-1}$.

Next, we consider (4) for $u = \psi_\mu$, so that $\hat{\mathbf{u}}_\lambda = \tilde{\psi}_\lambda(\psi_\mu)$. With $R : \ell_2(\Lambda) \rightarrow \ell_2(\Lambda)'$, $B : V \rightarrow \ell_2(\Lambda)'$ being defined by $(R\mathbf{c})(\mathbf{d}) = \langle \mathbf{c}, \mathbf{d} \rangle_{\ell_2(\Lambda)}$, $(Bw)(\mathbf{d}) = \langle \mathbf{d}^\top \Psi, w \rangle_V$, (4) with $u = \psi_\mu$ reads as

$$\begin{bmatrix} R & B \\ B' & 0 \end{bmatrix} \begin{bmatrix} \hat{\mathbf{u}} \\ w \end{bmatrix} = \begin{bmatrix} 0 \\ B'\mathbf{e}_\mu \end{bmatrix}$$

and so $\hat{\mathbf{u}} = R^{-1}B(B'R^{-1}B)^{-1}B'\mathbf{e}_\mu$. We conclude that

$$\tilde{\psi}_\lambda(\psi_\mu) = \hat{\mathbf{u}}_\lambda = (R\hat{\mathbf{u}})(\mathbf{e}_\lambda) = (B(B'R^{-1}B)^{-1}B'\mathbf{e}_\mu)(\mathbf{e}_\lambda) = (B(B'R^{-1}B)^{-1}B'\mathbf{e}_\lambda)(\mathbf{e}_\mu) = \tilde{\psi}_\mu(\psi_\lambda). \quad (5)$$

The second equation in (4) shows that $\sum_{\lambda \in \Lambda} \tilde{\psi}_\lambda(u)\psi_\lambda = u$ ($u \in V$), and so $\sum_{\lambda \in \Lambda} \tilde{\psi}_\lambda(u)\tilde{\psi}_\mu(\psi_\lambda) = \tilde{\psi}_\mu(u)$ ($u \in H$, $\mu \in \Lambda$), or $\sum_{\lambda \in \Lambda} \tilde{\psi}_\mu(\psi_\lambda)\tilde{\psi}_\lambda = \tilde{\psi}_\mu$. Replacing $\tilde{\psi}_\mu(\psi_\lambda)$ by $\tilde{\psi}_\lambda(\psi_\mu)$ in the last equality because of (5), it reads as $\tilde{F}'\tilde{F}\psi_\mu = \tilde{\psi}_\mu$, with \tilde{F} being the frame operator of $\tilde{\Psi}$. We conclude that Ψ is the canonical dual frame of $\tilde{\Psi}$, and thus in particular a frame, and which therefore has frame constants A_Ψ, B_Ψ . \square

2.3 Domain decomposition

A wavelet frame will be obtained by decomposing the domain Ω into affine overlapping images of the unit cube, $\Omega = \bigcup_{i=0}^{m-1} \Omega_i$. Let us assume that such a decomposition exists and that we have wavelet bases $\Psi_i = \{\psi_\lambda^{(i)} : \lambda \in \Lambda_i\}$ for the spaces

$$V_i := \{v \in (H_0^1(\Omega_i))^d, \operatorname{div} v = 0\}.$$

Having these bases at hand, we simply set $\Psi := \bigcup_{i=0}^{m-1} E_i\Psi_i$, where E_i is the zero extension from V_i to V . The index set belonging to Ψ is denoted by $\Lambda := \bigcup_{i=0}^{m-1} \{i\} \times \Lambda_i$, so we can write $\Psi = \{\psi_\lambda : \lambda \in \Lambda\}$. If the subdomains are overlapping, we can show that we indeed obtain a wavelet frame for V .

Lemma 2.2 *Assume the subdomains Ω_i are overlapping in the sense that*

$$H_0^1(\Omega) = H_0^1(\Omega_0) + \dots + H_0^1(\Omega_{m-1}).$$

Let Ψ_i be frames or Riesz bases for V_i . Then, $\Psi := \bigcup_{i=0}^{m-1} E_i\Psi_i$ is a frame for V .

Proof. We clearly have $H_0^1(\Omega)^d = H_0^1(\Omega_0)^d + \dots + H_0^1(\Omega_{m-1})^d$. Hence, from [12, Lemma 2], it follows that even $V = V_0 + \dots + V_{m-1}$. Now, from the partition lemma (see, for instance, [17]) we may conclude that there exists a stable splitting of V , which means that, uniformly in $v \in V$, we have

$$\begin{aligned} \|v\|_{H^1(\Omega)^d}^2 &\approx \inf_{\{(v_i)_i \in \prod_{i=0}^{m-1} V_i : v = \sum_{i=0}^{m-1} E_i v_i\}} \sum_{i=0}^{m-1} \|v_i\|_{H^1(\Omega_i)^d}^2 \\ &\approx \inf_{\{(v_i)_i \in \prod_{i=0}^{m-1} V_i : v = \sum_{i=0}^{m-1} E_i v_i\}} \sum_{i=0}^{m-1} \inf_{\{\mathbf{v}_i \in \ell_2(\Lambda_i) : \mathbf{v}_i^\top \Psi_i = v_i\}} \|\mathbf{v}_i\|_{\ell_2(\Lambda_i)}^2 \\ &\approx \inf_{\{\mathbf{v} \in \ell_2(\Lambda) : \mathbf{v}^\top \Psi = v\}} \|\mathbf{v}\|_{\ell_2(\Lambda)}^2, \end{aligned}$$

where we used Lemma 2.1 for the second \approx . The proof is completed by another application of this lemma. \square

Remark 2.3 *The condition in Lemma 2.2 is fulfilled if there exists a smooth partition of unity with respect to the domain decomposition $\Omega = \bigcup_{i=0}^{m-1} \Omega_i$. However, it can also be shown for situations where such a partition does not exist. One important case of this kind is the prototype of a non-convex polygonal domain in two space dimensions, the L-shaped domain $\Omega = (-1, 1)^2 \setminus [0, 1)^2$ with subdomains $\Omega_0 = (-1, 1) \times (-1, 0)$ and $\Omega_1 = (-1, 0) \times (-1, 1)$, see [17, 18].*

2.4 Divergence-free wavelet bases on the subdomains

We consider subdomains that are hypercubes. More general subdomains can then be treated by applying the Piola transform. Furthermore, we restrict ourselves to the two-dimensional case, i.e., to the unit square \mathcal{I}^2 , where $\mathcal{I} := (0, 1)$. Divergence-free wavelet bases for $\{v \in H_0^1(\mathcal{I}^d) : \operatorname{div} v = 0\}$ were constructed in [4] for any dimension $d \geq 2$. For $d > 2$, these bases consist of *anisotropic* wavelets, i.e., vectors of tensor products of univariate wavelets on arbitrary, unrelated scales. These anisotropic wavelet bases have the advantage that they give rise to approximation rates that are independent of the space dimension. On the other hand, the efficient approximate evaluation of nonlinear terms in anisotropic wavelet coordinates is yet not well understood. For that reason here we recall the construction of the *isotropic* divergence-free wavelet basis from [4] that applies to $d = 2$, cf. also [19].

The construction starts with collections of univariate primal and dual wavelets, and, for $\ell \in \mathbb{N}_0$, collections of univariate primal and dual scaling functions

$$\Psi = \{\psi_\lambda : \lambda \in J\}, \quad \tilde{\Psi} = \{\tilde{\psi}_\lambda : \lambda \in J\}, \quad \Phi_\ell = \{\phi_{\ell,k} : 1 \leq k \leq N_\ell\}, \quad \tilde{\Phi}_\ell = \{\tilde{\phi}_{\ell,k} : 1 \leq k \leq N_\ell\},$$

such that

- (a). $(\Psi, \tilde{\Psi})$ are $L_2(\mathcal{I})$ -biorthogonal,
- (b). $\{2^{-|\lambda|} \tilde{\psi}_\lambda : \lambda \in J\}$ is a Riesz basis for $H^1(\mathcal{I})$, where $|\lambda| \in \mathbb{N}_0$ is referred to as the level of λ ,
- (c). $\{2^{-|\lambda|} \psi_\lambda : \lambda \in J\}$ is a Riesz basis for $H_0^1(\mathcal{I})$,
- (d). Ψ is local, meaning that both $\operatorname{diam} \operatorname{supp} \psi_\lambda \lesssim 2^{-|\lambda|}$, and each interval of length $2^{-\ell}$ intersects the supports of an at most uniformly bounded number of ψ_λ for $|\lambda| = \ell$; and $\tilde{\Psi}$ is local,
- (e). there is a $\underline{\lambda} \in J$ with $|\underline{\lambda}| = 0$, such that $\tilde{\psi}_{\underline{\lambda}}$ is a multiple of the constant function $\mathbb{1}$,
- (f). $\operatorname{span}\{\psi_\lambda : \lambda \in J, |\lambda| \leq \ell\} = \operatorname{span} \Phi_\ell$, $\operatorname{span}\{\tilde{\psi}_\lambda : \lambda \in J, |\lambda| \leq \ell\} = \operatorname{span} \tilde{\Phi}_\ell$,
- (g). Φ_ℓ and $\tilde{\Phi}_\ell$ are biorthogonal, uniform (in ℓ) $L_2(\mathcal{I})$ -Riesz bases for their spans,
- (h). Φ_ℓ is uniformly (in ℓ) local, meaning that both $\operatorname{diam} \operatorname{supp} \phi_{\ell,k} \lesssim 2^{-\ell}$, and each interval of length $2^{-\ell}$ intersects the supports of an at most uniformly bounded number of $\phi_{\ell,k}$ for $|\lambda| = \ell$; and $\tilde{\Phi}$ is uniformly local.
- (i). for each ℓ , $\int_{\mathcal{I}} \phi_{\ell,k}$ is independent of $1 \leq k \leq N_\ell$, and $\inf \operatorname{supp} \phi_{k,\ell} \leq \inf \operatorname{supp} \phi_{k+1,\ell}$ for $1 \leq k \leq N_\ell - 1$.

With the exception of (e) and (i), all conditions are standard, and biorthogonal wavelets and scaling functions that satisfy them have been constructed in [20, 21, 22]. To satisfy (e), it is sufficient that $\mathbb{1} \in \operatorname{span}\{\psi_\lambda : \lambda \in J, |\lambda| = 0\}$, which in view of (b) is a natural condition that is satisfied by the constructions

in these references. Indeed, when this holds true then by means of a simple basis transformation, that only involves primal and dual wavelets on the lowest level, (e) is satisfied.

The second condition in (i) just fixes an ordering of the scaling functions by their supports. The first condition in (i) can always be satisfied by a rescaling of the scaling functions. However, in order that this rescaling does not jeopardize $\|\phi_{\ell,k}\|_{L_2(\mathcal{I})} \approx 1$, initially, for each ℓ , the $\int_{\mathcal{I}} \phi_{\ell,k}$'s should have comparable values. In view of $\int_{\mathcal{I}} \phi_{\ell,k} \lesssim 2^{-\ell/2}$, sufficient is $\int_{\mathcal{I}} \phi_{\ell,k} \gtrsim 2^{-\ell/2}$, which is satisfied by the B-spline scaling functions in the aforementioned references.

Remark 2.4 *From the interpolation space $[H^1(\mathcal{I})', H_0^1(\mathcal{I})]_{\frac{1}{2},2} = L_2(\mathcal{I})$, (a), (b), and (c) imply that Ψ , and so $\tilde{\Psi}$, are Riesz bases for $L_2(\mathcal{I})$.*

Next, from $(\Psi, \tilde{\Psi})$ we construct a new pair of biorthogonal wavelets $(\overset{\dagger}{\Psi}, \bar{\tilde{\Psi}})$, and corresponding primal and dual scaling functions, by means of integration or differentiation at primal or dual side, respectively. This generalizes the construction in [23, Proposition 7] for the shift-invariant case on \mathbb{R} .

Proposition 2.5 *With $\overset{\circ}{J} := J \setminus \{\lambda\}$, we define*

$$\overset{\dagger}{\Psi} = \{\overset{\dagger}{\psi}_{\lambda} : \lambda \in \overset{\circ}{J}\}, \quad \bar{\tilde{\Psi}} = \{\bar{\tilde{\psi}}_{\lambda} : \lambda \in \overset{\circ}{J}\},$$

by

$$\overset{\dagger}{\psi}_{\lambda} : x \mapsto \int_0^x 2^{|\lambda|} \psi_{\lambda}(y) dy, \quad \bar{\tilde{\psi}}_{\lambda} = -2^{-|\lambda|} \tilde{\psi}'_{\lambda}.$$

Then

- (i). $\overset{\dagger}{\Psi}, \bar{\tilde{\Psi}}$ are $L_2(\mathcal{I})$ -biorthogonal collections,
- (ii). $\overset{\dagger}{\Psi}$ is a Riesz basis for $L_2(\mathcal{I})$, and
- (iii). $\{4^{-|\lambda|} \overset{\dagger}{\psi}_{\lambda} : \lambda \in \overset{\circ}{J}\}$ is a Riesz basis for $H_0^2(\mathcal{I})$.
- (iv). $\overset{\dagger}{\Psi}$ and $\bar{\tilde{\Psi}}$ are local.

Proof. Obviously $\text{supp } \bar{\tilde{\psi}}_{\lambda} \subset \text{supp } \tilde{\psi}_{\lambda}$. Since for $\lambda \in \overset{\circ}{J}$, $\int_{\mathcal{I}} \psi_{\lambda} = 0$ by (a) and (e), we infer that $\text{supp } \overset{\dagger}{\psi}_{\lambda} \subset \text{convhull}(\text{supp } \psi_{\lambda})$, showing (iv), as well as $\overset{\dagger}{\psi}_{\lambda} \in H_0^1(\mathcal{I})$, the latter showing that

$$\langle \overset{\dagger}{\psi}_{\lambda}, \bar{\tilde{\psi}}_{\mu} \rangle_{L_2(\mathcal{I})} = \langle \overset{\dagger}{\psi}_{\lambda}, -2^{-|\mu|} \tilde{\psi}'_{\mu} \rangle_{L_2(\mathcal{I})} = 2^{|\lambda|-|\mu|} \langle \psi_{\lambda}, \tilde{\psi}_{\mu} \rangle_{L_2(\mathcal{I})},$$

and thus (i) by (a).

Since $H_0^1(\mathcal{I}) \cap \text{span}\{\mathbf{1}\}^{\perp L_2(\mathcal{I})} \rightarrow H_0^2(\mathcal{I}) : g \mapsto (x \mapsto \int_0^x g(y) dy)$ is bounded, with bounded inverse $f \mapsto f'$, $\{2^{-|\lambda|} \psi_{\lambda} : \lambda \in \overset{\circ}{J}\}$ being a Riesz basis for $H_0^1(\mathcal{I}) \cap \text{span}\{\mathbf{1}\}^{\perp L_2(\mathcal{I})}$, by (c), (e), (a), is equivalent to $\{4^{-|\lambda|} \overset{\dagger}{\psi}_{\lambda} : \lambda \in \overset{\circ}{J}\}$ being a Riesz basis for $H_0^2(\mathcal{I})$, i.e., (iii).

Since $H^1(\mathcal{I}) \cap \text{span}\{\psi_{\lambda}\}^{\perp L_2(\mathcal{I})} \rightarrow L_2(\mathcal{I}) : f \mapsto f'$ is bounded, with bounded inverse $g \mapsto \left(x \mapsto \int_0^x g(y) dy - \frac{\int_0^1 \int_0^z g(y) dy \psi_{\lambda}(z) dz}{\int_0^1 \psi_{\lambda}(z) dz}\right)$, $\{2^{-|\lambda|} \tilde{\psi}_{\lambda} : \lambda \in \overset{\circ}{J}\}$ being a Riesz basis for $H^1(\mathcal{I}) \cap \text{span}\{\psi_{\lambda}\}^{\perp L_2(\mathcal{I})}$, by (b) and (a), is equivalent to $\bar{\tilde{\Psi}}$ being a Riesz basis for $L_2(\mathcal{I})$, i.e., (ii) by (i). \square

Remark 2.6 *From $[L_2(\mathcal{I}), H_0^2(\mathcal{I})]_{\frac{1}{2},2} = H_0^1(\mathcal{I})$, (ii) and (iii) imply that $\{2^{-|\lambda|} \overset{\dagger}{\psi}_{\lambda} : \lambda \in \overset{\circ}{J}\}$ is a Riesz basis for $H_0^1(\mathcal{I})$.*

Proposition 2.7 *The collections $\overset{+}{\Phi}_\ell = \{\overset{+}{\phi}_{\ell,k} : 1 \leq k \leq N_\ell - 1\}$, $\bar{\Phi}_\ell = \{\bar{\phi}_{\ell,k} : 1 \leq k \leq N_\ell - 1\}$, defined by*

$$\overset{+}{\phi}_{\ell,k} : x \mapsto \int_0^x 2^{\ell+1}(\phi_{\ell,k+1}(y) - \phi_{\ell,k}(y))dy, \quad \bar{\phi}_{\ell,k} = -2^{-(\ell+1)} \sum_{p=k+1}^{N_\ell} \tilde{\phi}'_{\ell,p},$$

are biorthogonal, uniformly local, uniform $L_2(\mathcal{I})$ -Riesz bases for $\text{span}\{\overset{+}{\psi}_\lambda : \lambda \in \overset{\circ}{J}, |\lambda| \leq \ell\}$, $\text{span}\{\bar{\psi}_\lambda : \lambda \in \overset{\circ}{J}, |\lambda| \leq \ell\}$, respectively.

Proof. An application of a basis transform shows that

$$\{\phi_{\ell,1}, \phi_{\ell,2} - \phi_{\ell,1}, \dots, \phi_{\ell,N_\ell} - \phi_{\ell,N_\ell-1}\}, \quad \left\{ \sum_{k=1}^{N_\ell} \tilde{\phi}_{\ell,k}, \sum_{k=2}^{N_\ell} \tilde{\phi}_{\ell,k}, \dots, \tilde{\phi}_{\ell,N_\ell} \right\} \quad (6)$$

are biorthogonal bases for $\text{span } \Phi_\ell$, $\text{span } \bar{\Phi}_\ell$.

Because of $\int_{\mathcal{I}} \phi_{\ell,k+1} - \phi_{\ell,k} = 0$ by (i), and so $\overset{+}{\Phi}_\ell \subset H_0^1(\mathcal{I})$, integration by parts shows that $\overset{+}{\Phi}_\ell$, $\bar{\Phi}_\ell$ are biorthogonal.

Again (i) and $\mathbf{1} \in \text{span } \bar{\Phi}_\ell$, by (e), show that $\sum_{k=1}^{N_\ell} \tilde{\phi}_{\ell,k} \in \text{span}\{\mathbf{1}\}$. So for $\lambda \in \overset{\circ}{J}$, $|\lambda| \leq \ell$, we have $\psi_\lambda \in \text{span}\{\phi_{\ell,2} - \phi_{\ell,1}, \dots, \phi_{\ell,N_\ell} - \phi_{\ell,N_\ell-1}\}$, and so $\overset{+}{\psi}_\lambda \in \text{span } \overset{+}{\Phi}_\ell$. Since $\sum_{k=1}^{N_\ell} \tilde{\phi}'_{\ell,k} = 0$, for $\lambda \in \overset{\circ}{J}$, $|\lambda| \leq \ell$, we have $\bar{\psi}_\lambda \in \text{span } \bar{\Phi}_\ell$.

By $\text{supp } \overset{+}{\phi}_{\ell,k} \subset \text{convhull}(\text{supp } \phi_{\ell,k+1} \cup \text{supp } \phi_{\ell,k})$ and Φ_ℓ being uniformly local ((h)), we have that $\overset{+}{\Phi}_\ell$ is uniformly local. This property together with $\|\phi_{\ell,k}\|_{L_2(\mathcal{I})} \lesssim 1$ shows that $\|\overset{+}{\phi}_{\ell,k}\|_{L_2(\mathcal{I})} \lesssim 1$, and so, again by Φ_ℓ being uniformly local, that $\|\sum_{k=1}^{N_\ell-1} c_k \overset{+}{\phi}_{\ell,k}\|_{L_2(\mathcal{I})}^2 \lesssim \sum_{k=1}^{N_\ell-1} c_k^2$.

From $\sum_{k=1}^{N_\ell} \tilde{\phi}'_{\ell,k} = 0$, the ordering of the $\tilde{\phi}'_{\ell,k}$ by (i) and $\text{supp } \tilde{\phi}_{\ell,k} \cap \text{supp } \phi_{\ell,k} \neq \emptyset$, and $\bar{\Phi}_\ell$ being uniformly local ((h)), it follows that $\bar{\Phi}_\ell$ is uniformly local.

Any $u \in \text{span } \bar{\Phi}_\ell$ can be written as $\sum_{\{\lambda \in \overset{\circ}{J} : |\lambda| \leq \ell\}} c_\lambda 2^{-|\lambda|} \bar{\psi}_\lambda$. From $\{2^{-|\lambda|} \bar{\psi}_\lambda : \lambda \in \overset{\circ}{J}\}$ and $\bar{\Psi}$ being Riesz bases for $H^1(\mathcal{I})$ and $L_2(\mathcal{I})$, respectively, by (b) and Remark 2.4, we have $\|u\|_{H^1(\mathcal{I})}^2 \approx \sum_{\{\lambda \in \overset{\circ}{J} : |\lambda| \leq \ell\}} |c_\lambda|^2 \leq 4^\ell \sum_{\{\lambda \in \overset{\circ}{J} : |\lambda| \leq \ell\}} |c_\lambda 2^{-|\lambda|}|^2 \lesssim 4^\ell \|u\|_{L_2(\mathcal{I})}^2$.

From this so-called inverse inequality, $\|\tilde{\phi}_{\ell,k}\|_{L_2(\mathcal{I})} \lesssim 1$, and $\bar{\Phi}_\ell$ being uniformly local, it follows that $\|\bar{\phi}_{\ell,k}\|_{L_2(\mathcal{I})} \lesssim 1$, and so again from $\bar{\Phi}_\ell$ being local, that $\|\sum_{k=1}^{N_\ell-1} c_k \bar{\phi}_{\ell,k}\|_{L_2(\mathcal{I})}^2 \lesssim \sum_{k=1}^{N_\ell-1} c_k^2$, which by biorthogonality is equivalent to $\|\sum_{k=1}^{N_\ell-1} c_k \overset{+}{\phi}_{\ell,k}\|_{L_2(\mathcal{I})}^2 \gtrsim \sum_{k=1}^{N_\ell-1} c_k^2$. We conclude that $\overset{+}{\Phi}_\ell$, and so $\bar{\Phi}_\ell$, are uniform $L_2(\mathcal{I})$ -Riesz bases for their spans. \square

Having two biorthogonal multi-resolution analyses related by integration or differentiation at hand, we are ready to construct a wavelet Riesz basis for $\{v \in H_0^1(\mathcal{I}^2)^2 : \text{div } v = 0\}$, as well as a corresponding dual basis. Let us denote these bases here by Σ and $\tilde{\Sigma}$, respectively. With $\tilde{\Sigma}$ being a dual basis, we mean that $\tilde{\Sigma} \subset H^{-1}(\mathcal{I}^2)^2$, $\langle \tilde{\Sigma}, \Sigma \rangle_{H^{-1}(\mathcal{I}^2)^2 \times H_0^1(\mathcal{I}^2)^2} = \text{Id}$, and

$$v \mapsto \langle \tilde{\Sigma}, v \rangle_{H^{-1}(\mathcal{I}^2)^2 \times H_0^1(\mathcal{I}^2)^2} \in B(H_0^1(\mathcal{I}^2)^2, \ell_2(\tilde{\Sigma})). \quad (7)$$

Consequently, $v \mapsto \langle \tilde{\Sigma}, v \rangle_{H^{-1}(\mathcal{I}^2)^2 \times H_0^1(\mathcal{I}^2)^2} \Sigma$ is a bounded projection on $H_0^1(\mathcal{I}^2)^2$, with its image being equal to $\{v \in H_0^1(\mathcal{I}^2)^2 : \text{div } v = 0\}$. Note that such a dual basis is not unique.

Although our bases are similar to those constructed in [3] in the shift-invariant case on \mathbb{R}^d , working on a bounded domain causes some difficulties by which this construction of the isotropic divergence-free wavelets is restricted to two space dimensions. We refer to [4] for a further discussion of this point.

In the following, we set $I_\ell = \{1 \leq k \leq N_\ell - 1\}$, and for $\lambda \in \mathring{J}$, we write $\lambda = (\ell, k)$ where $\ell = |\lambda|$ and k runs over an index set J_ℓ , so that $\mathring{J} = \cup_{\ell=0}^{\infty} \{\ell\} \times J_\ell$.

Proposition 2.8 *The collection*

$$\begin{aligned} & \bigcup_{\ell \in \mathbb{N}_0} 2^{-\ell} \left(\left\{ [-\overset{\dagger}{\psi}_{\ell+1,k} \otimes (\phi_{\ell,m+1} - \phi_{\ell,m}), \psi_{\ell+1,k} \otimes \overset{\dagger}{\phi}_{\ell,m}]^\top : k \in J_{\ell+1}, m \in I_\ell \right\} \right. \\ & \quad \cup \left\{ [-\overset{\dagger}{\phi}_{\ell,k} \otimes \psi_{\ell+1,m}, (\phi_{\ell,k+1} - \phi_{\ell,k}) \otimes \overset{\dagger}{\psi}_{\ell+1,m}]^\top : k \in I_\ell, m \in J_{\ell+1} \right\} \\ & \quad \left. \cup \left\{ [-\overset{\dagger}{\psi}_{\ell,k} \otimes \psi_{\ell,m}, \psi_{\ell,k} \otimes \overset{\dagger}{\psi}_{\ell,m}]^\top : k \in J_\ell, m \in J_\ell \right\} \right) \end{aligned} \quad (8)$$

is a Riesz basis for $\{v \in H_0^1(\mathcal{I}^2)^2 : \operatorname{div} v = 0\}$.

A dual basis is given by

$$\begin{aligned} & \bigcup_{\ell \in \mathbb{N}_0} 2^\ell \left(\left\{ [0, \tilde{\psi}_{\ell+1,k} \otimes \bar{\phi}_{\ell,m}]^\top : k \in J_{\ell+1}, m \in I_\ell \right\} \right. \\ & \quad \cup \left\{ [-\bar{\phi}_{\ell,k} \otimes \tilde{\psi}_{\ell+1,m}, 0]^\top : k \in I_\ell, m \in J_{\ell+1} \right\} \\ & \quad \left. \cup \left\{ [-\bar{\psi}_{\ell,k} \otimes \tilde{\psi}_{\ell,m}, \tilde{\psi}_{\ell,k} \otimes \bar{\psi}_{\ell,m}]^\top : k \in J_\ell, m \in J_\ell \right\} \right). \end{aligned} \quad (9)$$

Both the primal and dual basis are local, meaning both that the diameter of the support of a wavelet on “level ℓ ” is $\lesssim 2^{-\ell}$, and that each ball of diameter $2^{-\ell}$ intersects the supports of at most uniformly bounded number of wavelets on level ℓ .

Proof. From $\overset{\dagger}{\Psi}$ and $\{4^{-|\lambda|} \overset{\dagger}{\psi}_\lambda : \lambda \in \mathring{J}\}$ being Riesz bases for $L_2(\mathcal{I})$ and $H_0^2(\mathcal{I})$, respectively, by (10), and $H_0^2(\mathcal{I}^2) \simeq H_0^2(\mathcal{I}) \otimes L_2(\mathcal{I}) \cap L_2(\mathcal{I}) \otimes H_0^2(\mathcal{I})$, we have that

$$\left\{ \left(\sum_{j=1}^2 16^{|\lambda_j|} \right)^{-\frac{1}{2}} \overset{\dagger}{\psi}_{\lambda_1} \otimes \overset{\dagger}{\psi}_{\lambda_2} : (\lambda_1, \lambda_2) \in \mathring{J} \times \mathring{J} \right\} \text{ is a Riesz basis for } H_0^2(\mathcal{I}^2). \quad (10)$$

By using that $\{\overset{\dagger}{\psi}_{\lambda_1} \otimes \overset{\dagger}{\psi}_{\lambda_2} : (\lambda_1, \lambda_2) \in \mathring{J} \times \mathring{J}\}$ is a Riesz basis for $L_2(\mathcal{I}^2)$, we infer that for $\ell \in \mathbb{N}_0$,

$$\|\cdot\|_{H^2(\mathcal{I}^2)} \approx 4^\ell \|\cdot\|_{L_2(\mathcal{I}^2)} \quad \text{on } \operatorname{span}\{\overset{\dagger}{\psi}_{\lambda_1} \otimes \overset{\dagger}{\psi}_{\lambda_2} : (\lambda_1, \lambda_2) \in \mathring{J} \times \mathring{J}, \max(|\lambda_1|, |\lambda_2|) = \ell\}. \quad (11)$$

With $\overset{\dagger}{\Psi}_{[\ell]} := \{\overset{\dagger}{\psi}_\lambda : \lambda \in \mathring{J}, |\lambda| = \ell\}$, for $\ell > 0$ an alternative, uniform $L_2(\mathcal{I}^2)$ -basis for the space from (11) is given by

$$\overset{\dagger}{\Psi}_{[\ell]} \otimes \overset{\dagger}{\Phi}_{\ell-1} \cup \overset{\dagger}{\Phi}_{\ell-1} \otimes \overset{\dagger}{\Psi}_{[\ell]} \cup \overset{\dagger}{\Psi}_\ell \otimes \overset{\dagger}{\Psi}_{[\ell]}.$$

The latter result, (11), and (10) show that

$$\bigcup_{\ell \in \mathbb{N}_0} 2^{-\ell} (2^{-(\ell+1)} \overset{\dagger}{\Psi}_{[\ell+1]} \otimes \overset{\dagger}{\Phi}_\ell \cup 2^{-(\ell+1)} \overset{\dagger}{\Phi}_\ell \otimes \overset{\dagger}{\Psi}_{[\ell+1]} \cup 2^{-\ell} \overset{\dagger}{\Psi}_{[\ell]} \otimes \overset{\dagger}{\Psi}_{[\ell]})$$

is a Riesz basis for $H_0^2(\mathcal{I}^2)$.

By applying minus **curl** to this collection, the collection in the statement of the proposition is obtained. Since, as follows from [13, §I.3.1], **curl** : $H_0^2(\mathcal{I}^2) \rightarrow \{v \in H_0^1(\mathcal{I}^2)^2 : \operatorname{div} v = 0\}$ is boundedly invertible, the first statement is proven.

The biorthogonality of the collections from (8) and (9) follows from the biorthogonality of $(\Psi, \tilde{\Psi})$, $(\check{\Psi}, \check{\tilde{\Psi}})$, $(\Phi_\ell, \check{\Phi}_\ell)$, $(\check{\Phi}_\ell, \check{\tilde{\Phi}}_\ell)$, and $\text{span}\{\check{\psi}_\lambda : \lambda \in J, |\lambda| \leq \ell\} = \text{span}\check{\Phi}_\ell$, $\text{span}\{\tilde{\psi}_\lambda : \lambda \in J, |\lambda| \leq \ell\} = \text{span}\check{\tilde{\Phi}}_\ell$, and (f).

The locality of both the primal and dual collections follows directly from the (uniform) locality of the primal and dual scaling functions and wavelets from both biorthogonal multiresolution analyses.

What remains to show is the property (7) for the dual collection (9). From Ψ and $\check{\Psi}$, and $\{2^{-|\lambda|}\psi_\lambda : \lambda \in J\}$ and $\{4^{-|\lambda|}\check{\psi}_\lambda : \lambda \in \check{J}\}$ being Riesz bases for $L_2(\mathcal{I})$ and $H_0^1(\mathcal{I})$, respectively, by (a), (c), (ii), and Remark 2.6, and $H_0^1(\mathcal{I}^2) \simeq H_0^1(\mathcal{I}) \otimes L_2(\mathcal{I}) \cap L_2(\mathcal{I}) \otimes H_0^1(\mathcal{I})$, we have that

$$\left\{ \left(\sum_{j=1}^2 4^{|\lambda_j|} \right)^{-\frac{1}{2}} \check{\psi}_{\lambda_1} \otimes \psi_{\lambda_2} : (\lambda_1, \lambda_2) \in \check{J} \times J \right\} \text{ is a Riesz basis for } H_0^1(\mathcal{I}^2). \quad (12)$$

By using that $\{\check{\psi}_{\lambda_1} \otimes \psi_{\lambda_2} : (\lambda_1, \lambda_2) \in \check{J} \times J\}$ is a Riesz basis for $L_2(\mathcal{I}^2)$, we infer that for $\ell \in \mathbb{N}_0$,

$$\|\cdot\|_{H^1(\mathcal{I}^2)} \approx 2^\ell \|\cdot\|_{L_2(\mathcal{I}^2)} \quad \text{on } \text{span}\{\check{\psi}_{\lambda_1} \otimes \psi_{\lambda_2} : (\lambda_1, \lambda_2) \in \check{J} \times J, \max(|\lambda_1|, |\lambda_2|) = \ell\}. \quad (13)$$

With $\Psi_{[\ell]} := \{\psi_\lambda : \lambda \in J, |\lambda| = \ell\}$, for $\ell > 0$ an alternative, uniform $L_2(\mathcal{I}^2)$ -basis for the space from (8) is given by

$$\check{\Psi}_{[\ell]} \otimes \Phi_{\ell-1} \cup \check{\Phi}_{\ell-1} \otimes \Psi_{[\ell]} \cup \check{\Psi}_\ell \otimes \Psi_{[\ell]}.$$

The latter result, (13), and (12) show that

$$\bigcup_{\ell \in \mathbb{N}_0} 2^{-\ell} (\check{\Psi}_{[\ell+1]} \otimes \Phi_\ell \cup \check{\Phi}_\ell \otimes \Psi_{[\ell+1]} \cup \check{\Psi}_{[\ell]} \otimes \Psi_{[\ell]})$$

is a Riesz basis for $H_0^1(\mathcal{I}^2)$. Its (unique) dual basis in $H^{-1}(\mathcal{I}^2)$ reads as

$$\bigcup_{\ell \in \mathbb{N}_0} 2^\ell (\check{\tilde{\Psi}}_{[\ell+1]} \otimes \check{\Phi}_\ell \cup \check{\tilde{\Phi}}_\ell \otimes \check{\tilde{\Psi}}_{[\ell+1]} \cup \check{\tilde{\Psi}}_{[\ell]} \otimes \check{\tilde{\Psi}}_{[\ell]}),$$

with the obvious definitions of $\check{\tilde{\Psi}}_{[\ell]}$ and $\check{\tilde{\Psi}}_{[\ell]}$.

We conclude that

$$\begin{aligned} v &\mapsto (\langle 2^\ell \check{\tilde{\Phi}}_{\ell,k} \otimes \check{\tilde{\psi}}_{\ell+1,m}, v \rangle_{H^{-1}(\mathcal{I}^2) \times H_0^1(\mathcal{I}^2)})_{k \in I_\ell, m \in J_{\ell+1}, \ell \in \mathbb{N}_0}, \\ v &\mapsto (\langle 2^\ell \check{\tilde{\psi}}_{\ell,k} \otimes \check{\tilde{\psi}}_{\ell,m}, v \rangle_{H^{-1}(\mathcal{I}^2) \times H_0^1(\mathcal{I}^2)})_{k \in J_\ell, m \in J_\ell, \ell \in \mathbb{N}_0} \end{aligned}$$

and, analogously,

$$\begin{aligned} v &\mapsto (\langle 2^\ell \check{\tilde{\psi}}_{\ell+1,k} \otimes \check{\tilde{\Phi}}_{\ell,m}, v \rangle_{H^{-1}(\mathcal{I}^2) \times H_0^1(\mathcal{I}^2)})_{k \in J_{\ell+1}, m \in I_\ell, \ell \in \mathbb{N}_0}, \\ v &\mapsto (\langle 2^\ell \check{\tilde{\psi}}_{\ell,k} \otimes \check{\tilde{\psi}}_{\ell,m}, v \rangle_{H^{-1}(\mathcal{I}^2) \times H_0^1(\mathcal{I}^2)})_{k \in J_\ell, m \in J_\ell, \ell \in \mathbb{N}_0} \end{aligned}$$

are bounded mappings from $H_0^1(\mathcal{I}^2)$ to the corresponding ℓ_2 -spaces, which proves (7). \square

2.5 The Navier-Stokes equations as an infinite system of scalar nonlinear equations

With the wavelet frame $\Psi = \{\psi_\lambda : \lambda \in \Lambda\} = \bigcup_{i=0}^{m-1} E_i \Psi_i$ for V , with the Ψ_i as given in (8), and corresponding frame operator $F : V' \rightarrow \ell_2(\Lambda)$, we can now reformulate equation (1) as an equivalent

infinite system of scalar nonlinear equations. Because of $\overline{\text{span } \Psi} = V$, the weak form is equivalent to finding a vector $\mathbf{u} \in \ell_2(\Lambda)$ such that

$$\mathbf{A}\mathbf{u} + \text{Re } \mathbf{G}(\mathbf{u}) = \text{Re } \mathbf{f},$$

where \mathbf{A} is the infinite-dimensional stiffness matrix $\mathbf{A} = \{a(\psi_\lambda, \psi_\mu)\}_{\mu, \lambda \in \Lambda}$, \mathbf{G} is the discrete nonlinearity $\mathbf{G}(\mathbf{u}) = (\int_\Omega \psi_\lambda \cdot (u \cdot \nabla)u)_{\lambda \in \Lambda}$, $u = \mathbf{u}^\top \Psi$ and \mathbf{f} is the discrete right-hand side $\mathbf{f} = (\int_\Omega f \psi_\lambda)_{\lambda \in \Lambda}$.

From a discrete solution \mathbf{u} , the continuous solution can be retrieved by $u = F'\mathbf{u} = \mathbf{u}^\top \Psi$. It is important to note that, because the operator F' is not injective unless Ψ is a basis, uniqueness of the continuous solution u does not imply uniqueness of the discrete solution \mathbf{u} .

2.6 Approximation spaces

In the following, we are going to outline the concept of asymptotic optimality of adaptive wavelet methods. Assume that the original problem has a solution $u = \mathbf{u}^\top \Psi$, which has *some* discrete representation $\mathbf{u} \in \ell_2(\Lambda)$ in the given wavelet frame Ψ that can be approximated with rate s with respect to the degrees of freedom, i.e.,

$$\sup_{N \in \mathbb{N}} N^s \inf_{\#\text{supp } \mathbf{v} \leq N} \|\mathbf{u} - \mathbf{v}\|_{\ell_2(\Lambda)} < \infty.$$

Then, we expect our algorithm to achieve the same rate s .

Dealing with nonlinear problems it has turned out that we have to confine ourselves to approximations supported on a tree-type index sets. In the present context of wavelet frames constructed from wavelet bases in the fashion described above, we say that a set $T = \bigcup_{i=0}^{m-1} \{i\} \times T_i \subset \Lambda = \bigcup_{i=0}^{m-1} \{i\} \times \Lambda_i$ has an *aggregated tree* structure, if all the T_i are trees. Here we call $T_i \subset \Lambda_i$ a *tree*, when for any $\lambda \in T_i$ with $|\lambda| > 0$, $\text{supp } \psi_\lambda^{(i)}$ is covered by the supports of $\psi_\mu^{(i)}$ for some $\mu \in T_i$ with $|\mu| = |\lambda| - 1$. In the literature, slightly different definitions of tree index sets can be found, but the differences are harmless.

Now we define

$$\Sigma_{N, \mathcal{AT}} := \{\mathbf{v} \in \ell_2(\Lambda), \#\text{supp } \mathbf{v} \leq N, \text{supp } \mathbf{v} \text{ has aggregated tree structure}\}$$

and set $\sigma_{N, \mathcal{AT}}(\mathbf{v}) := \inf_{\mathbf{w} \in \Sigma_{N, \mathcal{AT}}} \|\mathbf{v} - \mathbf{w}\|_{\ell_2(\Lambda)}$. From this we obtain the approximation space

$$\mathcal{A}_{\mathcal{AT}}^s := \{\mathbf{v} \in \ell_2(\Lambda) : \sigma_{N, \mathcal{AT}}(\mathbf{v}) \lesssim N^{-s}\}$$

of vectors in $\ell_2(\Lambda)$ that can be approximated with rate s in aggregated tree structure equipped with the quasi-norm $\|\mathbf{v}\|_{\mathcal{A}_{\mathcal{AT}}^s} := \sup_{N \in \mathbb{N}} N^s \sigma_{N, \mathcal{AT}}(\mathbf{v})$. Later, we are going to show that $\mathbf{u} \in \mathcal{A}_{\mathcal{AT}}^s$, where \mathbf{u} is *some* representation of the solution $u \in V$, implies that the adaptive algorithm we are going to construct converges with rate s . This property is called *asymptotic optimality*.

For use later, we also set

$$\Sigma_{N, \mathcal{T}_i} := \{\mathbf{v} \in \ell_2(\Lambda_i), \#\text{supp } \mathbf{v} \leq N, \text{supp } \mathbf{v} \text{ is a tree}\},$$

$\sigma_{N, \mathcal{T}_i}(\mathbf{v}) := \inf_{\mathbf{w} \in \Sigma_{N, \mathcal{T}_i}} \|\mathbf{v} - \mathbf{w}\|_{\ell_2(\Lambda_i)}$, $\mathcal{A}_{\mathcal{T}_i}^s := \{\mathbf{v} \in \ell_2(\Lambda_i) : \sigma_{N, \mathcal{T}_i}(\mathbf{v}) \lesssim N^{-s}\}$ equipped with the quasi-norm $\|\mathbf{v}\|_{\mathcal{A}_{\mathcal{T}_i}^s} := \sup_{N \in \mathbb{N}} N^s \sigma_{N, \mathcal{T}_i}(\mathbf{v})$, as well as $\Sigma_{N, i}$, $\sigma_{N, i}$, and \mathcal{A}_i^s equipped with $\|\mathbf{v}\|_{\mathcal{A}_i^s}$, defined by omitting the tree constrained on $\text{supp } \mathbf{v}$.

The question for which s we can find a discrete solution $\mathbf{u} \in \mathcal{A}_{\mathcal{AT}}^s$, and hence obtain convergence of an asymptotically optimal adaptive method with rate s , is linked to the Besov regularity of the continuous

solution in an appropriate scale, see [1, 2, 24]. In general Lipschitz domains, to our best knowledge, little is known about regularity of solutions to the incompressible Navier-Stokes equation with respect to this Besov scale. However, for the related Stokes equation and the three-dimensional Navier-Stokes equation in polyhedral cones, it can be shown that in many cases the Besov regularity indeed exceeds the Sobolev regularity, see [25] and [26], respectively. Moreover, numerical experiments from [6] suggest that, even in other cases, the regularity measured in this scale is significantly higher than in the Sobolev scale. Since the Sobolev regularity corresponds to the convergence rate of standard uniform methods, it seems reasonable to use adaptive methods in order to improve the convergence rate.

3 The adaptive algorithm

In this section, we are going to construct an adaptive wavelet Schwarz solver for equation (1) written in divergence-free wavelet frame coordinates. To do so, we are first going to briefly present some tools needed for the construction.

3.1 Building Blocks

For the design of an asymptotically optimal adaptive wavelet method, we need to have at hand a couple of elementary building blocks.

First of all, we require a solver for linear subproblems on the subdomains Ω_i . For the construction of such local solvers, we refer, e.g., to [1]. Even though these subproblems are fully linear in the unknowns, we will have to evaluate the nonlinear term $(v \cdot \nabla)v$ in wavelet coordinates to obtain the right-hand side for the subproblems. The evaluation of such nonlinearities is described in [2].

Furthermore, in order to guarantee an optimal balance between degrees of freedom and accuracy, we repeatedly remove very small entries from the discrete iterates. This will be done by the application of a method

$$\mathbf{COARSE}[\mathbf{v}, \varepsilon] : \ell_2(\Lambda) \rightarrow \ell_2(\Lambda)$$

that maps a finitely supported $\mathbf{v} \in \ell_2(\Lambda)$ to a near-smallest $\mathbf{v}_\varepsilon \in \ell_2(\Lambda)$ with $\|\mathbf{v} - \mathbf{v}_\varepsilon\|_{\ell_2(\Lambda)} \leq \varepsilon$. The construction of such a method involves sorting the entries of \mathbf{v} into buckets by their modulus. For details and further properties, see, for instance, [1, 24]. In particular, it is shown in [24] that there exists a constant $\vartheta \in (0, 1)$ such that for $\mathbf{v} \in \mathcal{A}_{\mathcal{AT}}^s$ and a finitely supported $\mathbf{w} \in \ell_2(\Lambda)$ with $\|\mathbf{v} - \mathbf{w}\|_{\ell_2(\Lambda)} \leq \vartheta\varepsilon$, it holds that $\tilde{\mathbf{w}} := \mathbf{COARSE}[\mathbf{w}, (1 - \vartheta)\varepsilon] \in \mathcal{A}_{\mathcal{AT}}^s$ with $\|\tilde{\mathbf{w}}\|_{\mathcal{A}_{\mathcal{AT}}^s} \lesssim \|\mathbf{w}\|_{\mathcal{A}_{\mathcal{AT}}^s}$ and $\#\text{supp } \tilde{\mathbf{w}} \lesssim \varepsilon^{-1/s} \|\mathbf{v}\|_{\mathcal{A}_{\mathcal{AT}}^s}^{1/s}$.

The second building block that we describe is designed to deal with the redundancy of a frame, i.e., with the fact that $\ker F' \neq \{0\}$. Any vector in $\ker F'$ is in the kernel of $\mathbf{u} \mapsto \mathbf{A}\mathbf{u} + \text{Re } \mathbf{G}\mathbf{u}$, and therefore is not affected by any iterative method to invert this operator. So components in $\ker F'$ which arise in the course of the iteration as a consequence of inexact evaluation of operators, or because of applications of \mathbf{COARSE} will never be reduced. Assuming u has some representation $u = \mathbf{u}^\top \Psi$ with $\mathbf{u} \in \mathcal{A}_{\mathcal{AT}}^s$, this may have as a consequence that the iterands converge to a representation that is not in $\mathcal{A}_{\mathcal{AT}}^s$, so that consequently an optimal rate s is not realized.

For a frame that is the union of Riesz bases on overlapping subdomains, a way to deal with this problem is, before solving on subdomain i , to remove terms in the expansion of the current iterand that are multiples of wavelets $\psi_\lambda^{(j)} \subset \Psi_j$ for $j \neq i$ with $\text{supp } \psi_\lambda^{(j)} \subset \Omega_i$. In any case for linear elliptic problems,

and assuming a sufficiently large overlap of the subdomains in relation to the maximal diameter of the support of any primal or dual wavelet, in [10] it was shown that for the multiplicative Schwarz method this approach yields an adaptive algorithm that converges with the optimal rate. In [18], it was shown that the same holds true for the additive Schwarz algorithm in case of having two subdomains, whereas numerical experiments indicate that this is also valid for more than two subdomains.

Since no proof of the latter is available, to cope with the redundancy, for completeness here we will resort on the technique introduced in [7]. Under some circumstances, however, the routine **PROJECTION** that will be introduced below can simply be omitted from the adaptive algorithm, whereas nevertheless optimal rates can be observed. We refer to [7, Thm. 3.12, §4.3] for an analysis in a restricted setting, and to [11, 27] for numerical results.

Let Z be a bounded right-inverse of F' , i.e., $Z \in B(V, \ell_2(\Lambda))$ with $F'Z = \text{Id}$, and with the projector $\mathbf{Q} := ZF' \in B(\ell_2(\Lambda), \ell_2(\Lambda))$, let \mathbf{Q} be bounded on $\mathcal{A}_{\mathcal{AT}}^s$. A suitable Z will be constructed below. The application of \mathbf{Q} to a vector in $\ell_2(\Lambda)$ does not change the function u it is representing, i.e., $F'\mathbf{Q} = F'$. Yet, if u has *some* representation $\mathbf{u} \in \mathcal{A}_{\mathcal{AT}}^s$, then the application of \mathbf{Q} to any representation \mathbf{v} of u yields a representation in $\mathcal{A}_{\mathcal{AT}}^s$, because $\mathbf{Q}\mathbf{v} = \mathbf{Q}\mathbf{u}$ and \mathbf{Q} is bounded on $\mathcal{A}_{\mathcal{AT}}^s$. In view of this property, we extend our adaptive wavelet method with a recurrent, inexact application of \mathbf{Q} in order to produce a sequence of iterands that is uniformly bounded in $\mathcal{A}_{\mathcal{AT}}^s$.

The routine that approximates the application of \mathbf{Q} within tolerance $\varepsilon > 0$ will be denoted as

$$\mathbf{PROJECTION}[\mathbf{v}, \varepsilon] : \ell_2(\Lambda) \rightarrow \ell_2(\Lambda).$$

It maps a $\mathbf{v} \in \ell_2(\Lambda)$ to a $\mathbf{w}_\varepsilon \in \ell_2(\Lambda)$ with $\|\mathbf{w}_\varepsilon - \mathbf{Q}\mathbf{v}\|_{\ell_2(\Lambda)} \leq \varepsilon$.

Now we come to the construction of a suitable Z and thus of \mathbf{Q} . For $\delta > 0$ and $0 \leq i \leq m-1$, we set $\Omega_i(-\delta) := \{x \in \Omega_i : B(x; \delta) \cap \Omega \subset \Omega_i\}$. We will assume a sufficiently large overlap of the subdomains in relation to the maximal diameter of any primal or dual wavelet from (8) and (9) in the sense that there exists a $\delta > 0$, that from here on will be fixed, such that

$$\text{diam supp } \psi_\lambda^{(i)} \cup \text{supp } \tilde{\psi}_\lambda^{(i)} \leq \delta/2, \quad (14)$$

$$\Omega \subset \bigcup_{i=0}^{m-1} \Omega_i(-(1 + \frac{m-1}{2})\delta). \quad (15)$$

Lemma 3.1 *With $\Lambda_i^\delta := \{\lambda \in \Lambda_i : \text{supp } \psi_\lambda^{(i)} \cap \Omega_i(-\delta) \neq \emptyset\}$, we have $v \mapsto (\tilde{\psi}_\lambda^{(i)}(v))_{\lambda \in \Lambda_i^\delta} \in B(V, \ell_2(\Lambda_i^\delta))$, and, for $v \in V$, $v - \sum_{\lambda \in \Lambda_i^\delta} \tilde{\psi}_\lambda^{(i)}(v) \psi_\lambda^{(i)}$ vanishes on $\Omega_i(-\delta)$.*

Proof. $\Omega = \Omega_i \cup (\Omega \setminus \Omega_i(-\delta/2))$ is an overlapping domain decomposition, and so, as in the proof of Lemma 2.2, for all $v \in V$ there exist $v_i \in V_i$ and $w_i \in \{w \in H_0^1(\Omega \setminus \Omega_i(-\delta/2)) : \text{div } w = 0\}$ with $v_i + w_i = v$ and $\|v_i\|_{H^1(\Omega)^d}^2 + \|w_i\|_{H^1(\Omega)^d}^2 \approx \|v\|_{H^1(\Omega)^d}^2$. Noting that by (14), $(\tilde{\psi}_\lambda^{(i)}(v))_{\lambda \in \Lambda_i^\delta}$ only depends on $v|_{\Omega_i(-\delta/2)}$, both statements follow from $v = v_i$ on $\Omega_i(-\delta/2)$, for the second statement using that $\Psi^{(i)}$ is a Riesz basis for V_i with dual basis $\tilde{\Psi}^{(i)}$, and the definition of Λ_i^δ . \square

Next, with $Z_{-1} = H_{-1} := 0$, for $0 \leq i \leq m-1$, we set

$$Z_i : V \rightarrow \ell_2(\Lambda_0) \times \cdots \times \ell_2(\Lambda_i) : v \mapsto (Z_{i-1}v, (\tilde{\psi}_\lambda^{(i)}(v - H_{i-1}Z_{i-1}v))_{\lambda \in \Lambda_i^\delta})$$

$$H_i : \ell_2(\Lambda_0) \times \cdots \times \ell_2(\Lambda_i) \rightarrow V : (\mathbf{v}_0, \dots, \mathbf{v}_i) \mapsto \sum_{j=0}^i \mathbf{v}_j^\top \Psi^{(j)}.$$

Proposition 3.2 *The mappings Z_i are bounded, and for $v \in V$, $v - H_i Z_i v$ vanishes on $\bigcup_{j=0}^i \Omega_j(-\frac{i}{2}\delta)$.*

Proof. The first statement follows easily from Lemma 3.1. For the second statement, we write

$$v - H_i Z_i v = (v - H_{i-1} Z_{i-1} v) - \sum_{\lambda \in \Lambda_i^\delta} \tilde{\psi}_\lambda^{(i)}(v - H_{i-1} Z_{i-1} v) \psi_\lambda^{(i)}.$$

The first term vanishes on $\bigcup_{j=0}^{i-1} \Omega_j(-\frac{i-1}{2}\delta)$, and so $v - H_i Z_i v$ vanishes on $\bigcup_{j=0}^{i-1} \Omega_j(-\frac{i}{2}\delta)$ by (14). By Lemma 3.1, $v - H_i Z_i v$ also vanishes on $\Omega_i(-\delta) \subset \Omega_i(-\frac{i}{2}\delta)$, which completes the proof. \square

Setting $Z = Z_{m-1}$, with a slight abuse of notation, we have $Z \in B(V, \ell_2(\Lambda))$, and, using (15), $F'Z = \text{Id}$.

Remark 3.3 *A small refinement of an argument used in the proof of Proposition 3.2 shows that (15) can be relaxed to $\Omega \subset \bigcup_{i=0}^{m-1} \Omega_i(-\frac{J}{2}\delta)$, where J is the maximal number of subdomains that have non-empty intersection.*

Finally, to show that \mathbf{Q} is bounded on $\mathcal{A}_{\mathcal{AT}}^s$, it suffices to show that each of the matrices $[\tilde{\psi}_\lambda^{(i)}(\psi_\mu^{(j)})]_{\lambda \in \Lambda_i^\delta, \mu \in \Lambda_j}$ is bounded from $\mathcal{A}_{\mathcal{T}_j}^s$ to $\mathcal{A}_{\mathcal{T}_i}^s$. Fixing $0 \leq i, j \leq m-1$, we know that $\mathbf{B} := [\tilde{\psi}_\lambda^{(i)}(\psi_\mu^{(j)})]_{\lambda \in \Lambda_i^\delta, \mu \in \Lambda_j} : \ell_2(\Lambda_j) \rightarrow \ell_2(\Lambda_i)$ is bounded. Let $\mathbf{v} \in \mathcal{A}_{\mathcal{T}_j}^s$ and $\varepsilon > 0$ be given. Then there exists a $\mathbf{v}_\varepsilon \in \ell_2(\Lambda_j)$ whose support is a tree, with $\|\mathbf{B}\|_{\ell_2(\Lambda_j) \rightarrow \ell_2(\Lambda_i)} \|\mathbf{v} - \mathbf{v}_\varepsilon\|_{\ell_2(\Lambda_j)} \leq \varepsilon/2$, $\#\text{supp } \mathbf{v}_\varepsilon \lesssim \varepsilon^{-1/s} \|\mathbf{v}\|_{\mathcal{A}_{\mathcal{T}_j}^s}^{1/s}$, and thus $\|\mathbf{v}_\varepsilon\|_{\mathcal{A}_{\mathcal{T}_j}^s} \lesssim \|\mathbf{v}\|_{\mathcal{A}_{\mathcal{T}_j}^s}$, cf. [2, Prop. 6.3].

In [7, §4.5], it was shown that for local $\Psi^{(j)}$ and $\tilde{\Psi}^{(i)}$, where the $\psi_\mu^{(j)}$ are spline wavelets, \mathbf{B} is a so-called s^* -compressible matrix, for a value of s^* that exceeds the best possible rate s for which membership $\mathbf{u} \in \mathcal{A}_{\mathcal{AT}}^s$ can be expected (assuming that this best possible rate is larger than $\frac{1}{2}$). Consequently, see e.g. [1, Corol. 3.10], when $s < s^*$ there exists a $\mathbf{w}_\varepsilon \in \ell_2(\Lambda_i)$ with $\|\mathbf{B}\mathbf{v}_\varepsilon - \mathbf{w}_\varepsilon\|_{\ell_2(\Lambda_i)} \leq \varepsilon/2$, thus $\|\mathbf{B}\mathbf{v} - \mathbf{w}_\varepsilon\|_{\ell_2(\Lambda_i)} \leq \varepsilon$, and $\#\text{supp } \mathbf{w}_\varepsilon \lesssim \varepsilon^{-1/s} \|\mathbf{v}_\varepsilon\|_{\mathcal{A}_{\mathcal{T}_j}^s}^{1/s} \leq \varepsilon^{-1/s} \|\mathbf{v}_\varepsilon\|_{\mathcal{A}_{\mathcal{T}_j}^s}^{1/s} \lesssim \varepsilon^{-1/s} \|\mathbf{v}\|_{\mathcal{A}_{\mathcal{T}_j}^s}^{1/s}$.

Let T_i be the smallest tree in Λ_i that contains $\text{supp } \mathbf{w}_\varepsilon$. To conclude that $\mathbf{B} : \mathcal{A}_{\mathcal{T}_j}^s \rightarrow \mathcal{A}_{\mathcal{T}_i}^s$ is bounded, it remains to show that $\#T_i \lesssim \varepsilon^{-1/s} \|\mathbf{v}\|_{\mathcal{A}_{\mathcal{T}_j}^s}^{1/s}$. Because of $\Psi^{(i)}$ being local, there exists a constant $C > 0$ such that

$$\bar{T}_i := \{\theta \in \Lambda_i : \exists \lambda \in \text{supp } \mathbf{w}_\varepsilon \text{ s.t. } |\lambda| \geq |\theta| \wedge \text{dist}(\text{supp } \psi_\lambda^{(i)}, \text{supp } \psi_\theta^{(i)}) \leq C2^{-|\theta|}\} \supset T_i.$$

Obviously, for all $\lambda \in \text{supp } \mathbf{w}_\varepsilon$, there exists a $\mu \in \text{supp } \mathbf{v}_\varepsilon$ with $\text{supp } \tilde{\psi}_\lambda^{(i)} \cap \text{supp } \psi_\mu^{(j)} \neq \emptyset$. By the construction of the sparse approximations for \mathbf{B} in [7], which are used as ingredients of the approximate matrix-vector routine **APPLY** developed in [1], we have that if $|\lambda| \geq |\mu|$, then for all $\nu \in \Lambda_i^\delta$ with $|\mu| \leq |\nu| \leq |\lambda|$ and $\text{supp } \tilde{\psi}_\nu^{(i)} \cap \text{supp } \psi_\mu^{(j)} \neq \emptyset$ it holds that $\nu \in \text{supp } \mathbf{w}_\varepsilon$. (“Coincidentally” $(\mathbf{w}_\varepsilon)_\nu$ might be zero, in which case formally $\nu \notin \text{supp } \mathbf{w}_\varepsilon$. The point is, however, that when determining the aforementioned upper bound for $\#\text{supp } \mathbf{w}_\varepsilon$, ν has been counted as being part of the support. Related to this, below we will use that for any $\ell \in \mathbb{N}_0$ with $|\mu| \leq \ell \leq |\lambda|$, there exists a $\nu \in \Lambda_i^\delta$ with $|\nu| = \ell$ and $\text{supp } \tilde{\psi}_\nu^{(i)} \cap \text{supp } \psi_\mu^{(j)} \neq \emptyset$. Although this would be a mild assumption on $\tilde{\Psi}^{(i)}$, it is not needed to impose this, since again by determining the upper bound for $\#\text{supp } \mathbf{w}_\varepsilon$, the existence of such a ν has been taken into account.)

Now considering an arbitrary $\theta \in \bar{T}_i$, let $\lambda \in \text{supp } \mathbf{w}_\varepsilon$ be as in the definition of \bar{T}_i , and let $\mu \in \text{supp } \mathbf{v}_\varepsilon$ be as above. If $|\theta| \leq |\mu|$, then by definition of \bar{T}_i and $\text{supp } \mathbf{v}_\varepsilon$ being a tree, there exists a $\gamma \in \text{supp } \mathbf{v}_\varepsilon$ with $|\theta| = |\gamma|$ and $\text{dist}(\text{supp } \psi_\theta^{(i)}, \text{supp } \psi_\gamma^{(j)}) \lesssim 2^{-|\theta|}$. Otherwise, so when $|\theta| > |\mu|$, there exists a $\nu \in \text{supp } \mathbf{w}_\varepsilon$ with $|\theta| = |\nu|$ and $\text{dist}(\text{supp } \psi_\theta^{(i)}, \text{supp } \tilde{\psi}_\nu^{(i)}) \lesssim 2^{-|\theta|}$. From both observations, and the fact that $\Psi^{(i)}, \tilde{\Psi}^{(i)}, \Psi^{(j)}$ are local, we conclude that $\#T_i \leq \#\bar{T}_i \lesssim \#\text{supp } \mathbf{v}_\varepsilon + \#\text{supp } \mathbf{w}_\varepsilon \lesssim \varepsilon^{-1/s} \|\mathbf{v}\|_{\mathcal{A}_{T_j}^s}^{1/s}$, which completes the proof of \mathbf{Q} being bounded on $\mathcal{A}_{\mathcal{AT}}^s$.

3.2 Construction of the algorithm

We are now ready to define the algorithm we are going to investigate. This method is the adaptive wavelet version of the algorithm proposed in [12]. Note that bold letters stand for discrete iterates while standard letters stand for their continuous representation, e.g. $v = \mathbf{v}^\top \Psi$. To explicitly formulate the algorithm, we need to fix some constants. Let

$$K := \sup_{0 \neq \mathbf{v} \in \ell_2(\Lambda)} \frac{\|\mathbf{v}^\top \Psi\|}{\|\mathbf{v}\|_{\ell_2(\Lambda)}}, \quad L := \sup_{0 \neq \mathbf{v} \in \ell_2(\Lambda)} \frac{\|\mathbf{Q}\mathbf{v}\|_{\ell_2(\Lambda)}}{\|\mathbf{v}^\top \Psi\|},$$

$$C := \sup_{0 \neq u, v \in V} \left\{ \frac{\|z\|}{\|u\| \|v\|} : a(z, w) = \int_{\Omega} w \cdot (u \cdot \nabla) v, (w \in V) \right\},$$

where $K \leq B_\Psi \sup_{0 \neq v \in V} \frac{\|v\|}{\|v\|_{H^1(\Omega)^d}} < \infty$, $L = \sup_{0 \neq v \in V} \frac{\|Zv\|_{\ell_2(\Lambda)}}{\|v\|} < \infty$ by $\mathbf{Q} = ZF'$ and $Z \in B(V, \ell_2(\Lambda))$, and where $C < \infty$ for $d \leq 4$ has been shown in [12, Lemma 1]. Let M be an upper bound for $\|u\|$.

By P_i , we denote the $a(\cdot, \cdot)$ -orthogonal projector from V onto V_i . It is known that the operator norm

$$\theta := \|I - \omega(P_0 + \dots + P_{m-1})\|$$

on $B(V, V)$ is smaller than 1 if $\omega > 0$ is sufficiently small. Then, for small Reynolds numbers, we have

$$\rho := \theta + 3\omega \text{Re } mCM < 1. \quad (16)$$

We are going to show a convergence rate $\tilde{\rho} := (1 + \rho)/2 < 1$.

Algorithm 1 AddSchw

```

% Let  $l^* \in \mathbb{N}$  be minimal such that  $\tilde{\rho}^{l^*} \leq \frac{1}{2KL} \vartheta \tilde{\rho}$ .
% Let  $\varepsilon_n := \tilde{\rho}^n M$ ,  $n \in \mathbb{N}$ .
 $u^{(0)} := 0$ 
for  $n = 0, 1, \dots$  do
   $v^{(n,0)} := u^{(n)}$ 
  for  $l = 0, \dots, l^* - 1$  do
    for  $i = 0, \dots, m - 1$  do
      Compute  $\tilde{d}_i^{(n,l)} \in V_i$  as an approximation to the solution  $d_i^{(n,l)} \in V_i$  of
       $a(d_i^{(n,l)}, v) = -\text{Re} \int_{\Omega_i} v \cdot (v^{(n,l)} \cdot \nabla) v^{(n,l)} + \text{Re} \int_{\Omega_i} f \cdot v - a(v^{(n,l)}, v)$  for all  $v \in V_i$ 
      with tolerance  $\|d_i^{(n,l)} - \tilde{d}_i^{(n,l)}\| \leq \frac{1-\rho}{2m\omega} \varepsilon_n \tilde{\rho}^l$ .
    end for
     $v^{(n,l+1)} := v^{(n,l)} + \omega \sum_{i=0}^{m-1} \tilde{d}_i^{(n,l)}$ 
  end for
   $\tilde{\mathbf{u}}^{(n+1)} := \text{PROJECTION}[\mathbf{v}^{(n,l^*)}, \frac{\vartheta}{2K} \varepsilon_{n+1}]$ 
   $\mathbf{u}^{(n+1)} := \text{COARSE}[\tilde{\mathbf{u}}^{(n+1)}, \frac{1-\vartheta}{K} \varepsilon_{n+1}]$ 
end for

```

Note that the subproblems on the subdomains are fully linear and the nonlinearity only appears on the right-hand side. Moreover, they are independent so they can be solved in parallel. The first result concerning Algorithm 1 shows convergence given that the Reynolds number is sufficiently small such that (16) holds.

Theorem 3.4 *Assume that $d \leq 4$ and that (16) is valid. Then, the iterates from Algorithm 1 fulfill*

$$\|u^{(n)} - u\| \leq \tilde{\rho}^n M.$$

Proof. We show the assertion by induction. The case $n = 0$ is clear by definition. We assume the assertion is true for some $n \in \mathbb{N}$. Denote by $\bar{v}^{(n,1)} := v^{(n,0)} + \sum_{i=0}^{m-1} d_i^{(n,0)}$ the result of the first iteration step with exact subdomain solvers. As in the proof of Theorem 4 in [12], the error after this step can be written as

$$\bar{v}^{(n,1)} - u = (I - \omega(P_0 + \dots + P_{m-1}))(u^{(n)} - u) + \omega \operatorname{Re} \sum_{i=0}^{m-1} F_i(u^{(n)}, u),$$

where $F_i(u^{(n)}, u)$ is defined as

$$F_i(u^{(n)}, u) = -S_i^{-1}((u^{(n)} \cdot \nabla)u^{(n)} - (u \cdot \nabla)u)$$

and $S_i^{-1} : (H^{-1}(\Omega_i))^d \rightarrow V_i$ the solution operator for the Stokes equation on Ω_i . In the proof of Theorem 1 in [12], using $d \leq 4$, we see that

$$\|F_i(u^{(n)}, u)\| \leq C(\|u^{(n)} - u\| + 2\|u\|)\|u^{(n)} - u\|,$$

hence with $u^{(n)} = v^{(n,0)}$ we obtain the estimate

$$\|\bar{v}^{(n,1)} - u\| \leq \theta \|v^{(n,0)} - u\| + \omega \operatorname{Re} m C (\|v^{(n,0)} - u\| + 2\|u\|) \|v^{(n,0)} - u\|.$$

By the induction hypothesis, it is $\|v^{(n,0)} - u\| = \|u^{(n)} - u\| \leq \|u^{(0)} - u\| = \|u\| \leq M$. From this we obtain

$$\|\bar{v}^{(n,1)} - u\| \leq \rho \|v^{(n,0)} - u\| \leq \rho \varepsilon_n,$$

where $\varepsilon_n = \tilde{\rho}^n M$. Taking into account the tolerance for the error in the inexact solutions of the local problems, we obtain

$$\|v^{(n,1)} - u\| \leq \|\bar{v}^{(n,1)} - u\| + m\omega \frac{1-\rho}{2m\omega} \varepsilon_n \leq \left(\rho + \frac{1-\rho}{2}\right) \varepsilon_n = \tilde{\rho} \varepsilon_n.$$

Iterating this argument over l yields $\|\bar{v}^{(n,l)} - u\| \leq \tilde{\rho}^l \varepsilon_n$. In particular, by the choice of l^* , we have $\|v^{(n,l^*)} - u\| \leq \frac{1}{2KL} \vartheta \varepsilon_{n+1}$. From this it follows that

$$\|\tilde{\mathbf{u}}^{(n+1)} - \mathbf{Q}\mathbf{u}\|_{\ell_2(\Lambda)} \leq \frac{\vartheta}{2K} \varepsilon_{n+1} + \|\mathbf{Q}(\mathbf{v}^{(n,l^*)} - \mathbf{u})\|_{\ell_2(\Lambda)} \leq \frac{\vartheta}{2K} \varepsilon_{n+1} + L \frac{1}{2KL} \vartheta \varepsilon_{n+1} = \frac{1}{K} \vartheta \varepsilon_{n+1}. \quad (17)$$

Therefore, we have $\|u^{(n+1)} - u\| \leq K \|\tilde{\mathbf{u}}^{(n+1)} - \mathbf{Q}\mathbf{u}\|_{\ell_2(\Lambda)} \leq K \left(\frac{1-\vartheta}{K} \varepsilon_{n+1} + \|\tilde{\mathbf{u}}^{(n+1)} - \mathbf{Q}\mathbf{u}\|_{\ell_2(\Lambda)}\right) \leq \varepsilon_{n+1}$. \square

Now, the properties of the coarsening and projection methods allow us to show that the algorithm is asymptotically optimal with respect to the degrees of freedom in the outer iterates $\mathbf{u}^{(n)}$. By this, we mean that we obtain the same rate as the best N -term approximation of any representation $\mathbf{u} \in \ell_2(\Lambda)$ of u that has the aggregated tree structure.

Theorem 3.5 *Assume that the solution u has some representation $\mathbf{u} \in \mathcal{A}_{\mathcal{AT}}^s$. Then, for the iterates $\mathbf{u}^{(n)}$ from Algorithm 1, it holds that*

$$\begin{aligned} \mathbf{u}^{(n)} &\in \mathcal{A}_{\mathcal{AT}}^s, \\ \#\text{supp } \mathbf{u}^{(n)} &\lesssim \varepsilon_n^{-1/s} \|\mathbf{u}\|_{\mathcal{A}_{\mathcal{AT}}^s}^{1/s}. \end{aligned}$$

Proof. From (17) and the properties of **COARSE**, for all $n \in \mathbb{N}$, we have $\mathbf{u}^{(n+1)} \in \mathcal{A}_{\mathcal{AT}}^s$ and

$$\#\text{supp } \mathbf{u}^{(n+1)} \lesssim \left(\frac{\varepsilon_{n+1}}{K}\right)^{-1/s} \|\mathbf{Q}\mathbf{u}\|_{\mathcal{A}_{\mathcal{AT}}^s}^{1/s}.$$

Using boundedness of \mathbf{Q} on $\mathcal{A}_{\mathcal{AT}}^s$, we obtain the result. \square

3.3 How to solve the local subproblems

Let us now describe how the local subproblems appearing in Algorithm 1 can be solved. This can be done in the same fashion as in [10, 11], based on the Richardson method in [28]. The construction principles from there carry over to the vector-valued setting, compare, for instance, [29]. For convenience, we are going to sketch the algorithm here.

Written in coordinates, the subproblems amount to solving the equations

$$\mathbf{A}^{(i,i)} \mathbf{d}_i^{(n,l)} = -\text{Re } \mathbf{G}(\mathbf{v}^{(n,l)})|_{\Lambda_i} + \mathbf{f}|_{\Lambda_i} - \sum_{j=0}^{m-1} \mathbf{A}^{(i,j)} \mathbf{v}^{(n,l)},$$

where $\mathbf{A}^{(i,j)} := \{a(\psi_\lambda, \psi_\mu)\}_{\lambda \in \Lambda_i, \mu \in \Lambda_j}$ denotes the (i, j) -th block of the matrix \mathbf{A} , and, for any vector $\mathbf{w} \in \ell_2(\Lambda)$, by $\mathbf{w}|_{\Lambda_i}$ we mean the restriction of \mathbf{w} to the index set Λ_i . The matrix $\mathbf{A}^{(i,i)}$ is positive definite. Hence, for a sufficiently small relaxation parameter $\omega > 0$ and with \mathbf{R} being the representation of the residual of the subproblems, the Richardson iteration

$$\mathbf{w}^{(l+1)} = \mathbf{w}^{(l)} - \omega \mathbf{R}(\mathbf{w}^{(l)})$$

converges linearly for any initial vector $\mathbf{w}^{(0)}$. This is still true even if the residual is only approximated up to a given, sufficiently small tolerance. To do so, we have to make use of the vector-valued versions of the methods presented in [1, 2, 24, 29] for approximating the infinite matrix-vector products, the nonlinear term and the right-hand side. Moreover, as in [10, 11] we can see that in each call of the local solver the number of iterations to achieve the prescribed tolerance $\frac{1-\rho}{2m\omega} \varepsilon_n \tilde{\rho}^l$ is a constant independent of n . Therefore, also the computational complexity of the inner iterations can be bounded by a constant multiple of ε_n .

References

- [1] A. Cohen, W. Dahmen and R. DeVore, Adaptive wavelet methods for elliptic operator equations: Convergence rates, *Math. Comp.* 70 (2001), no. 233, 27–75
- [2] A. Cohen, W. Dahmen and R. DeVore, Adaptive wavelet schemes for nonlinear variational problems, *SIAM J. Numer. Anal.* 41 (2003), no. 5, 1785–1823

- [3] P.-G. Lemarié-Rieusset, Analyses multi-résolutions non orthogonales, commutation entre projecteurs et dérivations et ondelettes vecteurs à divergence nulle, *Revista Mat. Iberoamer.* 8 (1992), 221–237
- [4] R.P. Stevenson. Divergence-free wavelets on the hypercube: General boundary conditions. ESI preprint 2417, Erwin Schrödinger Institute, Vienna, April 2013. Submitted.
- [5] K. Urban, Using divergence free wavelets for the numerical solution of the Stokes problem, In *Algebraic Multilevel Iterations*, O. Axelsson, B. Polman (ed.), Nijmegen, 1996, 259–278.
- [6] K. Urban, *Wavelets in numerical simulation – Problem adapted construction and applications*, Lecture Notes in Computational Science and Engineering, Springer, 2002
- [7] R.P. Stevenson, Adaptive solution of operator equations using wavelet frames, *SIAM J. Numer. Anal.* 41 (2003), no. 3, 1074–1100
- [8] M. Dryja and W. Hackbusch, On the nonlinear domain decomposition method, *BIT* 37 (1997), no. 2, 296–311.
- [9] X.-C. Tai and J. Xu, Global and uniform convergence of subspace correction methods for some convex optimization problems, *Math. Comp.* 71 (2002), no. 237, 105–124.
- [10] R.P. Stevenson and M. Werner, A multiplicative Schwarz adaptive wavelet method for elliptic boundary value problems, *Math. Comp.* 78 (2009), no. 266, 619–644
- [11] D. Lellek, Adaptive wavelet frame domain decomposition methods for nonlinear elliptic equations, *Numer. Methods Partial Differential Equations* 29 (2013), 297–319
- [12] S.H. Lui, On Schwarz alternating methods for the incompressible Navier-Stokes equations, *SIAM J. Sci. Comp.* 22 (2001), 1974–1986
- [13] V. Girault and P.-A. Raviart, *Finite element approximation of the Navier-Stokes equations*, Springer, 1979
- [14] N. Faustino, U. Kähler and G. Teschke, A Wavelet-Galerkin scheme for the Navier-Stokes equations, *Preprint, Konrad-Zuse-Zentrum Berlin*, 2006
- [15] R. Temam, *Navier-Stokes equations. Theory and numerical analysis*, North-Holland Publishing Co., 1977.
- [16] O. Christensen. *Frames and bases – An introductory course*, Applied and Numerical Harmonic Analysis, Birkhäuser, Boston, 2008
- [17] P. L. Lions, On the Schwarz alternating method I, *First International Symposium on Domain Decomposition Methods for Partial Differential Equations (Paris, 1987)*, 1–42, SIAM, Philadelphia, 1988.
- [18] M. Werner, *Adaptive wavelet frame domain decomposition methods for elliptic operator equations*, Logos Verlag Berlin, 2009
- [19] S. Kadri Harouna and V. Perrier, Effective construction of divergence-free wavelets on the square, *J. Comput. Appl. Math.*, 240 (2013), 74–86
- [20] W. Dahmen and R. Schneider. Wavelets with complementary boundary conditions—function spaces on the cube. *Results Math.*, 34(3-4):255–293, 1998.

- [21] T.J. Dijkema. *Adaptive tensor product wavelet methods for solving PDEs*. PhD thesis, Utrecht University, 2009.
- [22] M. Primbs. New stable biorthogonal spline-wavelets on the interval. *Results Math.*, 57(1-2):121–162, 2010.
- [23] P.-G. Lemarié-Rieusset, Fonctions à support compact dans les analyses multi-résolutions, *Revista Mat. Iberoamer.* 7 (1991), 157–182
- [24] J. Kapeei, Adaptive frame methods for nonlinear elliptic problems, *Appl. Anal.* 90 (2011), 1323–1353.
- [25] S. Dahlke, Besov regularity for the Stokes problem, Haußmann, Werner (ed.) et al., Advances in multivariate approximation. Proceedings of the 3rd international conference on multivariate approximation theory, Witten-Bommerholz, Germany, September 27-October 2, 1998. Berlin: Wiley-VCH. Math. Res. 107, 129–138, 1999.
- [26] F. Eckhardt, Besov-regularity for the Stokes and the Navier-Stokes system in polyhedral domains, *Bericht Nr. 2013-02, Philipps-Universität Marburg*
- [27] J. Kapeei, *Adaptive wavelet frame methods for nonlinear elliptic problems*, Logos Verlag Berlin, 2011.
- [28] A. Cohen, W. Dahmen and R. DeVore, Adaptive wavelet methods. II. Beyond the elliptic case, *Found. Comput. Math.* 2 (2002), no. 3, 203–245.
- [29] Y. Jiang, Divergence-free wavelet solution to the Stokes problem, *Anal. Theory Appl.*, 23 (2007), no. 1, 83–91

Stephan Dahlke and Dominik Lellek
 Philipps-Universität Marburg
 Mathematics and Computer Science
 Hans-Meerwein-Straße, 35032 Marburg, Germany
 {dahlke, lellek}@mathematik.uni-marburg.de

Shiu Hong Lui
 University of Manitoba
 Department of Mathematics
 Winnipeg, Manitoba, Canada R3T 2N2
 luish@cc.umanitoba.ca

Rob Stevenson
 Korteweg-de Vries (KdV) Institute for Mathematics
 University of Amsterdam
 P.O. Box 94248, 1090 GE Amsterdam, The Netherlands
 R.P.Stevenson@uva.nl

Preprint Series DFG-SPP 1324

<http://www.dfg-spp1324.de>

Reports

- [1] R. Ramlau, G. Teschke, and M. Zhariy. A Compressive Landweber Iteration for Solving Ill-Posed Inverse Problems. Preprint 1, DFG-SPP 1324, September 2008.
- [2] G. Plonka. The Easy Path Wavelet Transform: A New Adaptive Wavelet Transform for Sparse Representation of Two-dimensional Data. Preprint 2, DFG-SPP 1324, September 2008.
- [3] E. Novak and H. Woźniakowski. Optimal Order of Convergence and (In-) Tractability of Multivariate Approximation of Smooth Functions. Preprint 3, DFG-SPP 1324, October 2008.
- [4] M. Espig, L. Grasedyck, and W. Hackbusch. Black Box Low Tensor Rank Approximation Using Fibre-Crosses. Preprint 4, DFG-SPP 1324, October 2008.
- [5] T. Bonesky, S. Dahlke, P. Maass, and T. Raasch. Adaptive Wavelet Methods and Sparsity Reconstruction for Inverse Heat Conduction Problems. Preprint 5, DFG-SPP 1324, January 2009.
- [6] E. Novak and H. Woźniakowski. Approximation of Infinitely Differentiable Multivariate Functions Is Intractable. Preprint 6, DFG-SPP 1324, January 2009.
- [7] J. Ma and G. Plonka. A Review of Curvelets and Recent Applications. Preprint 7, DFG-SPP 1324, February 2009.
- [8] L. Denis, D. A. Lorenz, and D. Trede. Greedy Solution of Ill-Posed Problems: Error Bounds and Exact Inversion. Preprint 8, DFG-SPP 1324, April 2009.
- [9] U. Friedrich. A Two Parameter Generalization of Lions' Nonoverlapping Domain Decomposition Method for Linear Elliptic PDEs. Preprint 9, DFG-SPP 1324, April 2009.
- [10] K. Bredies and D. A. Lorenz. Minimization of Non-smooth, Non-convex Functionals by Iterative Thresholding. Preprint 10, DFG-SPP 1324, April 2009.
- [11] K. Bredies and D. A. Lorenz. Regularization with Non-convex Separable Constraints. Preprint 11, DFG-SPP 1324, April 2009.

- [12] M. Döhler, S. Kunis, and D. Potts. Nonequispaced Hyperbolic Cross Fast Fourier Transform. Preprint 12, DFG-SPP 1324, April 2009.
- [13] C. Bender. Dual Pricing of Multi-Exercise Options under Volume Constraints. Preprint 13, DFG-SPP 1324, April 2009.
- [14] T. Müller-Gronbach and K. Ritter. Variable Subspace Sampling and Multi-level Algorithms. Preprint 14, DFG-SPP 1324, May 2009.
- [15] G. Plonka, S. Tenorth, and A. Iske. Optimally Sparse Image Representation by the Easy Path Wavelet Transform. Preprint 15, DFG-SPP 1324, May 2009.
- [16] S. Dahlke, E. Novak, and W. Sickel. Optimal Approximation of Elliptic Problems by Linear and Nonlinear Mappings IV: Errors in L_2 and Other Norms. Preprint 16, DFG-SPP 1324, June 2009.
- [17] B. Jin, T. Khan, P. Maass, and M. Pidcock. Function Spaces and Optimal Currents in Impedance Tomography. Preprint 17, DFG-SPP 1324, June 2009.
- [18] G. Plonka and J. Ma. Curvelet-Wavelet Regularized Split Bregman Iteration for Compressed Sensing. Preprint 18, DFG-SPP 1324, June 2009.
- [19] G. Teschke and C. Borries. Accelerated Projected Steepest Descent Method for Nonlinear Inverse Problems with Sparsity Constraints. Preprint 19, DFG-SPP 1324, July 2009.
- [20] L. Grasedyck. Hierarchical Singular Value Decomposition of Tensors. Preprint 20, DFG-SPP 1324, July 2009.
- [21] D. Rudolf. Error Bounds for Computing the Expectation by Markov Chain Monte Carlo. Preprint 21, DFG-SPP 1324, July 2009.
- [22] M. Hansen and W. Sickel. Best m -term Approximation and Lizorkin-Triebel Spaces. Preprint 22, DFG-SPP 1324, August 2009.
- [23] F.J. Hickernell, T. Müller-Gronbach, B. Niu, and K. Ritter. Multi-level Monte Carlo Algorithms for Infinite-dimensional Integration on \mathbb{R}^N . Preprint 23, DFG-SPP 1324, August 2009.
- [24] S. Dereich and F. Heidenreich. A Multilevel Monte Carlo Algorithm for Lévy Driven Stochastic Differential Equations. Preprint 24, DFG-SPP 1324, August 2009.
- [25] S. Dahlke, M. Fornasier, and T. Raasch. Multilevel Preconditioning for Adaptive Sparse Optimization. Preprint 25, DFG-SPP 1324, August 2009.

- [26] S. Dereich. Multilevel Monte Carlo Algorithms for Lévy-driven SDEs with Gaussian Correction. Preprint 26, DFG-SPP 1324, August 2009.
- [27] G. Plonka, S. Tenorth, and D. Roşca. A New Hybrid Method for Image Approximation using the Easy Path Wavelet Transform. Preprint 27, DFG-SPP 1324, October 2009.
- [28] O. Koch and C. Lubich. Dynamical Low-rank Approximation of Tensors. Preprint 28, DFG-SPP 1324, November 2009.
- [29] E. Faou, V. Gradinaru, and C. Lubich. Computing Semi-classical Quantum Dynamics with Hagedorn Wavepackets. Preprint 29, DFG-SPP 1324, November 2009.
- [30] D. Conte and C. Lubich. An Error Analysis of the Multi-configuration Time-dependent Hartree Method of Quantum Dynamics. Preprint 30, DFG-SPP 1324, November 2009.
- [31] C. E. Powell and E. Ullmann. Preconditioning Stochastic Galerkin Saddle Point Problems. Preprint 31, DFG-SPP 1324, November 2009.
- [32] O. G. Ernst and E. Ullmann. Stochastic Galerkin Matrices. Preprint 32, DFG-SPP 1324, November 2009.
- [33] F. Lindner and R. L. Schilling. Weak Order for the Discretization of the Stochastic Heat Equation Driven by Impulsive Noise. Preprint 33, DFG-SPP 1324, November 2009.
- [34] L. Kämmerer and S. Kunis. On the Stability of the Hyperbolic Cross Discrete Fourier Transform. Preprint 34, DFG-SPP 1324, December 2009.
- [35] P. Cerejeiras, M. Ferreira, U. Kähler, and G. Teschke. Inversion of the noisy Radon transform on $SO(3)$ by Gabor frames and sparse recovery principles. Preprint 35, DFG-SPP 1324, January 2010.
- [36] T. Jahnke and T. Udrescu. Solving Chemical Master Equations by Adaptive Wavelet Compression. Preprint 36, DFG-SPP 1324, January 2010.
- [37] P. Kittipoom, G. Kutyniok, and W.-Q. Lim. Irregular Shearlet Frames: Geometry and Approximation Properties. Preprint 37, DFG-SPP 1324, February 2010.
- [38] G. Kutyniok and W.-Q. Lim. Compactly Supported Shearlets are Optimally Sparse. Preprint 38, DFG-SPP 1324, February 2010.

- [39] M. Hansen and W. Sickel. Best m -Term Approximation and Tensor Products of Sobolev and Besov Spaces – the Case of Non-compact Embeddings. Preprint 39, DFG-SPP 1324, March 2010.
- [40] B. Niu, F.J. Hickernell, T. Müller-Gronbach, and K. Ritter. Deterministic Multi-level Algorithms for Infinite-dimensional Integration on $\mathbb{R}^{\mathbb{N}}$. Preprint 40, DFG-SPP 1324, March 2010.
- [41] P. Kittipoom, G. Kutyniok, and W.-Q. Lim. Construction of Compactly Supported Shearlet Frames. Preprint 41, DFG-SPP 1324, March 2010.
- [42] C. Bender and J. Steiner. Error Criteria for Numerical Solutions of Backward SDEs. Preprint 42, DFG-SPP 1324, April 2010.
- [43] L. Grasedyck. Polynomial Approximation in Hierarchical Tucker Format by Vector-Tensorization. Preprint 43, DFG-SPP 1324, April 2010.
- [44] M. Hansen und W. Sickel. Best m -Term Approximation and Sobolev-Besov Spaces of Dominating Mixed Smoothness - the Case of Compact Embeddings. Preprint 44, DFG-SPP 1324, April 2010.
- [45] P. Binev, W. Dahmen, and P. Lamby. Fast High-Dimensional Approximation with Sparse Occupancy Trees. Preprint 45, DFG-SPP 1324, May 2010.
- [46] J. Ballani and L. Grasedyck. A Projection Method to Solve Linear Systems in Tensor Format. Preprint 46, DFG-SPP 1324, May 2010.
- [47] P. Binev, A. Cohen, W. Dahmen, R. DeVore, G. Petrova, and P. Wojtaszczyk. Convergence Rates for Greedy Algorithms in Reduced Basis Methods. Preprint 47, DFG-SPP 1324, May 2010.
- [48] S. Kestler and K. Urban. Adaptive Wavelet Methods on Unbounded Domains. Preprint 48, DFG-SPP 1324, June 2010.
- [49] H. Yserentant. The Mixed Regularity of Electronic Wave Functions Multiplied by Explicit Correlation Factors. Preprint 49, DFG-SPP 1324, June 2010.
- [50] H. Yserentant. On the Complexity of the Electronic Schrödinger Equation. Preprint 50, DFG-SPP 1324, June 2010.
- [51] M. Guillemard and A. Iske. Curvature Analysis of Frequency Modulated Manifolds in Dimensionality Reduction. Preprint 51, DFG-SPP 1324, June 2010.
- [52] E. Herrholz and G. Teschke. Compressive Sensing Principles and Iterative Sparse Recovery for Inverse and Ill-Posed Problems. Preprint 52, DFG-SPP 1324, July 2010.

- [53] L. Kämmerer, S. Kunis, and D. Potts. Interpolation Lattices for Hyperbolic Cross Trigonometric Polynomials. Preprint 53, DFG-SPP 1324, July 2010.
- [54] G. Kutyniok and W.-Q Lim. Shearlets on Bounded Domains. Preprint 54, DFG-SPP 1324, July 2010.
- [55] A. Zeiser. Wavelet Approximation in Weighted Sobolev Spaces of Mixed Order with Applications to the Electronic Schrödinger Equation. Preprint 55, DFG-SPP 1324, July 2010.
- [56] G. Kutyniok, J. Lemvig, and W.-Q Lim. Compactly Supported Shearlets. Preprint 56, DFG-SPP 1324, July 2010.
- [57] A. Zeiser. On the Optimality of the Inexact Inverse Iteration Coupled with Adaptive Finite Element Methods. Preprint 57, DFG-SPP 1324, July 2010.
- [58] S. Jokar. Sparse Recovery and Kronecker Products. Preprint 58, DFG-SPP 1324, August 2010.
- [59] T. Aboiyar, E. H. Georgoulis, and A. Iske. Adaptive ADER Methods Using Kernel-Based Polyharmonic Spline WENO Reconstruction. Preprint 59, DFG-SPP 1324, August 2010.
- [60] O. G. Ernst, A. Mugler, H.-J. Starkloff, and E. Ullmann. On the Convergence of Generalized Polynomial Chaos Expansions. Preprint 60, DFG-SPP 1324, August 2010.
- [61] S. Holtz, T. Rohwedder, and R. Schneider. On Manifolds of Tensors of Fixed TT-Rank. Preprint 61, DFG-SPP 1324, September 2010.
- [62] J. Ballani, L. Grasedyck, and M. Kluge. Black Box Approximation of Tensors in Hierarchical Tucker Format. Preprint 62, DFG-SPP 1324, October 2010.
- [63] M. Hansen. On Tensor Products of Quasi-Banach Spaces. Preprint 63, DFG-SPP 1324, October 2010.
- [64] S. Dahlke, G. Steidl, and G. Teschke. Shearlet Coorbit Spaces: Compactly Supported Analyzing Shearlets, Traces and Embeddings. Preprint 64, DFG-SPP 1324, October 2010.
- [65] W. Hackbusch. Tensorisation of Vectors and their Efficient Convolution. Preprint 65, DFG-SPP 1324, November 2010.
- [66] P. A. Cioica, S. Dahlke, S. Kinzel, F. Lindner, T. Raasch, K. Ritter, and R. L. Schilling. Spatial Besov Regularity for Stochastic Partial Differential Equations on Lipschitz Domains. Preprint 66, DFG-SPP 1324, November 2010.

- [67] E. Novak and H. Woźniakowski. On the Power of Function Values for the Approximation Problem in Various Settings. Preprint 67, DFG-SPP 1324, November 2010.
- [68] A. Hinrichs, E. Novak, and H. Woźniakowski. The Curse of Dimensionality for Monotone and Convex Functions of Many Variables. Preprint 68, DFG-SPP 1324, November 2010.
- [69] G. Kutyniok and W.-Q. Lim. Image Separation Using Shearlets. Preprint 69, DFG-SPP 1324, November 2010.
- [70] B. Jin and P. Maass. An Analysis of Electrical Impedance Tomography with Applications to Tikhonov Regularization. Preprint 70, DFG-SPP 1324, December 2010.
- [71] S. Holtz, T. Rohwedder, and R. Schneider. The Alternating Linear Scheme for Tensor Optimisation in the TT Format. Preprint 71, DFG-SPP 1324, December 2010.
- [72] T. Müller-Gronbach and K. Ritter. A Local Refinement Strategy for Constructive Quantization of Scalar SDEs. Preprint 72, DFG-SPP 1324, December 2010.
- [73] T. Rohwedder and R. Schneider. An Analysis for the DIIS Acceleration Method used in Quantum Chemistry Calculations. Preprint 73, DFG-SPP 1324, December 2010.
- [74] C. Bender and J. Steiner. Least-Squares Monte Carlo for Backward SDEs. Preprint 74, DFG-SPP 1324, December 2010.
- [75] C. Bender. Primal and Dual Pricing of Multiple Exercise Options in Continuous Time. Preprint 75, DFG-SPP 1324, December 2010.
- [76] H. Harbrecht, M. Peters, and R. Schneider. On the Low-rank Approximation by the Pivoted Cholesky Decomposition. Preprint 76, DFG-SPP 1324, December 2010.
- [77] P. A. Cioica, S. Dahlke, N. Döhring, S. Kinzel, F. Lindner, T. Raasch, K. Ritter, and R. L. Schilling. Adaptive Wavelet Methods for Elliptic Stochastic Partial Differential Equations. Preprint 77, DFG-SPP 1324, January 2011.
- [78] G. Plonka, S. Tenorth, and A. Iske. Optimal Representation of Piecewise Hölder Smooth Bivariate Functions by the Easy Path Wavelet Transform. Preprint 78, DFG-SPP 1324, January 2011.

- [79] A. Mugler and H.-J. Starkloff. On Elliptic Partial Differential Equations with Random Coefficients. Preprint 79, DFG-SPP 1324, January 2011.
- [80] T. Müller-Gronbach, K. Ritter, and L. Yaroslavtseva. A Derandomization of the Euler Scheme for Scalar Stochastic Differential Equations. Preprint 80, DFG-SPP 1324, January 2011.
- [81] W. Dahmen, C. Huang, C. Schwab, and G. Welper. Adaptive Petrov-Galerkin methods for first order transport equations. Preprint 81, DFG-SPP 1324, January 2011.
- [82] K. Grella and C. Schwab. Sparse Tensor Spherical Harmonics Approximation in Radiative Transfer. Preprint 82, DFG-SPP 1324, January 2011.
- [83] D.A. Lorenz, S. Schiffler, and D. Tiede. Beyond Convergence Rates: Exact Inversion With Tikhonov Regularization With Sparsity Constraints. Preprint 83, DFG-SPP 1324, January 2011.
- [84] S. Dereich, M. Scheutzow, and R. Schottstedt. Constructive quantization: Approximation by empirical measures. Preprint 84, DFG-SPP 1324, January 2011.
- [85] S. Dahlke and W. Sickel. On Besov Regularity of Solutions to Nonlinear Elliptic Partial Differential Equations. Preprint 85, DFG-SPP 1324, January 2011.
- [86] S. Dahlke, U. Friedrich, P. Maass, T. Raasch, and R.A. Ressel. An adaptive wavelet method for parameter identification problems in parabolic partial differential equations. Preprint 86, DFG-SPP 1324, January 2011.
- [87] A. Cohen, W. Dahmen, and G. Welper. Adaptivity and Variational Stabilization for Convection-Diffusion Equations. Preprint 87, DFG-SPP 1324, January 2011.
- [88] T. Jahnke. On Reduced Models for the Chemical Master Equation. Preprint 88, DFG-SPP 1324, January 2011.
- [89] P. Binev, W. Dahmen, R. DeVore, P. Lamby, D. Savu, and R. Sharpley. Compressed Sensing and Electron Microscopy. Preprint 89, DFG-SPP 1324, March 2011.
- [90] P. Binev, F. Blanco-Silva, D. Blom, W. Dahmen, P. Lamby, R. Sharpley, and T. Vogt. High Quality Image Formation by Nonlocal Means Applied to High-Angle Annular Dark Field Scanning Transmission Electron Microscopy (HAADF-STEM). Preprint 90, DFG-SPP 1324, March 2011.
- [91] R. A. Ressel. A Parameter Identification Problem for a Nonlinear Parabolic Differential Equation. Preprint 91, DFG-SPP 1324, May 2011.

- [92] G. Kutyniok. Data Separation by Sparse Representations. Preprint 92, DFG-SPP 1324, May 2011.
- [93] M. A. Davenport, M. F. Duarte, Y. C. Eldar, and G. Kutyniok. Introduction to Compressed Sensing. Preprint 93, DFG-SPP 1324, May 2011.
- [94] H.-C. Kreuzler and H. Yserentant. The Mixed Regularity of Electronic Wave Functions in Fractional Order and Weighted Sobolev Spaces. Preprint 94, DFG-SPP 1324, June 2011.
- [95] E. Ullmann, H. C. Elman, and O. G. Ernst. Efficient Iterative Solvers for Stochastic Galerkin Discretizations of Log-Transformed Random Diffusion Problems. Preprint 95, DFG-SPP 1324, June 2011.
- [96] S. Kunis and I. Melzer. On the Butterfly Sparse Fourier Transform. Preprint 96, DFG-SPP 1324, June 2011.
- [97] T. Rohwedder. The Continuous Coupled Cluster Formulation for the Electronic Schrödinger Equation. Preprint 97, DFG-SPP 1324, June 2011.
- [98] T. Rohwedder and R. Schneider. Error Estimates for the Coupled Cluster Method. Preprint 98, DFG-SPP 1324, June 2011.
- [99] P. A. Cioica and S. Dahlke. Spatial Besov Regularity for Semilinear Stochastic Partial Differential Equations on Bounded Lipschitz Domains. Preprint 99, DFG-SPP 1324, July 2011.
- [100] L. Grasedyck and W. Hackbusch. An Introduction to Hierarchical (H-) Rank and TT-Rank of Tensors with Examples. Preprint 100, DFG-SPP 1324, August 2011.
- [101] N. Chegini, S. Dahlke, U. Friedrich, and R. Stevenson. Piecewise Tensor Product Wavelet Bases by Extensions and Approximation Rates. Preprint 101, DFG-SPP 1324, September 2011.
- [102] S. Dahlke, P. Oswald, and T. Raasch. A Note on Quarkonial Systems and Multi-level Partition of Unity Methods. Preprint 102, DFG-SPP 1324, September 2011.
- [103] A. Uschmajew. Local Convergence of the Alternating Least Squares Algorithm For Canonical Tensor Approximation. Preprint 103, DFG-SPP 1324, September 2011.
- [104] S. Kvaal. Multiconfigurational time-dependent Hartree method for describing particle loss due to absorbing boundary conditions. Preprint 104, DFG-SPP 1324, September 2011.

- [105] M. Guillemard and A. Iske. On Groupoid C^* -Algebras, Persistent Homology and Time-Frequency Analysis. Preprint 105, DFG-SPP 1324, September 2011.
- [106] A. Hinrichs, E. Novak, and H. Woźniakowski. Discontinuous information in the worst case and randomized settings. Preprint 106, DFG-SPP 1324, September 2011.
- [107] M. Espig, W. Hackbusch, A. Litvinenko, H. Matthies, and E. Zander. Efficient Analysis of High Dimensional Data in Tensor Formats. Preprint 107, DFG-SPP 1324, September 2011.
- [108] M. Espig, W. Hackbusch, S. Handschuh, and R. Schneider. Optimization Problems in Contracted Tensor Networks. Preprint 108, DFG-SPP 1324, October 2011.
- [109] S. Dereich, T. Müller-Gronbach, and K. Ritter. On the Complexity of Computing Quadrature Formulas for SDEs. Preprint 109, DFG-SPP 1324, October 2011.
- [110] D. Belomestny. Solving optimal stopping problems by empirical dual optimization and penalization. Preprint 110, DFG-SPP 1324, November 2011.
- [111] D. Belomestny and J. Schoenmakers. Multilevel dual approach for pricing American style derivatives. Preprint 111, DFG-SPP 1324, November 2011.
- [112] T. Rohwedder and A. Uschmajew. Local convergence of alternating schemes for optimization of convex problems in the TT format. Preprint 112, DFG-SPP 1324, December 2011.
- [113] T. Görner, R. Hielscher, and S. Kunis. Efficient and accurate computation of spherical mean values at scattered center points. Preprint 113, DFG-SPP 1324, December 2011.
- [114] Y. Dong, T. Görner, and S. Kunis. An iterative reconstruction scheme for photoacoustic imaging. Preprint 114, DFG-SPP 1324, December 2011.
- [115] L. Kämmerer. Reconstructing hyperbolic cross trigonometric polynomials by sampling along generated sets. Preprint 115, DFG-SPP 1324, February 2012.
- [116] H. Chen and R. Schneider. Numerical analysis of augmented plane waves methods for full-potential electronic structure calculations. Preprint 116, DFG-SPP 1324, February 2012.
- [117] J. Ma, G. Plonka, and M.Y. Hussaini. Compressive Video Sampling with Approximate Message Passing Decoding. Preprint 117, DFG-SPP 1324, February 2012.

- [118] D. Heinen and G. Plonka. Wavelet shrinkage on paths for scattered data denoising. Preprint 118, DFG-SPP 1324, February 2012.
- [119] T. Jahnke and M. Kreim. Error bound for piecewise deterministic processes modeling stochastic reaction systems. Preprint 119, DFG-SPP 1324, March 2012.
- [120] C. Bender and J. Steiner. A-posteriori estimates for backward SDEs. Preprint 120, DFG-SPP 1324, April 2012.
- [121] M. Espig, W. Hackbusch, A. Litvinenkoy, H.G. Matthiesy, and P. Wähnert. Efficient low-rank approximation of the stochastic Galerkin matrix in tensor formats. Preprint 121, DFG-SPP 1324, May 2012.
- [122] O. Bokanowski, J. Garcke, M. Griebel, and I. Klompmaker. An adaptive sparse grid semi-Lagrangian scheme for first order Hamilton-Jacobi Bellman equations. Preprint 122, DFG-SPP 1324, June 2012.
- [123] A. Mugler and H.-J. Starkloff. On the convergence of the stochastic Galerkin method for random elliptic partial differential equations. Preprint 123, DFG-SPP 1324, June 2012.
- [124] P.A. Cioica, S. Dahlke, N. Döhning, U. Friedrich, S. Kinzel, F. Lindner, T. Raasch, K. Ritter, and R.L. Schilling. On the convergence analysis of Rothe's method. Preprint 124, DFG-SPP 1324, July 2012.
- [125] P. Binev, A. Cohen, W. Dahmen, and R. DeVore. Classification Algorithms using Adaptive Partitioning. Preprint 125, DFG-SPP 1324, July 2012.
- [126] C. Lubich, T. Rohwedder, R. Schneider, and B. Vandereycken. Dynamical approximation of hierarchical Tucker and Tensor-Train tensors. Preprint 126, DFG-SPP 1324, July 2012.
- [127] M. Kovács, S. Larsson, and K. Urban. On Wavelet-Galerkin methods for semilinear parabolic equations with additive noise. Preprint 127, DFG-SPP 1324, August 2012.
- [128] M. Bachmayr, H. Chen, and R. Schneider. Numerical analysis of Gaussian approximations in quantum chemistry. Preprint 128, DFG-SPP 1324, August 2012.
- [129] D. Rudolf. Explicit error bounds for Markov chain Monte Carlo. Preprint 129, DFG-SPP 1324, August 2012.
- [130] P.A. Cioica, K.-H. Kim, K. Lee, and F. Lindner. On the $L_q(L_p)$ -regularity and Besov smoothness of stochastic parabolic equations on bounded Lipschitz domains. Preprint 130, DFG-SPP 1324, December 2012.

- [131] M. Hansen. n -term Approximation Rates and Besov Regularity for Elliptic PDEs on Polyhedral Domains. Preprint 131, DFG-SPP 1324, December 2012.
- [132] R. E. Bank and H. Yserentant. On the H^1 -stability of the L_2 -projection onto finite element spaces. Preprint 132, DFG-SPP 1324, December 2012.
- [133] M. Gnewuch, S. Mayer, and K. Ritter. On Weighted Hilbert Spaces and Integration of Functions of Infinitely Many Variables. Preprint 133, DFG-SPP 1324, December 2012.
- [134] D. Crisan, J. Diehl, P.K. Friz, and H. Oberhauser. Robust Filtering: Correlated Noise and Multidimensional Observation. Preprint 134, DFG-SPP 1324, January 2013.
- [135] Wolfgang Dahmen, Christian Plesken, and Gerrit Welper. Double Greedy Algorithms: Reduced Basis Methods for Transport Dominated Problems. Preprint 135, DFG-SPP 1324, February 2013.
- [136] Aicke Hinrichs, Erich Novak, Mario Ullrich, and Henryk Wozniakowski. The Curse of Dimensionality for Numerical Integration of Smooth Functions. Preprint 136, DFG-SPP 1324, February 2013.
- [137] Markus Bachmayr, Wolfgang Dahmen, Ronald DeVore, and Lars Grasedyck. Approximation of High-Dimensional Rank One Tensors. Preprint 137, DFG-SPP 1324, March 2013.
- [138] Markus Bachmayr and Wolfgang Dahmen. Adaptive Near-Optimal Rank Tensor Approximation for High-Dimensional Operator Equations. Preprint 138, DFG-SPP 1324, April 2013.
- [139] Felix Lindner. Singular Behavior of the Solution to the Stochastic Heat Equation on a Polygonal Domain. Preprint 139, DFG-SPP 1324, May 2013.
- [140] Stephan Dahlke, Dominik Lellek, Shiu Hong Lui, and Rob Stevenson. Adaptive Wavelet Schwarz Methods for the Navier-Stokes Equation. Preprint 140, DFG-SPP 1324, May 2013.