


DFG-Schwerpunktprogramm 1324

„Extraktion quantifizierbarer Information aus komplexen Systemen“

Approximation of multivariate periodic functions by
trigonometric polynomials based on sampling along
rank-1 lattice with generating vector of Korobov
form

L. Kämmerer, D. Potts, T. Volkmer

Preprint 159



Edited by

AG Numerik/Optimierung
Fachbereich 12 - Mathematik und Informatik
Philipps-Universität Marburg
Hans-Meerwein-Str.
35032 Marburg

DFG-Schwerpunktprogramm 1324

„Extraktion quantifizierbarer Information aus komplexen Systemen“

Approximation of multivariate periodic functions by
trigonometric polynomials based on sampling along
rank-1 lattice with generating vector of Korobov
form

L. Kämmerer, D. Potts, T. Volkmer

Preprint 159



The consecutive numbering of the publications is determined by their chronological order.

The aim of this preprint series is to make new research rapidly available for scientific discussion. Therefore, the responsibility for the contents is solely due to the authors. The publications will be distributed by the authors.

Approximation of multivariate periodic functions by trigonometric polynomials based on sampling along rank-1 lattice with generating vector of Korobov form

Lutz Kämmerer* Daniel Potts* Toni Volkmer*

In this paper, we present error estimates for the approximation of multivariate periodic functions in periodic Hilbert spaces of isotropic and dominating mixed smoothness by trigonometric polynomials. The approximation is based on sampling of the multivariate functions on rank-1 lattices. We use reconstructing rank-1 lattices with generating vectors of Korobov form for the sampling and generalize the technique from [24], in order to show that the aliasing error of that approximation is of the same order as the error of the approximation using the partial sum of the Fourier series. The main advantage of our method is that the computation of the Fourier coefficients of such a trigonometric polynomial, which we use as approximant, is based mainly on a one-dimensional fast Fourier transform, i.e., the arithmetic complexity of the computation depends only on the cardinality of the support of the trigonometric polynomial in the frequency domain. Numerical results are presented up to dimension $d = 10$.

Keywords and phrases : approximation of multivariate functions, trigonometric polynomials, hyperbolic cross, lattice rule, rank-1 lattice, fast Fourier transform

2000 AMS Mathematics Subject Classification : 65T40, 42A10,

*Technische Universität Chemnitz, Faculty of Mathematics, 09107 Chemnitz, Germany
{lutz.kaemmerer, daniel.potts, toni.volkmer}@mathematik.tu-chemnitz.de

1 Introduction

We approximate functions $f \in \mathcal{H}^\omega(\mathbb{T}^d)$ from the Hilbert space

$$\mathcal{H}^\omega(\mathbb{T}^d) := \left\{ f \in C(\mathbb{T}^d) : \|f\|_{\mathcal{H}^\omega(\mathbb{T}^d)} := \sqrt{\sum_{\mathbf{k} \in \mathbb{Z}^d} \omega(\mathbf{k})^2 |\hat{f}_{\mathbf{k}}|^2} < \infty \right\},$$

where $\omega: \mathbb{Z}^d \rightarrow (0, \infty]$ is a weight function, by trigonometric polynomials p with frequencies supported on an index set $I \subset \mathbb{Z}^d$ of finite cardinality, $p(\mathbf{x}) := \sum_{\mathbf{k} \in I} \hat{p}_{\mathbf{k}} e^{2\pi i \mathbf{k} \mathbf{x}}$. Thereby, we are especially interested in the higher-dimensional cases, i.e., $d \geq 4$. As usual, we denote the Fourier coefficients of the function $f \in L^2(\mathbb{T}^d)$ by

$$\hat{f}_{\mathbf{k}} := \int_{\mathbb{T}^d} f(\mathbf{x}) e^{-2\pi i \mathbf{k} \mathbf{x}} d\mathbf{x}, \quad \mathbf{k} \in \mathbb{Z}^d.$$

We remark that for the special choice $\omega \equiv 1$, we have $\mathcal{H}^\omega(\mathbb{T}^d) = C(\mathbb{T}^d) \cap L^2(\mathbb{T}^d)$. One theoretical possibility to obtain such a trigonometric polynomial p is to formally approximate the function f by the Fourier partial sum

$$S_I f := \sum_{\mathbf{k} \in I} \hat{f}_{\mathbf{k}} e^{2\pi i \mathbf{k} \mathbf{x}},$$

where $I \subset \mathbb{Z}^d$ is a frequency index set of finite cardinality. Since $S_I f$ is the truncated Fourier partial sum of the function f , this approximation causes a truncation error $\|f - S_I f\|$, where $\|\cdot\|$ is an arbitrarily chosen norm. For a function $f \in L^2(\mathbb{T}^d) \cap \mathcal{H}^\omega(\mathbb{T}^d)$ we choose a frequency index set $I = I_N := \{\mathbf{k} \in \mathbb{Z}^d : \omega(\mathbf{k})^{1/\nu} \leq N\}$ of refinement $N \in \mathbb{R}$, $N \geq 1$, $\nu > 0$, and obtain

$$\|f - S_{I_N} f\|_{L^2(\mathbb{T}^d)} \leq N^{-\nu} \|f\|_{\mathcal{H}^\omega(\mathbb{T}^d)},$$

see Lemma 3.3. We stress the fact that $S_{I_N} f$ is the best approximation of the function f with respect to the $L^2(\mathbb{T}^d)$ norm in the space $\Pi_{I_N} := \text{span}\{e^{2\pi i \mathbf{k} \mathbf{x}} : \mathbf{k} \in I_N\}$ of trigonometric polynomials with frequencies supported on the index set I_N and that the operator $S_{I_N}: L^1(\mathbb{T}^d) \rightarrow \Pi_{I_N}$ only depends on the frequency index set I_N .

Since, in general, we do not know the Fourier coefficients $\hat{f}_{\mathbf{k}}$, we are going to approximate the function f from samples using the approximated Fourier partial sum

$$\tilde{S}_{I_N} f := \sum_{\mathbf{k} \in I_N} \tilde{\hat{f}}_{\mathbf{k}} e^{2\pi i \mathbf{k} \mathbf{x}}.$$

We compute the approximated Fourier coefficients $\tilde{\hat{f}}_{\mathbf{k}} \in \mathbb{C}$, $\mathbf{k} \in I_N$, of the function f using a lattice rule by

$$\tilde{\hat{f}}_{\mathbf{k}} := \frac{1}{M} \sum_{j=0}^{M-1} f(\mathbf{x}_j) e^{-2\pi i \mathbf{k} \mathbf{x}_j} \quad \text{for } \mathbf{k} \in I_N, \quad (1.1)$$

where the sampling nodes $\mathbf{x}_j := \frac{j}{M} \mathbf{z} \bmod 1$ are the nodes of a so-called reconstructing rank-1 lattice $\Lambda(\mathbf{z}, M, I_N)$ with generating vector $\mathbf{z} \in \mathbb{Z}^d$ and rank-1 lattice size $M \in \mathbb{N}$ for the frequency index set I_N , see Section 2.2 for the definition. Lattice rules have extensively been investigated for the integration of functions of many variables for a long time, cf. e.g., [22, 4, 5]

and the extensive reference list therein. Especially, rank-1 lattice rules have also been studied for the approximation of multivariate functions of suitable smoothness, cf. [24, 18, 17, 19]. Furthermore, there exist already comprehensive tractability results for numerical integration and approximation using rank-1 lattices, see [20, 17].

Since we consider the partial sum $\tilde{S}_{I_N} f$ of the approximated Fourier coefficients $\tilde{f}_{\mathbf{k}}$ instead of the Fourier partial sum $S_{I_N} f$ of Fourier coefficients $\hat{f}_{\mathbf{k}}$, we obtain an additional error. We estimate the approximation error $\|f - \tilde{S}_{I_N} f\|_{L^2(\mathbb{T}^d)}$ using the triangle inequality $\|f - \tilde{S}_{I_N} f\|_{L^2(\mathbb{T}^d)} \leq \|f - S_{I_N} f\|_{L^2(\mathbb{T}^d)} + \|S_{I_N} f - \tilde{S}_{I_N} f\|_{L^2(\mathbb{T}^d)}$, where $\|f - S_{I_N} f\|_{L^2(\mathbb{T}^d)}$ is called the truncation error and $\|S_{I_N} f - \tilde{S}_{I_N} f\|_{L^2(\mathbb{T}^d)}$ is called the aliasing error.

In this paper, we consider frequency index sets I_N of special structure and show that there exists a reconstructing rank-1 lattice $\Lambda(\mathbf{z}, M, I_N)$ of reasonable size M , see Section 2.2 for the definition, such that the order of the aliasing error $\|S_{I_N} f - \tilde{S}_{I_N} f\|_{L^2(\mathbb{T}^d)}$ is bounded by the order of the truncation error $\|f - S_{I_N} f\|_{L^2(\mathbb{T}^d)}$. To this end, we use the highly structured rank-1 lattice rules with generating vector of Korobov form. This allows us to generalize the ideas of V. N. Temlyakov, see [24], in order to estimate the aliasing error. We consider, similar to [7] and as in [14], functions f from the Hilbert space

$$\mathcal{H}^\omega(\mathbb{T}^d) = \mathcal{H}^{\alpha, \beta}(\mathbb{T}^d) := \left\{ f \in C(\mathbb{T}^d) : \|f\|_{\mathcal{H}^{\alpha, \beta}(\mathbb{T}^d)} := \sqrt{\sum_{\mathbf{k} \in \mathbb{Z}^d} \omega^{\alpha, \beta}(\mathbf{k})^2 |\hat{f}_{\mathbf{k}}|^2} < \infty \right\},$$

where the parameter $\beta \in \mathbb{R}$, $\beta \geq 0$, characterizes the dominating mixed smoothness, the parameter $\alpha \in \mathbb{R}$, $\alpha > -\beta$, characterizes the isotropic smoothness, and the weights $\omega(\mathbf{k}) = \omega^{\alpha, \beta}(\mathbf{k})$ are given by

$$\omega^{\alpha, \beta}(\mathbf{k}) := \max(1, \|\mathbf{k}\|_1)^\alpha \prod_{s=1}^d \max(1, |k_s|)^\beta, \quad \mathbf{k} := (k_1, \dots, k_d)^\top.$$

We remark that one can use various equivalent weights $\omega(\mathbf{k})$ which have different approximation properties for large dimensions d , cf. [16]. Furthermore, we define the corresponding frequency index sets $I_N = I_N^{d, T}$, $N \in \mathbb{R}$, $N \geq 1$, $T \in \mathbb{R}$, $-\infty < T < 1$, by

$$I_N^{d, T} := \left\{ \mathbf{k} \in \mathbb{Z}^d : \omega^{-\frac{T}{1-T}, \frac{1}{1-T}}(\mathbf{k}) = \max(1, \|\mathbf{k}\|_1)^{-\frac{T}{1-T}} \prod_{s=1}^d \max(1, |k_s|)^{\frac{1}{1-T}} \leq N \right\}.$$

In the cases $0 < T < 1$, the frequency index sets $I_N^{d, T}$ are called energy-norm based hyperbolic crosses, see [2, 3], and in the case $T = 0$ symmetric hyperbolic crosses. As a natural extension for $T = -\infty$, we define the frequency index set $I_N^{d, -\infty}$ as the d -dimensional ℓ_1 -ball of size N ,

$$I_N^{d, -\infty} := \left\{ \mathbf{k} \in \mathbb{Z}^d : \max(1, \|\mathbf{k}\|_1) \leq N \right\}.$$

Figure 1.1 illustrates the frequency index sets $I_N^{d, T}$ in the two-dimensional case for different choices $-\infty \leq T < 1$ of the parameter T . The cardinalities of the frequency index sets $I_N^{d, T}$ are given in Lemma 4.1, which reads for fixed $d \in \mathbb{N}$ and $T := -\alpha/\beta$ as follows

$$\left| I_N^{d, -\alpha/\beta} \right| = \begin{cases} \Theta(N^d) & \text{for } \alpha > 0 \text{ and } \beta = 0 \quad (\iff T = -\infty), \\ \Theta(N^{d \frac{\beta + \alpha}{d\beta + \alpha}}) & \text{for } \alpha > 0 \text{ and } \beta > 0 \quad (\iff -\infty < T < 0), \\ \Theta(N \log^{d-1} N) & \text{for } \alpha = 0 \text{ and } \beta > 0 \quad (\iff T = 0), \\ \Theta(N) & \text{for } \alpha < 0 \text{ and } \beta > -\alpha \quad (\iff 0 < T < 1). \end{cases} \quad (1.2)$$




Figure 1.1: Two-dimensional frequency index sets $I_{32}^{2,T}$ for $T \in \{-\infty, -5, 0, \frac{1}{2}\}$.




Figure 1.2: Visualization of the admissible values of α and β in the case $d = 2$, such that (1.3) and (1.4) are valid. We set the corresponding values $T := -\alpha/\beta$.

In this setting, we obtain that the $L^2(\mathbb{T}^d)$ truncation error is bounded by

$$\|f - S_{I_N^{d,-\alpha/\beta}} f\|_{L^2(\mathbb{T}^d)} \leq N^{-(\alpha+\beta)} \|f\|_{\mathcal{H}^{\alpha,\beta}(\mathbb{T}^d)},$$

see Lemma 4.4. The main result of this paper is, that for fixed dimension $d \in \mathbb{N}$, $d \geq 2$ there exists a reconstructing rank-1 lattice $\Lambda(\mathbf{z}, M, I_N^{d,-\alpha/\beta})$ with generating vector $\mathbf{z} := (1, a, \dots, a^{d-1})^\top \in \mathbb{Z}^d$ of Korobov form and size

$$M = \begin{cases} \mathcal{O}(N^d) & \text{for } \alpha > \frac{d}{2} \text{ and } \beta = 0, \\ \mathcal{O}(N^{\frac{2d\beta+\alpha(\beta+\alpha)}{(d\beta+\alpha)^2}}) & \text{for } \beta > 0 \text{ and } \alpha > \max(0, (\frac{1}{4} - \beta + \frac{1}{4}\sqrt{8\beta+1})d), \\ \mathcal{O}(N^2 \log^{d-1} N) & \text{for } \alpha = 0 \text{ and } \beta > 1, \\ \mathcal{O}(N^2) & \text{for } \alpha < 0 \text{ and } \beta > 1 - \alpha, \end{cases} \quad (1.3)$$

such that the aliasing error is bounded by

$$\|S_{I_N^{d,-\alpha/\beta}} f - \tilde{S}_{I_N^{d,-\alpha/\beta}} f\|_{L^2(\mathbb{T}^d)} \leq C(d, \alpha, \beta) N^{-(\alpha+\beta)} \|f\|_{\mathcal{H}^{\alpha,\beta}(\mathbb{T}^d)}, \quad (1.4)$$

where $C(d, \alpha, \beta) > 0$ is a constant which only depends on d , α , β . In the cases where $\alpha \geq 0$, we obtain estimate (1.4) from Theorem 4.7 and the lower bound for the size M in (1.3) due to (4.4) and (4.1). For $\alpha < 0$, we infer estimate (1.4) from Theorem 4.10 and the lower bound for the size M in (1.3) due to (4.5) and (4.1). Figure 1.2 visualizes the

different cases for the admissible values of the isotropic smoothness α and the dominating mixed smoothness β in (1.3) and (1.4) in the two-dimensional case and gives the corresponding values of the parameter T . In Figure 1.3, the admissible values of α and β are shown for the cases $d = 2, 6, 10$. Comparing the number M of sampling nodes \mathbf{x}_j in (1.3) and the




Figure 1.3: Visualization of the admissible values of α and β in the cases $d = 2, 6, 10$, such that (1.3) and (1.4) are valid.

number $|I_N^{d,-\alpha/\beta}|$ of frequency indices in (1.2), our results yield in general an oversampling, i.e., $M > |I_N^{d,-\alpha/\beta}|$. In the case $\alpha > \frac{d}{2}$ and $\beta = 0$, where the frequency index sets $I_N^{d,-\infty}$ are l_1 -balls, the asymptotic order of M and $|I_N^{d,-\infty}|$ in N is obviously identical. Considering the case $\alpha < 0$ and $\beta > 1 - \alpha$, where the frequency index sets $I_N^{d,-\alpha/\beta}$ are energy-norm based hyperbolic crosses, we obtain a gap between M and $|I_N^{d,-\alpha/\beta}|$ in the asymptotic order in N . However, this gap is necessary in order to obtain an orthogonal Fourier transform as given by (1.1), cf. [10, Lemma 2.1]. Note that in the case $\alpha = 0$, the oversampling factors $M/|I_N^{d,0}|$, i.e., ratios of the rank-1 lattice sizes M and the cardinalities of the symmetric hyperbolic cross index sets $I_N^{d,0}$ are still moderate for reasonable problem sizes compared to the asymptotic statement $\mathcal{O}(N)$ in (1.3) and (1.2), see Table 5.1.

Let us mention that sampling on (generalized) sparse grids, see [25, 1, 30, 8, 26, 2, 23, 3, 6, 21, 9, 7], is another intensively studied approach used to approximate functions of the classes $\mathcal{H}^{\alpha,\beta}(\mathbb{T}^d)$. One advantage of this method is that only $|I_N^{d,-\alpha/\beta}|$ many samples are required. Furthermore, for $\alpha = 0$, there exists a fast algorithm for computing the approximation of the Fourier partial sum $S_{I_N^{d,0}}$ of a function f in $\mathcal{O}(N \log^d N)$ arithmetic operations. However, the computation may be numerically unstable in this setting, cf. [12]. Known upper bounds for the approximation errors are discussed in Section 4.3. We stress again, that the outstanding property of the sampling method (1.1) discussed in this paper is that the computation of the approximated Fourier coefficients $\hat{f}_{\mathbf{k}}, \mathbf{k} \in I_N^{d,0}$, is perfectly stable and takes $\mathcal{O}(N^2 \log^d N)$ arithmetic operations, since it is mainly based on a one-dimensional fast Fourier transform (FFT), cf. [13] and [14, Algorithm 3].

The paper is organized as follows: We discuss the exact reconstruction of trigonometric polynomials from samples along a rank-1 lattice in Section 2 and prove the existence of a special rank-1 lattice with certain properties. Based on these special properties, we show general estimates for the aliasing error for general frequency index sets I_N in Section 3. Then, in Section 4, we consider the approximation error $\|f - \tilde{S}_{I_N^{d,-\alpha/\beta}} f\|_{L^2(\mathbb{T}^d)}$. Therefore,

we present the estimates for the truncation error in Section 4.1. In Section 4.2, we prove the results (1.3) and (1.4). We compare these results with previously known ones in Section 4.3. Finally, we present numerical tests in Section 5 in order to illustrate the theoretical results and we give some concluding remarks in Section 6.

2 Approximation based on rank-1 lattice sampling

2.1 Reconstruction of trigonometric polynomials from samples

As already discussed in Section 1, we want to approximate a function $f \in \mathcal{H}^\omega(\mathbb{T}^d)$ using a trigonometric polynomial p . Here, we use the following approach. For a given frequency index set $I \subset \mathbb{Z}^d$ of finite cardinality, we want to be able to exactly reconstruct the Fourier coefficients $\hat{p}_{\mathbf{k}}$, $\mathbf{k} \in I$, of an arbitrarily chosen trigonometric polynomial $p(\mathbf{x}) := \sum_{\mathbf{k} \in I} \hat{p}_{\mathbf{k}} e^{2\pi i \mathbf{k} \mathbf{x}}$ with frequencies supported on I from sampling values $p(\mathbf{x}_j)$. As sampling nodes \mathbf{x}_j , $j = 0, \dots, M-1$, we use the nodes of a rank-1 lattice $\Lambda(\mathbf{z}, M)$ with generating vector $\mathbf{z} \in \mathbb{Z}^d$ of size $M \in \mathbb{N}$, i.e., we set $\mathbf{x}_j := \frac{j}{M} \mathbf{z} \bmod 1$, $j = 0, \dots, M-1$. Formally, the Fourier coefficients $\hat{p}_{\mathbf{k}}$ are given by

$$\hat{p}_{\mathbf{k}} := \int_{\mathbb{T}^d} p(\mathbf{x}) e^{-2\pi i \mathbf{k} \mathbf{x}} d\mathbf{x}$$

and we approximate this integral by the (rank-1) lattice rule

$$\frac{1}{M} \sum_{j=0}^{M-1} p(\mathbf{x}_j) e^{-2\pi i \mathbf{k} \mathbf{x}_j} = \frac{1}{M} \sum_{j=0}^{M-1} p\left(\frac{j}{M} \mathbf{z}\right) e^{-2\pi i j \mathbf{k} \mathbf{z} / M} =: \hat{\hat{p}}_{\mathbf{k}}.$$

Now, we ask for the exactness of this cubature formula, i.e., when is $\hat{\hat{p}}_{\mathbf{k}} = \hat{p}_{\mathbf{k}} \forall \mathbf{k} \in I$. Since we have

$$\hat{\hat{p}}_{\mathbf{k}} = \frac{1}{M} \sum_{j=0}^{M-1} \sum_{\mathbf{k}' \in I} \hat{p}_{\mathbf{k}'} e^{2\pi i j \mathbf{k}' \mathbf{z} / M} e^{-2\pi i j \mathbf{k} \mathbf{z} / M} = \sum_{\mathbf{k}' \in I} \hat{p}_{\mathbf{k}'} \frac{1}{M} \sum_{j=0}^{M-1} e^{2\pi i j (\mathbf{k}' - \mathbf{k}) \mathbf{z} / M},$$

we need the condition

$$\frac{1}{M} \sum_{j=0}^{M-1} e^{2\pi i j (\mathbf{k}' - \mathbf{k}) \mathbf{z} / M} = \begin{cases} 1 & \text{for } \mathbf{k} = \mathbf{k}' \\ 0 & \text{for } \mathbf{k} \neq \mathbf{k}', \mathbf{k}, \mathbf{k}' \in I, \end{cases} \quad (2.1)$$

to be fulfilled. This is the case if and only if

$$(\mathbf{k}' - \mathbf{k}) \mathbf{z} \not\equiv 0 \pmod{M} \quad \forall \mathbf{k}, \mathbf{k}' \in I, \mathbf{k} \neq \mathbf{k}', \quad (2.2)$$

$$\iff \mathbf{k} \cdot \mathbf{z} \not\equiv \mathbf{k}' \cdot \mathbf{z} \pmod{M} \quad \forall \mathbf{k}, \mathbf{k}' \in I, \mathbf{k} \neq \mathbf{k}', \quad (2.3)$$

see [11, Section 2]. Introducing the difference set $\mathcal{D}(I)$ for the index set I , $\mathcal{D}(I) := \{\mathbf{k} - \mathbf{k}' : \mathbf{k}, \mathbf{k}' \in I\}$, we can rewrite the above conditions to

$$m \mathbf{z} \not\equiv 0 \pmod{M} \quad \forall m \in \mathcal{D}(I) \setminus \{\mathbf{0}\}. \quad (2.4)$$

2.2 Reconstructing rank-1 lattices

A rank-1 lattice $\Lambda(\mathbf{z}, M)$ which fulfills one of the (equivalent) conditions (2.1),(2.2),(2.3),(2.4) for a given frequency index set I will be called reconstructing rank-1 lattice $\Lambda(\mathbf{z}, M, I)$. Under mild assumptions, e.g., $I \subset \mathbb{Z}^d \cap (-M/2, M/2)^d$, there always exists a reconstructing rank-1 lattice $\Lambda(\mathbf{z}, M, I)$ of size $\frac{|\mathcal{D}(I)|}{2} \leq M \leq |\mathcal{D}(I)|$ due to [11, Corollary 1] and Bertrand's postulate. We remark that we can compute the approximated Fourier coefficients $\tilde{f}_{\mathbf{k}}, \mathbf{k} \in I$, from (1.1) in $\mathcal{O}(M \log M + d|I|)$ arithmetic operations using a single one-dimensional fast Fourier transform of length M and by computing the scalar products $\mathbf{k}\mathbf{z}$ for $\mathbf{k} \in I$. The generating vector \mathbf{z} can be constructed using a component-by-component approach, see [11].

2.3 Existence of a special reconstructing rank-1 lattice

In this section, we proof a generalisation of [24, Lemma 2] and [27, Lemma 4.1]. Conceptually, $I = I_N \subset \mathbb{Z}^d$ represent frequency index sets used as support for the approximation of a function f by a trigonometric polynomial p based on sampling values $f(\mathbf{x}_j)$. The index sets $\mathcal{I} = \mathcal{I}_N \subset \mathbb{Z}^d$ may be considered as a superset of the difference set $\mathcal{D}(I) := \{\mathbf{k} - \mathbf{k}' : \mathbf{k}, \mathbf{k}' \in I\}$. As sampling nodes \mathbf{x}_j , we use the nodes $\mathbf{x}_j := \frac{j}{M}\mathbf{z} \bmod 1, j = 0, \dots, M-1$, of a reconstructing rank-1 lattice $\Lambda(\mathbf{z}, M, I)$ with generating vector $\mathbf{z} := (1, a, \dots, a^{d-1})^\top \in \mathbb{Z}^d$ of Korobov form, i.e., the condition $\mathbf{m}\mathbf{z} \not\equiv 0 \pmod{M}$ has to be fulfilled for all $\mathbf{m} \in \mathcal{D}(I) \setminus \{\mathbf{0}\}$ or one of the other (equivalent) conditions (2.1),(2.2),(2.3).

Lemma 2.1. *Let a sequence of frequency index sets $\mathcal{I}_N \subset \mathbb{Z}^d, d \in \mathbb{N}$, of finite cardinality $|\mathcal{I}_N|$ be given, which may depend on the refinement $N \in \mathbb{R}, N \geq 1$. For fixed refinement $N \in \mathbb{R}, N \geq 1$, and arbitrarily chosen parameter $\kappa \in \mathbb{R}, \kappa > 0$, let $M \in \mathbb{N}$ be a prime such that*

$$M > \frac{d|\mathcal{I}_N|}{1 - 2^{-\kappa}} + 1 \quad (2.5)$$

and

$$\mathcal{I}_N \cap M\mathbb{Z}^d = \{\mathbf{0}\}. \quad (2.6)$$

For an arbitrarily chosen monotonic increasing function $\varphi : \mathbb{N} \cup \{0\} \rightarrow [1, \infty)$ with $\varphi(0) = 1$, we define the shells $F_l(N) := \mathcal{I}_{N\varphi(l)} \setminus \mathcal{I}_{N\varphi(l-1)}, N \in \mathbb{R}, N \geq 1, l \in \mathbb{N}$, and for each $a \in \{1, \dots, M-1\}$ the sets

$$M_a^l := \{\mathbf{m} \in F_l(N) : m_1 + m_2 a + \dots + m_d a^{d-1} \equiv 0 \pmod{M} \text{ and } \mathbf{m} \neq M\mathbf{m}' \forall \mathbf{m}' \in \mathbb{Z}^d\}.$$

Then, there exists a number $a \in \{1, \dots, M-1\}$, such that

$$m_1 + m_2 a + \dots + m_d a^{d-1} \not\equiv 0 \pmod{M} \text{ for all } \mathbf{m} \in \mathcal{I}_N \setminus \{\mathbf{0}\} \quad (2.7)$$

and

$$|M_a^l| \leq A_l^N := |F_l(N)| d 2^{(l+1)\kappa} (2^\kappa - 1)^{-1} (M-1)^{-1}, \quad l \in \mathbb{N}. \quad (2.8)$$

Proof. This proof is a generalisation of the proofs of [24, Lemma 2] and [27, Lemma 4.1]. We remark that $F_l(N) = \emptyset$ may occur for some or all $l \in \mathbb{N}$ and then also $M_a^l = \emptyset$ follows. The idea is to prove that the number of integers $a \in \{1, \dots, M-1\}$ for which the statement of the lemma is not valid is less than $M-1$ and consequently, at least one $a \in \{1, \dots, M-1\}$ fulfills the statement. We consider the congruence

$$m_1 + m_2 a + \dots + m_d a^{d-1} \equiv 0 \pmod{M}. \quad (2.9)$$

For a fixed frequency $\mathbf{m} \in \mathbb{Z}^d$, we denote the set of natural numbers $a \in \{1, \dots, M-1\}$ which are solutions of congruence (2.9) by $A_M(\mathbf{m})$, i.e.,

$$A_M(\mathbf{m}) := \{a \in \{1, \dots, M-1\} : m_1 + m_2 a + \dots + m_d a^{d-1} \equiv 0 \pmod{M}\}.$$

Let a frequency $\mathbf{m} \in \mathcal{I}_N \setminus \{\mathbf{0}\}$ be given. Due to condition (2.6), at least one component fulfills $m_{s'} \not\equiv 0 \pmod{M}$ and we can apply Lagrange's Theorem. This yields that the congruence (2.9) has at most $d-1$ roots modulo M . Therefore, we have

$$|A_M(\mathbf{m})| \leq d-1 < d \quad (2.10)$$

for all $\mathbf{m} \in \mathcal{I}_N \setminus \{\mathbf{0}\}$. Next, we estimate the number of integers $a \in \{1, \dots, M-1\}$ for which the relation (2.7) is not valid for at least one $\mathbf{m} \in \mathcal{I}_N \setminus \{\mathbf{0}\}$. Therefore, we denote by G_0 the set of numbers $a \in \{1, \dots, M-1\}$ which are solutions of congruence (2.9) for at least one frequency $\mathbf{m} \in \mathcal{I}_N \setminus \{\mathbf{0}\}$,

$$G_0 = \bigcup_{\mathbf{m} \in \mathcal{I}_N \setminus \{\mathbf{0}\}} A_M(\mathbf{m}).$$

Since $|A_M(\mathbf{m})| < d$ by (2.10) and due to (2.5), we obtain

$$|G_0| \leq \sum_{\mathbf{m} \in \mathcal{I}_N \setminus \{\mathbf{0}\}} |A_M(\mathbf{m})| < d |\mathcal{I}_N| < (M-1)(1-2^{-\kappa}). \quad (2.11)$$

This means, for any $a \in \{1, \dots, M-1\} \setminus G_0$, the relations (2.7) are valid and $|\{1, \dots, M-1\} \setminus G_0| > M-1 - (M-1)(1-2^{-\kappa}) = (M-1)2^{-\kappa} > 0$.

Next, we consider the inequalities (2.8). For each $l \in \mathbb{N}$, we estimate the number of integers $a \in \{1, \dots, M-1\}$ for which $|M_a^l| > A_l^N$, i.e., for which the inequalities (2.8) are not fulfilled. Therefor, we define the sets $G_l := \{a \in \{1, \dots, M-1\} : |M_a^l| > A_l^N\}$, $l \in \mathbb{N}$. If $F_l(N) = \emptyset$, then obviously $|G_l| = 0$. Otherwise for $F_l(N) \neq \emptyset$, we have

$$\sum_{a \in G_l} |M_a^l| > \sum_{a \in G_l} A_l^N = |G_l| A_l^N. \quad (2.12)$$

We note that estimate (2.10) is also true for all $\mathbf{m} \in M_a^l$ due to Lagrange's Theorem, i.e., there exist at most $d-1$ many numbers $a \in \{1, \dots, M-1\}$ satisfying (2.9) for fixed $\mathbf{m} \in M_a^l$. Consequently, for fixed $\mathbf{m} \in M_a^l$, there exist at most $d-1$ sets M_a^l which contain \mathbf{m} . Thus, each $\mathbf{m} \in F_l(N)$ can belong to at most $d-1$ different sets M_a^l and therefore

$$\sum_{a \in G_l} |M_a^l| \leq (d-1) |F_l(N)| < d |F_l(N)|. \quad (2.13)$$

Comparing (2.12) and (2.13), we obtain $|G_l| A_l^N < d |F_l(N)|$ and by inserting the definition of A_l^N from (2.8), we infer

$$|G_l| < d |F_l(N)| / A_l^N = 2^{-(l+1)\kappa} (2^\kappa - 1)(M-1) = 2^{-l\kappa} (M-1)(1-2^{-\kappa}), \quad l \in \mathbb{N}, \quad (2.14)$$

if $F_l(N) \neq \emptyset$. Alltogether, relation (2.11) as well as relation (2.14) if $F_l(N) \neq \emptyset$ and $|G_l| = 0$ if $F_l(N) = \emptyset$ yield

$$\begin{aligned} \sum_{l=0}^{\infty} |G_l| &< \sum_{l=0}^{\infty} 2^{-l\kappa} (M-1)(1-2^{-\kappa}) = (M-1)(1-2^{-\kappa}) \sum_{l=0}^{\infty} (2^{-\kappa})^l \\ &= (M-1)(1-2^{-\kappa}) \frac{1}{1-2^{-\kappa}} = M-1. \end{aligned}$$

This means that the number of integers $a \in \{1, \dots, M-1\}$ for which the statement of the lemma is not valid is less than $M-1$. Since the cardinality $|\{1, \dots, M-1\}| = M-1$, there exists at least one $a \in \{1, \dots, M-1\}$ for which relations (2.7) and (2.8) are valid. \blacksquare

3 Aliasing error for rank-1 lattice sampling and general frequency index sets

Based on Lemma 2.1, we proof general statements for the aliasing error for arbitrary frequency index sets $I \subset \mathbb{Z}^d$ of finite cardinality. We are going to use the results of this section extensively in Section 4. The following lemma was proven in [24], see [24, Property 2°].

Lemma 3.1. *Let the dimensionality $d \in \mathbb{N}$, $d \geq 2$, a frequency index set $I \subset \mathbb{Z}^d$ of finite cardinality and a reconstructing rank-1 lattice $\Lambda(\mathbf{z}, M, I)$ with the nodes $\mathbf{x}_j := \frac{j}{M}\mathbf{z} \bmod 1$, $j = 0, \dots, M-1$, be given. We denote the Dirichlet kernel with frequencies supported on the index set I by $D_I(\mathbf{x}) := \sum_{\mathbf{k} \in I} e^{2\pi i \mathbf{k} \mathbf{x}}$. For an arbitrary vector $\mathbf{b} := (b_0, \dots, b_{M-1})^\top \in \mathbb{C}^M$, we have*

$$\left\| \frac{1}{M} \sum_{j=0}^{M-1} b_j D_I(\circ - \mathbf{x}_j) \Big|_{L_2(T^d)} \right\| \leq \left(\frac{1}{M} \sum_{j=0}^{M-1} |b_j|^2 \right)^{1/2} = \|\mathbf{b}/\sqrt{M}\|_2. \quad (3.1)$$

Additionally, for an arbitrary trigonometric polynomial $p: \mathbb{T}^d \rightarrow \mathbb{C}$ with frequencies supported on the index set I , $p(\mathbf{x}) := \sum_{\mathbf{k} \in I} \hat{p}_{\mathbf{k}} e^{2\pi i \mathbf{k} \mathbf{x}}$, $\hat{p}_{\mathbf{k}} \in \mathbb{C}$, we have $\tilde{S}_I p = p$.

Proof. Due to $\int_{\mathbb{T}^d} e^{2\pi i \mathbf{k} \mathbf{x}} d\mathbf{x} = \begin{cases} 1 & \text{for } \mathbf{k} = \mathbf{0}, \\ 0 & \text{for } \mathbf{k} \in \mathbb{Z}^d \setminus \{\mathbf{0}\}, \end{cases}$ we obtain

$$\begin{aligned} & \left\| \frac{1}{M} \sum_{j=0}^{M-1} b_j D_I(\circ - \mathbf{x}_j) \Big|_{L_2(\mathbb{T}^d)} \right\|^2 = \frac{1}{M^2} \int_{\mathbb{T}^d} \left| \sum_{j=0}^{M-1} b_j D_I(\mathbf{x} - \mathbf{x}_j) \right|^2 d\mathbf{x} \\ & = \frac{1}{M^2} \sum_{0 \leq j, j' \leq M-1} b_j D_I(\mathbf{x}_{j'} - \mathbf{x}_j) \overline{b_{j'}}. \end{aligned}$$

We can rewrite this as a quadratic form of the vector \mathbf{b}/\sqrt{M} and the matrix $\mathcal{D} = (\mathcal{D}_{j',j})_{j',j=0}^{M-1}$

with elements $\mathcal{D}_{j',j} := \frac{1}{M} D_I(\mathbf{x}_{j'} - \mathbf{x}_j)$, $\left\| \frac{1}{M} \sum_{j=0}^{M-1} b_j D_I(\circ - \mathbf{x}_j) \Big|_{L_2(T^d)} \right\|^2 = \left(\frac{\mathbf{b}}{\sqrt{M}} \right)^\mathsf{H} \mathcal{D} \left(\frac{\mathbf{b}}{\sqrt{M}} \right)$.

Next, we consider the matrix $\mathcal{D}^2 = \mathcal{D} \cdot \mathcal{D} := ((\mathcal{D}^2)_{j',j})_{j',j=0}^{M-1}$ with the elements $(\mathcal{D}^2)_{j',j}$. We obtain by using the reconstructing property (2.1) of the reconstructing rank-1 lattice

$\Lambda(\mathbf{z}, M, I)$ that $(\mathcal{D}^2)_{j',j} = \frac{1}{M^2} \sum_{\rho=0}^{M-1} D_I(\mathbf{x}_{j'} - \mathbf{x}_\rho) D_I(\mathbf{x}_\rho - \mathbf{x}_j) \stackrel{(2.1)}{=} \frac{1}{M} \sum_{\mathbf{k} \in I} e^{2\pi i \mathbf{k}(\mathbf{x}_{j'} - \mathbf{x}_j)} = \mathcal{D}_{j',j}$,

i.e., $\mathcal{D}^2 = \mathcal{D}$. Furthermore, we have $\mathcal{D}^\mathsf{H} = \mathcal{D}$ and therefore, $\mathcal{D} = \mathcal{D}^2 = \mathcal{D}^\mathsf{H} \mathcal{D}$ follows. Consequently, we infer

$$\left\| \frac{1}{M} \sum_{j=0}^{M-1} b_j D_I(\circ - \mathbf{x}_j) \Big|_{L_2(T^d)} \right\|^2 \leq \|\mathcal{D}\|_2^2 \left\| \left(\frac{\mathbf{b}}{\sqrt{M}} \right) \right\|_2^2 = \sigma_{\max}(\mathcal{D})^2 \left\| \frac{\mathbf{b}}{\sqrt{M}} \right\|_2^2,$$

where $\sigma_{\max}(\mathbf{D})$ denotes the largest singular value of the matrix \mathbf{D} . Last, we show $\sigma_{\max}(\mathbf{D}) \leq 1$. Let $\mathbf{D} = \mathbf{U}\mathbf{\Sigma}\mathbf{V}^H$ be a singular value decomposition of the matrix \mathbf{D} , where \mathbf{U}, \mathbf{V} are unitary matrices and $\mathbf{\Sigma} = \text{diag}((\sigma_1, \dots, \sigma_M))$ is a diagonal matrix of the singular values $\sigma_j \geq 0, j = 1, \dots, M$, of the matrix \mathbf{D} . Then, we infer from $\mathbf{U}\mathbf{\Sigma}\mathbf{V}^H = \mathbf{D} = \mathbf{D}^2 = \mathbf{D}^H\mathbf{D} = \mathbf{U}\mathbf{\Sigma}^2\mathbf{V}^H$ that $\sigma_j^2 = \sigma_j, j = 1, \dots, M$. Therefore, each singular value $\sigma_j \in \{0, 1\}$ and we obtain $\sigma_{\max}(\mathbf{D}) \leq 1$.

For $p(\mathbf{x}) = \sum_{\mathbf{k} \in I} \hat{p}_{\mathbf{k}} e^{2\pi i \mathbf{k} \mathbf{x}}$, we infer

$$\begin{aligned} \tilde{S}_I p(\mathbf{x}) &= \sum_{\mathbf{k} \in I} \frac{1}{M} \sum_{j=0}^{M-1} p(\mathbf{x}_j) e^{-2\pi i \mathbf{k} \mathbf{x}_j} e^{2\pi i \mathbf{k} \mathbf{x}} \\ &= \sum_{\mathbf{k} \in I} \left(\sum_{\mathbf{k}' \in I} \hat{p}_{\mathbf{k}'} \frac{1}{M} \sum_{j=0}^{M-1} e^{2\pi i j(\mathbf{k}' - \mathbf{k}) \mathbf{z} / M} \right) e^{2\pi i \mathbf{k} \mathbf{x}} \stackrel{(2.1)}{=} \sum_{\mathbf{k} \in I} \hat{p}_{\mathbf{k}} e^{2\pi i \mathbf{k} \mathbf{x}} = p(\mathbf{x}) \end{aligned}$$

since we use the sampling nodes $\mathbf{x}_j := \frac{j}{M} \mathbf{z} \bmod 1, j = 0, \dots, M-1$, from a reconstructing rank-1 lattice $\Lambda(\mathbf{z}, M, I)$. \blacksquare

Lemma 3.2. *Let the dimensionality $d \in \mathbb{N}, d \geq 2$, a function $f \in C(\mathbb{T}^d) \cap L^2(\mathbb{T}^d)$ with point-wise convergent Fourier series, a frequency index set $I \subset \mathbb{Z}^d$ of finite cardinality and a reconstructing rank-1 lattice $\Lambda(\mathbf{z}, M, I)$ with the nodes $\mathbf{x}_j := \frac{j}{M} \mathbf{z} \bmod 1, j = 0, \dots, M-1$, be given. Additionally, we define shells $U_l \subset \mathbb{Z}^d, l \in \mathbb{N} \cup \{0\}$, with the properties $U_{l'} \cap U_{l''} = \emptyset$ for $l' \neq l''$ and $\text{supp} \hat{f} \setminus I \subset \bigcup_{l=0}^{\infty} U_l$, where $\text{supp} \hat{f} := \{\mathbf{k} \in \mathbb{Z}^d : \hat{f}_{\mathbf{k}} \neq 0\}$. Then, we have*

$$\|\tilde{S}_I(f - S_I f) |L^2(\mathbb{T}^d)\| \leq \sum_{l=0}^{\infty} \sigma_l, \quad \sigma_l := \left(\frac{1}{M} \sum_{j=0}^{M-1} |S_{U_l} f(\mathbf{x}_j)|^2 \right)^{1/2}.$$

Proof. By definition, we have

$$\begin{aligned} \tilde{S}_I(f - S_I f) &= \sum_{\mathbf{h} \in I} \frac{1}{M} \sum_{j=0}^{M-1} (f - S_I f)(\mathbf{x}_j) e^{-2\pi i \mathbf{h} \mathbf{x}_j} e^{2\pi i \mathbf{h} \mathbf{o}} \\ &= \frac{1}{M} \sum_{j=0}^{M-1} S_{\text{supp} \hat{f} \setminus I} f(\mathbf{x}_j) D_I(\mathbf{o} - \mathbf{x}_j) \\ &= \frac{1}{M} \sum_{j=0}^{M-1} \left(\sum_{\mathbf{k} \in \text{supp} \hat{f} \setminus I} \hat{f}_{\mathbf{k}} e^{2\pi i \mathbf{k} \mathbf{x}_j} \right) D_I(\mathbf{o} - \mathbf{x}_j) \\ &= \frac{1}{M} \sum_{j=0}^{M-1} \left(\sum_{l=0}^{\infty} \sum_{\mathbf{k} \in U_l} \hat{f}_{\mathbf{k}} e^{2\pi i \mathbf{k} \mathbf{x}_j} \right) D_I(\mathbf{o} - \mathbf{x}_j) \\ &= \sum_{l=0}^{\infty} \frac{1}{M} \sum_{j=0}^{M-1} \sum_{\mathbf{k} \in U_l} \hat{f}_{\mathbf{k}} e^{2\pi i \mathbf{k} \mathbf{x}_j} D_I(\mathbf{o} - \mathbf{x}_j) \\ &= \sum_{l=0}^{\infty} \frac{1}{M} \sum_{j=0}^{M-1} S_{U_l} f(\mathbf{x}_j) D_I(\mathbf{o} - \mathbf{x}_j). \end{aligned}$$

We apply the Minkowski inequality and Lemma 3.1 with $b_j := S_{U_l} f(\mathbf{x}_j)$. This yields

$$\begin{aligned}
\|\tilde{S}_I(f - S_I f)|_{L^2(\mathbb{T}^d)}\| &= \left\| \sum_{l=0}^{\infty} \frac{1}{M} \sum_{j=0}^{M-1} S_{U_l} f(\mathbf{x}_j) D_I(\circ - \mathbf{x}_j) \right\|_{L^2(\mathbb{T}^d)} \\
&\leq \sum_{l=0}^{\infty} \left\| \frac{1}{M} \sum_{j=0}^{M-1} S_{U_l} f(\mathbf{x}_j) D_I(\circ - \mathbf{x}_j) \right\|_{L^2(\mathbb{T}^d)} \\
&\stackrel{(3.1)}{\leq} \sum_{l=0}^{\infty} \left(\frac{1}{M} \sum_{j=0}^{M-1} |S_{U_l} f(\mathbf{x}_j)|^2 \right)^{1/2} = \sum_{l=0}^{\infty} \sigma_l
\end{aligned}$$

and the assertion follows. \blacksquare

Lemma 3.3. *Let the dimensionality $d \in \mathbb{N}$, a weight function $\omega : \mathbb{Z}^d \rightarrow (0, \infty]$, a smoothness parameter $\nu > 0$, the sequence of frequency index sets $I_N := \{\mathbf{k} \in \mathbb{Z}^d : \omega(\mathbf{k})^{1/\nu} \leq N\}$ of refinement $N \in \mathbb{R}$, $N \geq 1$, and a function $f \in L^2(\mathbb{T}^d) \cap \mathcal{H}^\omega(\mathbb{T}^d)$ be given. Then, the truncation error is bounded by*

$$\|f - S_{I_N} f\|_{L^2(\mathbb{T}^d)}^2 \leq N^{-\nu} \|f\|_{\mathcal{H}^\omega(\mathbb{T}^d)}.$$

Proof. We have

$$\mathbb{Z}^d \setminus I_N = \{\mathbf{k} \in \mathbb{Z}^d : \omega(\mathbf{k})^{1/\nu} > N\} = \{\mathbf{k} \in \mathbb{Z}^d : \frac{1}{\omega(\mathbf{k})^{1/\nu}} < \frac{1}{N}\} = \{\mathbf{k} \in \mathbb{Z}^d : \frac{1}{\omega(\mathbf{k})^2} < N^{-2\nu}\}$$

and this yields the assertion since

$$\begin{aligned}
\|f - S_{I_N} f\|_{L^2(\mathbb{T}^d)}^2 &= \sum_{\mathbf{k} \in \mathbb{Z}^d \setminus I_N} \frac{\omega(\mathbf{k})^2}{\omega(\mathbf{k})^2} |\hat{f}_{\mathbf{k}}|^2 \leq \max_{\mathbf{k} \in \mathbb{Z}^d \setminus I_N} \frac{1}{\omega(\mathbf{k})^2} \sum_{\mathbf{k} \in \mathbb{Z}^d \setminus I_N} \omega(\mathbf{k})^2 |\hat{f}_{\mathbf{k}}|^2 \\
&\leq N^{-2\nu} \sum_{\mathbf{k} \in \mathbb{Z}^d \setminus I_N} \omega(\mathbf{k})^2 |\hat{f}_{\mathbf{k}}|^2 \leq N^{-2\nu} \|f\|_{\mathcal{H}^\omega(\mathbb{T}^d)}^2.
\end{aligned}$$

\blacksquare

Theorem 3.4. *Let the dimensionality $d \in \mathbb{N}$, $d \geq 2$, a function $f \in L^2(\mathbb{T}^d) \cap \mathcal{H}^\omega(\mathbb{T}^d)$ with point-wise convergent Fourier series, a smoothness parameter $\nu > 0$ and the sequence of frequency index sets $I_N := \{\mathbf{k} \in \mathbb{Z}^d : \omega(\mathbf{k})^{1/\nu} \leq N\}$ with refinement $N \in \mathbb{R}$, $N \geq 1$, be given, where $\omega : \mathbb{Z}^d \rightarrow (0, \infty]$ is a weight function. Furthermore, let $\mathcal{I}_N \subset \mathbb{Z}^d$ be a nested sequence of frequency index sets with refinement $N \in \mathbb{R}$, $N \geq 1$,*

$$\mathcal{I}_{N'} \subset \mathcal{I}_{N''} \text{ for } N' \leq N'', \tag{3.2}$$

such that the inclusion $\mathcal{I}_N \supset \mathcal{D}(I_N) := \{\mathbf{k} - \mathbf{k}' : \mathbf{k}, \mathbf{k}' \in I_N\}$ is valid for all $N \in \mathbb{R}$, $N \geq 1$. For each fixed $N \in \mathbb{R}$, $N \geq 1$, let a parameter $\kappa > 0$ and a prime number $M \in \mathbb{N}$,

$$M > \frac{d |\mathcal{I}_N|}{1 - 2^{-\kappa}} + 1, \tag{3.3}$$

be given. Additionally, let the inequality

$$|\{\mathbf{m} \in \mathcal{I}_{N2^l} : \exists \mathbf{m}' \in \mathbb{Z}^d \text{ such that } \mathbf{m} = M\mathbf{m}'\}| \leq C \frac{|\mathcal{I}_{N2^l}|}{M} \psi(l) + 1 \quad \forall l \in \mathbb{N} \quad (3.4)$$

be valid, where $\psi : [0, \infty) \rightarrow [1, \infty)$ and $C > 0$ is a constant which does not depend on N or M . Then, there exists a reconstructing rank-1 lattice $\Lambda(\mathbf{z}, M, I_N)$ with generating vector $\mathbf{z} := (1, a, \dots, a^{d-1})^\top \in \mathbb{Z}^d$ of Korobov form such that the aliasing error is bounded by

$$\begin{aligned} \|S_{I_N} f - \tilde{S}_{I_N} f\|_{L^2(\mathbb{T}^d)} &\leq 2^\nu N^{-\nu} \|f\|_{\mathcal{H}^\omega(\mathbb{T}^d)} \\ &\quad \cdot \sum_{l=0}^{\infty} \sqrt{2(2 + (1 - 2^{-\kappa})C\psi(l+1))} 2^{(l+1)(\frac{\kappa}{2} - \nu)} \sqrt{\frac{|\mathcal{I}_{N2^{l+1}}|}{|\mathcal{I}_N|}}. \end{aligned}$$

Proof. This proof is a generalisation of [24, Theorem 2]). Since inequality (3.3) is valid, we apply Lemma 2.1 and obtain that there exists a number $a \in \{1, \dots, M-1\}$ which fulfills properties (2.7) and (2.8). Since property (2.7) is valid, the rank-1 lattice $\Lambda(\mathbf{z}, M)$ with the generating vector $\mathbf{z} := (1, a, \dots, a^{d-1})^\top$ and the nodes $\mathbf{x}_j := \frac{j}{M}\mathbf{z} \bmod 1$, $j = 0, \dots, M-1$, is a reconstructing rank-1 lattice $\Lambda(\mathbf{z}, M, I_N)$. We use this special rank-1 lattice $\Lambda(\mathbf{z}, M, I_N)$ for computing the approximated Fourier coefficients $\hat{f}_{\mathbf{k}}$, $\mathbf{k} \in I_N$, from the sampling values $f(\mathbf{x}_j)$. Since the Fourier partial sum $S_{I_N} f$ of the function f is a trigonometric polynomial with frequencies supported on the index set I_N and by applying Lemma 3.1, we obtain $\tilde{S}_{I_N}(S_{I_N} f) = S_{I_N} f$. This yields $S_{I_N} f - \tilde{S}_{I_N} f = \tilde{S}_{I_N}(f - S_{I_N} f)$. Next, we set the shells $U_l := I_{N2^{l+1}} \setminus I_{N2^l}$, $l = 0, 1, \dots$, and consequently, the property $U_l \cap U_{l'} = \emptyset \quad \forall l \neq l'$ is valid. We apply Lemma 3.2 and we obtain $\|\tilde{S}_{I_N}(f - S_{I_N} f)\|_{L^2(\mathbb{T}^d)} \leq \sum_{l=0}^{\infty} \sigma_l$, where

$$\sigma_l := \left(\frac{1}{M} \sum_{j=0}^{M-1} |S_{U_l} f(\mathbf{x}_j)|^2 \right)^{1/2}, \quad l \in \mathbb{N} \cup \{0\}.$$

Next, we want to estimate

$$\sigma_l^2 \leq B_l \sum_{\mathbf{k} \in U_l} |\hat{f}_{\mathbf{k}}|^2,$$

with numbers $B_l \geq 0$, which have to be determined. We have

$$\sigma_l^2 = \frac{1}{M} \sum_{j=0}^{M-1} \left| \sum_{\mathbf{k} \in U_l} \hat{f}_{\mathbf{k}} e^{2\pi i \mathbf{k} \mathbf{x}_j} \right|^2 = \frac{1}{M} \sum_{j=0}^{M-1} \sum_{\mathbf{k}, \mathbf{h} \in U_l} \hat{f}_{\mathbf{k}} \overline{\hat{f}_{\mathbf{h}}} e^{2\pi i (\mathbf{k} - \mathbf{h}) \mathbf{x}_j} = \sum_{\mathbf{k}, \mathbf{h} \in U_l} \hat{f}_{\mathbf{k}} \overline{\hat{f}_{\mathbf{h}}} \Delta_M(\mathbf{k} - \mathbf{h}),$$

where

$$\Delta_M(\mathbf{m}) := \frac{1}{M} \sum_{j=0}^{M-1} e^{2\pi i j \mathbf{m} \mathbf{z} / M} = \begin{cases} 1 & \text{for } m_1 + m_2 a + \dots + m_d a^{d-1} \equiv 0 \pmod{M}, \\ 0 & \text{for } m_1 + m_2 a + \dots + m_d a^{d-1} \not\equiv 0 \pmod{M}. \end{cases}$$

For fixed frequency $\mathbf{k} \in U_l$, we define the set of frequencies

$$\theta_{\mathbf{k}} := \{\mathbf{h} \in U_l : \Delta_M(\mathbf{k} - \mathbf{h}) = 1\},$$

and by applying the Cauchy Schwarz inequality twice, we obtain

$$\begin{aligned}
\sigma_l^2 &= \sum_{\mathbf{k} \in U_l} \hat{f}_{\mathbf{k}} \sum_{\mathbf{h} \in \theta_{\mathbf{k}}} \overline{\hat{f}_{\mathbf{h}}} \leq \left(\sum_{\mathbf{k} \in U_l} |\hat{f}_{\mathbf{k}}|^2 \right)^{1/2} \left(\sum_{\mathbf{k} \in U_l} \left| \sum_{\mathbf{h} \in \theta_{\mathbf{k}}} \overline{\hat{f}_{\mathbf{h}}} \right|^2 \right)^{1/2} \\
&\leq \left(\sum_{\mathbf{k} \in U_l} |\hat{f}_{\mathbf{k}}|^2 \right)^{1/2} \left(\sum_{\mathbf{k} \in U_l} \left(\sum_{\mathbf{h} \in \theta_{\mathbf{k}}} 1 \cdot |\hat{f}_{\mathbf{h}}| \right)^2 \right)^{1/2} \\
&\leq \left(\sum_{\mathbf{k} \in U_l} |\hat{f}_{\mathbf{k}}|^2 \right)^{1/2} \left(\sum_{\mathbf{k} \in U_l} |\theta_{\mathbf{k}}| \sum_{\mathbf{h} \in \theta_{\mathbf{k}}} |\hat{f}_{\mathbf{h}}|^2 \right)^{1/2}.
\end{aligned}$$

We have $\mathbf{k} - \mathbf{h} \in \mathcal{D}(I_{N2^{l+1}}) \subset \mathcal{I}_{N2^{l+1}}$ for $\mathbf{k}, \mathbf{h} \in U_l$ and this yields

$$|\theta_{\mathbf{k}}| \leq |\{\mathbf{m} \in \mathcal{I}_{N2^{l+1}} : m_1 + m_2 a + \dots + m_d a^{d-1} \equiv 0 \pmod{M}\}|.$$

We define the function $\varphi(l) := 2^l$ for $l \in \mathbb{N} \cup \{0\}$. Due to property (2.8) in Lemma 2.1, we obtain

$$\begin{aligned}
&\left| \{\mathbf{m} \in I_{N\varphi(l+1)}^{d,0} : m_1 + m_2 a + \dots + m_d a^{d-1} \equiv 0 \pmod{M} \text{ and } \mathbf{m} \neq M\mathbf{m}' \forall \mathbf{m}' \in \mathbb{Z}^d\} \right| \\
&= \left| \bigcup_{j=1}^{l+1} F_j(N) \right| \leq \sum_{j=1}^{l+1} A_j^N.
\end{aligned}$$

Then, we have

$$|\theta_{\mathbf{k}}| \leq B_l := \sum_{j=1}^{l+1} A_j^N + C \frac{|\mathcal{I}_{N2^{l+1}}|}{M} \psi(l+1) + 1 \quad (3.5)$$

and

$$\sigma_l^2 \leq \left(\sum_{\mathbf{k} \in U_l} |\hat{f}_{\mathbf{k}}|^2 \right)^{1/2} \left(B_l \sum_{\mathbf{k} \in U_l} \sum_{\mathbf{h} \in \theta_{\mathbf{k}}} |\hat{f}_{\mathbf{h}}|^2 \right)^{1/2}.$$

For an arbitrarily chosen $\mathbf{k} \in U_l$, let $\mathbf{h} \in \theta_{\mathbf{k}}$. This means, we have $(\mathbf{h} - \mathbf{k})\mathbf{z} \equiv 0 \pmod{M}$. If $\mathbf{h} \in \theta_{\mathbf{k}'}$ for another $\mathbf{k}' \in U_l$, $\mathbf{k}' \neq \mathbf{k}$, then $(\mathbf{h} - \mathbf{k}')\mathbf{z} \equiv 0 \pmod{M}$ is valid and $(\mathbf{k} - \mathbf{k}')\mathbf{z} \equiv 0 \pmod{M}$ follows. This yields $\mathbf{k}' \in \theta_{\mathbf{k}}$. Especially, we have $\mathbf{k} \in \theta_{\mathbf{k}}$. Therefore, each frequency $\mathbf{h}' \in U_l$ is element of at most B_l many distinct sets $\theta_{\mathbf{k}}$. This means, we obtain

$$\sum_{\mathbf{k} \in U_l} \sum_{\mathbf{h} \in \theta_{\mathbf{k}}} |\hat{f}_{\mathbf{h}}|^2 \leq \sum_{\mathbf{k} \in U_l} B_l |\hat{f}_{\mathbf{k}}|^2$$

and

$$\begin{aligned}
\sigma_l^2 &\leq \left(\sum_{\mathbf{k} \in U_l} |\hat{f}_{\mathbf{k}}|^2 \right)^{1/2} \left(B_l^2 \sum_{\mathbf{k} \in U_l} |\hat{f}_{\mathbf{k}}|^2 \right)^{1/2} = B_l \sum_{\mathbf{k} \in U_l} |\hat{f}_{\mathbf{k}}|^2 \leq B_l \sum_{\mathbf{k} \in \mathbb{Z}^d \setminus I_{N2^l}} |\hat{f}_{\mathbf{k}}|^2 \\
&= B_l \|f - S_{I_{N2^l}} f\|_{L^2(\mathbb{T}^d)}^2 \leq B_l (N2^l)^{-2\nu} \|f\|_{\mathcal{H}^\omega(\mathbb{T}^d)}^2.
\end{aligned}$$

Next, we estimate B_l . Using the inequality $\frac{1}{M-1} \leq \frac{2}{M}$ for $M \geq 2$ as well as (3.5) and (2.8), we infer

$$\begin{aligned}
B_l &= \sum_{j=1}^{l+1} |\mathcal{I}_{N\varphi(j)} \setminus \mathcal{I}_{N\varphi(j-1)}| d 2^{(j+1)\kappa} (2^\kappa - 1)^{-1} (M-1)^{-1} + C \frac{|\mathcal{I}_{N2^{l+1}}|}{M} \psi(l+1) + 1 \\
&\leq d \frac{2^\kappa}{2^\kappa - 1} 2^{(l+1)\kappa} \frac{2}{M} \sum_{j=1}^{l+1} |\mathcal{I}_{N2^j} \setminus \mathcal{I}_{N2^{j-1}}| + C \psi(l+1) |\mathcal{I}_{N2^{l+1}}| / M + 1 \\
&\stackrel{(3.2)}{\leq} d 2^{(l+1)\kappa} \frac{|\mathcal{I}_{N2^{l+1}}|}{M} \left(\frac{2}{1-2^{-\kappa}} + C \psi(l+1) \right) + 1 \\
&\stackrel{(3.3)}{\leq} d 2^{(l+1)\kappa} \frac{|\mathcal{I}_{N2^{l+1}}|}{\frac{d|\mathcal{I}_N|}{1-2^{-\kappa}} + 1} \left(\frac{2}{1-2^{-\kappa}} + C \psi(l+1) \right) + 1 \\
&\leq 2^{(l+1)\kappa+1} \frac{|\mathcal{I}_{N2^{l+1}}|}{|\mathcal{I}_N|} (2 + (1-2^{-\kappa}) C \psi(l+1))
\end{aligned}$$

and this yields

$$\begin{aligned}
\sigma_l &\leq (N2^l)^{-\nu} \|f|_{\mathcal{H}^\omega(\mathbb{T}^d)}\| \sqrt{B_l} \\
&\leq \sqrt{2(2 + (1-2^{-\kappa}) C \psi(l+1))} 2^\nu N^{-\nu} \|f|_{\mathcal{H}^\omega(\mathbb{T}^d)}\| 2^{(l+1)(\frac{\kappa}{2}-\nu)} \sqrt{\frac{|\mathcal{I}_{N2^{l+1}}|}{|\mathcal{I}_N|}}.
\end{aligned}$$

■

4 Approximation error for rank-1 lattice sampling and frequency index sets $I_N^{d,T}$

Next, we apply the general results from Section 2 and Section 3. Therefor, we use the index sets $I = I_N = I_N^{d,T}$. In the case $-\infty \leq T \leq 0$, we set $\mathcal{I}_N := I_N^{d,T} \frac{2^{\frac{d-T}{1-T}}}{N^{1+\frac{d}{d-T}}}$, where $\mathcal{I}_N \supset \mathcal{D}(I_N^{d,T})$, see Lemma 4.2. This means, we cover the difference set $\mathcal{D}(I_N^{d,T})$ with an index set $I_N^{d,T} \frac{2^{\frac{d-T}{1-T}}}{N^{1+\frac{d}{d-T}}}$ of larger refinement $2^{\frac{d-T}{1-T}} N^{1+\frac{d}{d-T}}$. In the case $0 < T < 1$, we set $\mathcal{I}_N := \mathcal{D}(I_N^{d,T})$. Before we estimate the truncation error $\|f - S_{I_N^{d,T}} f|_{L^2(\mathbb{T}^d)}\|$ and the aliasing error $\|S_{I_N^{d,T}} f - \tilde{S}_{I_N^{d,T}} f|_{L^2(\mathbb{T}^d)}\|$, we show preliminary lemmata for the cardinalities and embeddings of the frequency index sets $I_N^{d,T}$.

Lemma 4.1. *Let the dimensionality $d \in \mathbb{N}$, and a parameter T , $-\infty \leq T < 1$, be given. Then, the cardinalities of the frequency index sets $I_N^{d,T}$ are*

$$|I_N^{d,T}| = \begin{cases} \Theta(N^d) & \text{for } T = -\infty, \\ \Theta(N^{\frac{T-1}{T/d-1}}) & \text{for } -\infty < T < 0, \\ \Theta(N \log^{d-1} N) & \text{for } T = 0, \\ \Theta(N) & \text{for } 0 < T < 1, \end{cases} \quad (4.1)$$

for fixed $d \in \mathbb{N}$.

Proof. The upper bounds follow from [14, Lemma 2.6] and [15, Section 3.3 Lemma 2]. Therefore, we still have to discuss the lower bounds.

- Case $T = -\infty$. Since the volume of an l_1 ball with radius N is $\frac{2^d}{d!} N^d$ (e.g. see [29]), the lower bound $|I_N^{d,-\infty}| = \Omega(N^d)$ follows.
- Case $-\infty < T < 0$. First, we show $I_N^{d,-\infty} \subset I_N^{d,T}$. For arbitrary $\mathbf{k} \in I_N^{d,-\infty}$, we have

$$N^{\frac{1-T}{d-T}} \geq \max(1, \|\mathbf{k}\|_1) = \max(1, \|\mathbf{k}\|_1)^{-\frac{T}{d-T}} \max(1, \|\mathbf{k}\|_1)^{1+\frac{T}{d-T}}.$$

Since $\max(1, \|\mathbf{k}\|_1)^d \geq \max(1, \|\mathbf{k}\|_\infty) \geq \prod_{s=1}^d \max(1, |k_s|)$, we infer

$$\begin{aligned} N^{\frac{1-T}{d-T}} &\geq \max(1, \|\mathbf{k}\|_1)^{-\frac{T}{d-T}} \prod_{s=1}^d \max(1, |k_s|)^{\frac{1}{d}(1+\frac{T}{d-T})} \\ &= \max(1, \|\mathbf{k}\|_1)^{-\frac{T}{1-T} \frac{1-T}{d-T}} \prod_{s=1}^d \max(1, |k_s|)^{\frac{1}{1-T} \frac{1-T}{d-T}} \end{aligned}$$

and consequently $\max(1, \|\mathbf{k}\|_1)^{-\frac{T}{1-T}} \prod_{s=1}^d \max(1, |k_s|)^{\frac{1}{1-T}} \leq N$. This means, we have $\mathbf{k} \in I_N^{d,T}$ and therefore we obtain $I_N^{d,-\infty} \subset I_N^{d,T}$. Since we have $|I_N^{d,-\infty}| \geq c_1(d) N^{\frac{1-T}{d-T} d}$, where $c_1(d)$ is a constant which only depends on the dimensionality d , we obtain $|I_N^{d,T}| = \Omega(N^{\frac{T-1}{T/d-1}})$ due to $|I_N^{d,T}| \geq |I_N^{d,-\infty}| \geq c_1(d) N^{\frac{1-T}{d-T} d} = c_1(d) N^{\frac{T-1}{T/d-1}}$.

- Case $T = 0$. For the lower bound $|I_N^{d,0}| = \Omega(N \log^{d-1} N)$, we refer to [8, Section 5.3].
- Case $0 < T < 1$. Since the frequencies on the coordinate axis from $-[N]$ to $[N]$ are elements of $I_N^{d,T}$, we obtain $|I_N^{d,T}| \geq 2d[N] + 1 \geq 2d(N-1) + 1$.

These lower bounds yield the assertion. ■

Lemma 4.2. *Let the dimensionality $d \in \mathbb{N}$, and a parameter T , $-\infty \leq T \leq 0$, be given. We consider the difference set $\mathcal{D}(I_N^{d,T}) := \{\mathbf{k}' - \mathbf{k} : \mathbf{k}, \mathbf{k}' \in I_N^{d,T}\}$. Then, we have the inclusion*

$$\mathcal{D}(I_N^{d,T}) \subset I_{2^{\frac{d-T}{1-T}} N^{1+\frac{d}{d-T}}}^{d,T}. \quad (4.2)$$

Proof. For $\mathbf{k} \in I_N^{d,T}$, we have $\max(1, \|\mathbf{k}\|_1)^{-\frac{T}{1-T}} \prod_{s=1}^d \max(1, |k_s|)^{\frac{1}{1-T}} \leq N$ by definition. Consequently, for $\mathbf{k}, \mathbf{k}' \in I_N^{d,T}$ and $-\infty \leq T < 0$, we infer

$$\begin{aligned} &\max(1, \|\mathbf{k} - \mathbf{k}'\|_1) \prod_{s=1}^d \max(1, |k_s - k'_s|)^{-\frac{1}{T}} \\ &\leq (\max(1, \|\mathbf{k}\|_1) + \max(1, \|\mathbf{k}'\|_1)) \prod_{s=1}^d (\max(1, |k_s|) + \max(1, |k'_s|))^{-\frac{1}{T}} \\ &\leq (\max(1, \|\mathbf{k}\|_1) + \max(1, \|\mathbf{k}'\|_1)) 2^{-\frac{d}{T}} \prod_{s=1}^d \max(1, |k_s|)^{-\frac{1}{T}} \max(1, |k'_s|)^{-\frac{1}{T}} \\ &\leq 2^{-\frac{d}{T}} N^{-\frac{1-T}{T}} \left(\prod_{s=1}^d \max(1, |k'_s|)^{-\frac{1}{T}} + \prod_{s=1}^d \max(1, |k_s|)^{-\frac{1}{T}} \right). \end{aligned}$$

Next, we estimate dominating mixed smoothness by isotropic smoothness. Since we have $\prod_{s=1}^d \max(1, |k_s|) \leq \max(1, \|\mathbf{k}\|_\infty)^d \leq \max(1, \|\mathbf{k}\|_1)^d$ for $\mathbf{k} \in \mathbb{Z}^d$, we obtain

$$\begin{aligned} \prod_{s=1}^d \max(1, |k_s|)^{-\frac{1}{T}} &= \prod_{s=1}^d \max(1, |k_s|)^{\frac{1}{d-T}} \prod_{s=1}^d \max(1, |k_s|)^{-\frac{1}{T} - \frac{1}{d-T}} \\ &\leq \max(1, \|\mathbf{k}\|_1)^{\frac{d}{d-T}} \prod_{s=1}^d \max(1, |k_s|)^{-\frac{d}{T(d-T)}} \\ &= \left(\max(1, \|\mathbf{k}\|_1)^{-\frac{T}{1-T}} \prod_{s=1}^d \max(1, |k_s|)^{\frac{1}{1-T}} \right)^{-\frac{1-T}{T} \frac{d}{d-T}} \\ &\leq N^{-\frac{1-T}{T} \frac{d}{d-T}} \end{aligned}$$

and analogously $\prod_{s=1}^d \max(1, |k'_s|)^{-\frac{1}{T}} \leq N^{-\frac{1-T}{T} \frac{d}{d-T}}$. For $T = 0$, we have

$$\prod_{s=1}^d \max(1, |k_s - k'_s|) \leq 2^d \prod_{s=1}^d \max(1, |k_s|) \prod_{s=1}^d \max(1, |k'_s|) \leq 2^d N^2.$$

These results yield

$$\max(1, \|\mathbf{k} - \mathbf{k}'\|_1)^{-\frac{T}{1-T}} \prod_{s=1}^d \max(1, |k_s - k'_s|)^{\frac{1}{1-T}} \leq 2^{\frac{d-T}{1-T}} N^{1+\frac{d}{d-T}} \text{ for all } \mathbf{k}, \mathbf{k}' \in I_N^{d,T}$$

and inclusion (4.2) follows. ■

4.1 Truncation error

We estimate the truncation error $\|f - S_{I_N^{d,T}} f\|_{L^2(\mathbb{T}^d)}$, since this error is part of the approximation error $\|f - \tilde{S}_{I_N^{d,T}} f\|_{L^2(\mathbb{T}^d)}$ and since we also need the result as a prerequisite for Theorem 3.4. First, we show $\mathcal{H}^{\alpha,\beta}(\mathbb{T}^d) \subset L^2(\mathbb{T}^d)$ for $\beta \geq 0$ and $\alpha > -\beta$.

Lemma 4.3. *Let the parameter $\alpha, \beta \in \mathbb{R}$, $\beta \geq 0$, $\alpha > -\beta$ be given. Then, $\mathcal{H}^{\alpha,\beta}(\mathbb{T}^d) \subset L^2(\mathbb{T}^d)$.*

Proof. In the case $\alpha \geq 0$, we obviously have $\omega^{\alpha,\beta}(\mathbf{k}) \geq 1$ for all $\mathbf{k} \in \mathbb{Z}^d$. In the case $\alpha < 0$, due to $\prod_{s=1}^d \max(1, |k_s|) \leq \max(1, \|\mathbf{k}\|_1)^d$ for $\mathbf{k} \in \mathbb{Z}^d$ and $\beta + \frac{\alpha}{d} > \alpha + \beta > 0$, we infer

$$\omega^{\alpha,\beta}(\mathbf{k}) := \max(1, \|\mathbf{k}\|_1)^\alpha \prod_{s=1}^d \max(1, |k_s|)^\beta \geq \prod_{s=1}^d \max(1, |k_s|)^{\beta + \frac{\alpha}{d}} \geq 1 \text{ for all } \mathbf{k} \in \mathbb{Z}^d.$$

Consequently, we obtain

$$\|f\|_{L^2(\mathbb{T}^d)} = \sqrt{\sum_{\mathbf{k} \in \mathbb{Z}^d} |\hat{f}_{\mathbf{k}}|^2} \leq \sqrt{\sum_{\mathbf{k} \in \mathbb{Z}^d} \omega^{\alpha,\beta}(\mathbf{k})^2 |\hat{f}_{\mathbf{k}}|^2} = \|f\|_{\mathcal{H}^{\alpha,\beta}(\mathbb{T}^d)} < \infty$$

for an arbitrarily chosen function $f \in \mathcal{H}^{\alpha,\beta}(\mathbb{T}^d)$. ■

Next, we estimate the truncation error $\|f - S_{I_N^{d,T}} f\|_{L^2(\mathbb{T}^d)}$. The following lemma is a special case of [14, Lemma 3.5].

Lemma 4.4. *Let the dimensionality $d \in \mathbb{N}$, a function $f \in \mathcal{H}^{\alpha,\beta}(\mathbb{T}^d)$ and the d -dimensional index set $I_N^{d,T}$ of refinement $N \in \mathbb{R}$, $N \geq 1$, be given, where $\beta \geq 0$, $\alpha > -\beta$ and $T := -\alpha/\beta$. Then, the truncation error is bounded by*

$$\|f - S_{I_N^{d,T}} f\|_{L^2(\mathbb{T}^d)} \leq N^{-(\alpha+\beta)} \|f\|_{\mathcal{H}^{\alpha,\beta}(\mathbb{T}^d)}.$$

Proof. We set $\omega(\mathbf{k}) := \omega^{\alpha,\beta}(\mathbf{k})$, $\nu := \alpha + \beta$ and $I_N := I_N^{d,T}$. Since $T := -\alpha/\beta$, the conditions $\beta \geq 0$ and $\alpha > -\beta$ ensure that $-\infty \leq T < 1$. From Lemma 4.3, we obtain $\mathcal{H}^{\alpha,\beta}(\mathbb{T}^d) \subset L^2(\mathbb{T}^d)$. Next, we apply Lemma 3.3. Due to $\omega(\mathbf{k})^{1/\nu} = \max(1, \|\mathbf{k}\|_1)^{\frac{\alpha}{\alpha+\beta}} \prod_{s=1}^d \max(1, |k_s|)^{\frac{\beta}{\alpha+\beta}} = \omega^{-\frac{T}{1-T}, \frac{1}{1-T}}(\mathbf{k})$, we obtain $\|f - S_{I_N} f\|_{L^2(\mathbb{T}^d)} \leq N^{-\nu} \|f\|_{\mathcal{H}^\omega(\mathbb{T}^d)} = N^{-(\alpha+\beta)} \|f\|_{\mathcal{H}^{\alpha,\beta}(\mathbb{T}^d)}$. ■

4.2 Aliasing error

We are going to apply Theorem 3.4 for the frequency index sets $I_N = I_N^{d,T}$ in order to estimate the aliasing error $\|S_{I_N} f - \tilde{S}_{I_N} f\|_{L^2(\mathbb{T}^d)}$. Therefor, we show that condition (3.4) is fulfilled for the frequency index sets $I_N^{d,T}$ of refinements $N \in \mathbb{R}$, $N \geq 2$, and parameters $-\infty \leq T < 1$.

4.2.1 Cases $-\infty \leq T \leq 0$

Lemma 4.5. *Let the dimensionality $d \in \mathbb{N}$, $d \geq 2$, a parameter T , $-\infty \leq T \leq 0$, and $M \in \mathbb{N}$, $M \geq 2$, be given. Then, we have*

$$|\{\mathbf{m} \in I_{N 2^{(l+1)(1+\frac{d}{d-T})}}^{d,T} : \exists \mathbf{m}' \in \mathbb{Z}^d \text{ such that } \mathbf{m} = M\mathbf{m}'\}| \leq C_A(d, T) |I_{N 2^{(l+1)(1+\frac{d}{d-T})}}^{d,T}| / M + 1$$

for all refinements $N \in \mathbb{R}$, $N \geq 1$, and levels $l \in \mathbb{N} \cup \{0\}$, where $C_A(d, T) \geq 1$ is a constant which only depends on d and T .

Proof. We denote $A_{N 2^{(l+1)(1+\frac{d}{d-T})}}^{d,T} := \{\mathbf{m} \in I_{N 2^{(l+1)(1+\frac{d}{d-T})}}^{d,T} : \exists \mathbf{m}' \in \mathbb{Z}^d \text{ such that } \mathbf{m} = M\mathbf{m}'\}$ and we group the indices $\mathbf{m} \in A_{N 2^{(l+1)(1+\frac{d}{d-T})}}^{d,T}$, where all components are zero, exactly one component is non-zero, \dots , $d-1$ components are non-zero, and all d components are non-zero. For $t = 0, \dots, d$, we denote

$$A_{N 2^{(l+1)(1+\frac{d}{d-T})}, t}^{d,T} := \left\{ \mathbf{m} \in A_{N 2^{(l+1)(1+\frac{d}{d-T})}}^{d,T} : \text{exactly } t \text{ components of } \mathbf{m} \text{ are non-zero} \right\}.$$

- Case $t = 0$. We have $A_{N 2^{(l+1)(1+\frac{d}{d-T})}, 0}^{d,T} = \{\mathbf{0}\}$.
- Case $1 \leq t \leq d$. If exactly the components m_{i_1}, \dots, m_{i_t} of $\mathbf{m} \in A_{N 2^{(l+1)(1+\frac{d}{d-T})}}^{d,T}$ are

non-zero, $i_1, \dots, i_t \in \{1, \dots, d\}$, $i_j \neq i_{j'}$ for $j \neq j'$, we have

$$\begin{aligned}
& \omega^{-\frac{T}{1-T}, \frac{1}{1-T}}(\mathbf{m}) \\
&= \max(1, M(|m'_{i_1}| + \dots + |m'_{i_t}|))^{-\frac{T}{1-T}} \prod_{\tau=1}^t \max(1, M|m'_{i_\tau}|)^{\frac{1}{1-T}} \\
&= M^{-\frac{T}{1-T}} \max(1, |m'_{i_1}| + \dots + |m'_{i_t}|)^{-\frac{T}{1-T}} M^{\frac{t}{1-T}} \prod_{\tau=1}^t \max(1, M|m'_{i_\tau}|)^{\frac{1}{1-T}} \\
&= M^{\frac{t-T}{1-T}} \omega^{-\frac{T}{1-T}, \frac{1}{1-T}}(\mathbf{m}') \leq N 2^{(l+1)(1+\frac{d}{d-T})} \iff \omega^{-\frac{T}{1-T}, \frac{1}{1-T}}(\mathbf{m}') \leq \frac{N 2^{(l+1)(1+\frac{d}{d-T})}}{M^{\frac{t-T}{1-T}}}.
\end{aligned}$$

Since there are $\binom{d}{t}$ choices for the non-zero components, we have

$$\left| A_{N 2^{(l+1)(1+\frac{d}{d-T})}, t}^{d, T} \right| = \binom{d}{t} \cdot \begin{cases} \mathcal{O}\left(\left(\frac{N 2^{(l+1)}}{M}\right)^t\right) & \text{for } T = -\infty, \\ \mathcal{O}\left(\frac{\left(N 2^{(l+1)(1+\frac{d}{d-T})}\right)^{\frac{T-1}{T/t-1}}}{M^t}\right) & \text{for } -\infty < T < 0, \\ \mathcal{O}\left(\left(\frac{N 2^{(l+1)^2}}{M^t}\right) \log^{t-1}\left(\frac{N 2^{(l+1)^2}}{M^t}\right)\right) & \text{for } T = 0, \end{cases}$$

for fixed $d \in \mathbb{N}$.

This means

- for $T = -\infty$

$$\begin{aligned}
|A_{N 2^{l+1}}^{d, -\infty}| &\leq 1 + \sum_{t=1}^d \binom{d}{t} C_1(d) \left(\frac{N 2^{(l+1)}}{M}\right)^t \leq 1 + \frac{(N 2^{(l+1)})^d}{M} C_1(d) (2^d - 1) \\
&\leq 1 + \frac{|I_{N 2^{l+1}}^{d, -\infty}|}{M} \frac{C_1(d)}{c_1(d)} (2^d - 1)
\end{aligned}$$

due to $|I_{N 2^{l+1}}^{d, -\infty}| \geq c_1(d)(N 2^{(l+1)})^d$ as stated in Lemma 4.1,

- for $-\infty < T < 0$

$$\begin{aligned}
\left| A_{N 2^{(l+1)(1+\frac{d}{d-T})}}^{d, T} \right| &\leq 1 + \sum_{t=1}^d \binom{d}{t} C_2(d) \left(\frac{N 2^{(l+1)(1+\frac{d}{d-T})}}{M^{\frac{t-T}{1-T}}}\right)^{\frac{T-1}{T/t-1}} \\
&= 1 + \sum_{t=1}^d \binom{d}{t} C_2(d) \frac{\left(N 2^{(l+1)(1+\frac{d}{d-T})}\right)^{\frac{t(1-T)}{t-T}}}{M^t} \\
&\leq 1 + C_2(d) \frac{\left(N 2^{(l+1)(1+\frac{d}{d-T})}\right)^{\frac{d(1-T)}{d-T}}}{M} (2^d - 1) \\
&\leq 1 + \frac{C_2(d)}{c_1(d)} \frac{|I_{N 2^{(l+1)(1+\frac{d}{d-T})}}^{d, T}|}{M} (2^d - 1)
\end{aligned}$$

due to $|I_{N 2^{(l+1)(1+\frac{d}{d-T})}}^{d,T}| \geq c_1(d) \left(N 2^{(l+1)(1+\frac{d}{d-T})} \right)^{\frac{T-1}{T/d-1}} = c_1(d) \left(N 2^{(l+1)(1+\frac{d}{d-T})} \right)^{\frac{d(1-T)}{d-T}}$
as stated in Lemma 4.1,

- for $T = 0$

$$\begin{aligned} \left| A_{N 2^{(l+1)2}}^{d,0} \right| &\leq 1 + \sum_{t=1}^d \binom{d}{t} C_3(d) \left(\frac{N 2^{(l+1)2}}{M^t} \right) \log^{t-1} \left(\frac{N 2^{(l+1)2}}{M^t} \right) \\ &\leq 1 + C_3(d) \frac{N 2^{(l+1)2}}{M} \log^{d-1} \left(N 2^{(l+1)2} \right) (2^d - 1) \\ &\leq 1 + \frac{|I_{N 2^{(l+1)2}}^{d,0}|}{M} \frac{C_3(d)}{c_3(d)} (2^d - 1) \end{aligned}$$

due to $|I_{N 2^{(l+1)2}}^{d,0}| \geq c_3(d) N 2^{(l+1)2} \log^{d-1} (N 2^{(l+1)2})$ as stated in Lemma 4.1.

We set

$$C_A(d, T) := (2^d - 1) \cdot \begin{cases} C_1(d)/c_1(d) & \text{for } T = -\infty, \\ C_2(d)/c_1(d) & \text{for } -\infty < T < 0, \\ C_3(d)/c_3(d) & \text{for } T = 0, \end{cases}$$

and this yields the assertion. ■

Lemma 4.6. *Let the dimensionality $d \in \mathbb{N}$, $d \geq 2$ and a function $f \in \mathcal{H}^{\alpha, \beta}(\mathbb{T}^d)$, where $\alpha, \beta \geq 0$ and $\alpha > d(\frac{1}{2} - \beta)$. Then, the function f has an absolutely converging Fourier series,*

$$\sum_{\mathbf{k} \in \mathbb{Z}^d} |\hat{f}_{\mathbf{k}}| < \infty.$$

Proof. Applying the Cauchy-Schwarz inequality yields

$$\begin{aligned} \sum_{\mathbf{k} \in \mathbb{Z}^d} |\hat{f}_{\mathbf{k}}| &= \sum_{\mathbf{k} \in \mathbb{Z}^d} \frac{\omega^{\alpha, \beta}(\mathbf{k})}{\omega^{\alpha, \beta}(\mathbf{k})} |\hat{f}_{\mathbf{k}}| \leq \sqrt{\sum_{\mathbf{k} \in \mathbb{Z}^d} \frac{1}{\omega^{\alpha, \beta}(\mathbf{k})^2}} \sqrt{\sum_{\mathbf{k} \in \mathbb{Z}^d} \omega^{\alpha, \beta}(\mathbf{k})^2 |\hat{f}_{\mathbf{k}}|^2} \\ &= \sqrt{\sum_{\mathbf{k} \in \mathbb{Z}^d} \frac{1}{\max(1, \|\mathbf{k}\|_1)^{2\alpha} \prod_{s=1}^d \max(1, |k_s|)^{2\beta}}} \|f\|_{\mathcal{H}^{\alpha, \beta}(\mathbb{T}^d)}. \end{aligned}$$

Due to $\prod_{s=1}^d \max(1, |k_s|) \leq \max(1, \|\mathbf{k}\|_1)^d$ for $\mathbf{k} \in \mathbb{Z}^d$, we infer

$$\begin{aligned} \sum_{\mathbf{k} \in \mathbb{Z}^d} |\hat{f}_{\mathbf{k}}| &\leq \sqrt{\sum_{\mathbf{k} \in \mathbb{Z}^d} \prod_{s=1}^d \frac{1}{\max(1, |k_s|)^{2(\beta + \frac{\alpha}{d})}}} \|f\|_{\mathcal{H}^{\alpha, \beta}(\mathbb{T}^d)} \\ &= \left(1 + 2\zeta \left(2 \left(\beta + \frac{\alpha}{d} \right) \right) \right)^{\frac{d}{2}} \|f\|_{\mathcal{H}^{\alpha, \beta}(\mathbb{T}^d)}, \end{aligned}$$

where ζ is the Riemann zeta function. Since $\beta \geq 0$ and $\alpha > d(\frac{1}{2} - \beta)$, we obtain $2(\beta + \frac{\alpha}{d}) > 2(\beta + \frac{1}{2} - \beta) = 1$. Due to this and since $f \in \mathcal{H}^{\alpha, \beta}(\mathbb{T}^d)$, we infer $\sum_{\mathbf{k} \in \mathbb{Z}^d} |\hat{f}_{\mathbf{k}}| < \infty$. ■

Theorem 4.7. Let the dimensionality $d \in \mathbb{N}$, $d \geq 2$, a function $f \in \mathcal{H}^{\alpha, \beta}(\mathbb{T}^d)$ and a refinement $N \in \mathbb{R}$, $N \geq 2$ be given, where $\beta \geq 0$, $\alpha \geq 0$,

$$\alpha + \beta > \left(1 + \frac{d}{d-T}\right) \frac{T-1}{T/d-1} \frac{1}{2} \quad (4.3)$$

and the parameter $T := -\alpha/\beta$. Additionally, let a prime number $M \in \mathbb{N}$,

$$M > \frac{d \left| I_{\frac{d-T}{2^{1-T}}}^{d,T} N^{1+\frac{d}{d-T}} \right|}{1-2^{-\kappa}} + 1, \quad (4.4)$$

be given, where we set the parameter $\kappa := \alpha + \beta - \left(1 + \frac{d}{d-T}\right) \frac{T-1}{T/d-1} \frac{1}{2}$. Then, there exists a reconstructing rank-1 lattice $\Lambda(\mathbf{z}, M, I_N^{d,T})$ with generating vector $\mathbf{z} := (1, a, \dots, a^{d-1})^\top \in \mathbb{Z}^d$ of Korobov form and nodes $\mathbf{x}_j := \frac{j}{M} \mathbf{z} \bmod 1$, $j = 0, \dots, M-1$, such that the aliasing error is bounded by

$$\|S_{I_N^{d,T}} f - \tilde{S}_{I_N^{d,T}} f\|_{L^2(\mathbb{T}^d)} \leq C(d, \alpha, \beta) N^{-(\alpha+\beta)} \|f\|_{\mathcal{H}^{\alpha, \beta}(\mathbb{T}^d)},$$

where $C(d, \alpha, \beta) > 0$ is a constant which only depends on d , α , β .

Proof. We are going to apply Theorem 3.4. Therefore, we set $\omega(\mathbf{k}) := \omega^{\alpha, \beta}(\mathbf{k})$, $\nu := \alpha + \beta$ and $I_N := I_N^{d,T}$. Due to $d \geq 2$, $\alpha \geq 0$ and $\beta \geq 0$, we have $\left(1 + \frac{d}{d-T}\right) \frac{T-1}{T/d-1} \frac{1}{2} = \frac{d}{2} \frac{2d\beta + \alpha}{d\beta + \alpha} \frac{\alpha + \beta}{d\beta + \alpha} > 0$ and consequently, $\nu = \alpha + \beta > 0$ follows from condition (4.3). From Lemma 4.3, we obtain $\mathcal{H}^{\alpha, \beta}(\mathbb{T}^d) \subset L^2(\mathbb{T}^d)$. Furthermore, we obtain $\mathcal{D}(I_N^{d,T}) \subset I_{\frac{d-T}{2^{1-T}}}^{d,T} N^{1+\frac{d}{d-T}}$ from Lemma 4.2. Thus, we set $\mathcal{I}_N := I_{\frac{d-T}{2^{1-T}}}^{d,T} N^{1+\frac{d}{d-T}}$ for all $N \in \mathbb{R}$, $N \geq 1$. Applying Lemma 4.5, we infer

$$\begin{aligned} & |\{\mathbf{m} \in \mathcal{I}_{N2^l} : \exists \mathbf{m}' \in \mathbb{Z}^d \text{ such that } \mathbf{m} = M\mathbf{m}'\}| \\ &= |\{\mathbf{m} \in I_{\frac{d-T}{2^{1-T}}}^{d,T} N^{1+\frac{d}{d-T}} 2^{l(1+\frac{d}{d-T})} : \exists \mathbf{m}' \in \mathbb{Z}^d \text{ such that } \mathbf{m} = M\mathbf{m}'\}| \\ &\leq C_A(d, T) \frac{\left| I_{\frac{d-T}{2^{1-T}}}^{d,T} N^{1+\frac{d}{d-T}} 2^{l(1+\frac{d}{d-T})} \right|}{M} + 1 \quad \text{for all } l \in \mathbb{N}. \end{aligned}$$

In order to apply Lemma 4.6, we first show $\alpha > \frac{d}{2} - d\beta$. Due to (4.3), we have $\alpha + \beta > \frac{d}{2} \frac{2d\beta + \alpha}{d\beta + \alpha} \frac{\alpha + \beta}{d\beta + \alpha}$. This is equivalent to the condition $2(d\beta + \alpha)^2 > d(2d\beta + \alpha)$ since $d\beta + \alpha \geq \alpha + \beta > 0$. Due to $2d\beta \geq d\beta$, we obtain $2(d\beta + \alpha)^2 > d(d\beta + \alpha)$. Consequently, we have $\alpha > \frac{d}{2} - d\beta$ such that we can apply Lemma 4.6 and we obtain that f has an absolutely converging Fourier series, whereby f has a point-wise convergent Fourier series.

Next, we apply Theorem 3.4 with $\psi \equiv 1$ and we obtain that there exists a reconstructing rank-1 lattice $\Lambda(\mathbf{z}, M, I_N)$ with generating vector $\mathbf{z} := (1, a, \dots, a^{d-1})^\top \in \mathbb{Z}^d$ of Korobov form, such that the aliasing error is bounded by

$$\begin{aligned} \|S_{I_N} f - \tilde{S}_{I_N} f\|_{L^2(\mathbb{T}^d)} &\leq 2^{\alpha+\beta} N^{-(\alpha+\beta)} \|f\|_{\mathcal{H}^{\alpha, \beta}(\mathbb{T}^d)} \\ &\quad \cdot \sum_{l=0}^{\infty} 2^{(l+1)(\frac{\kappa}{2} - (\alpha+\beta))} \sqrt{\frac{|\mathcal{I}_{N2^{l+1}}|}{|\mathcal{I}_N|}} \sqrt{2(2 + (1-2^{-\kappa}) C_A(d, T))}. \end{aligned}$$

- Case $T = -\infty$, i.e., $\beta = 0$ and $\alpha > \frac{d}{2}$. Due to

$\sqrt{\frac{|\mathcal{I}_{N2^{l+1}}|}{|\mathcal{I}_N|}} = \sqrt{\frac{|I_{2N2^{l+1}}^{d,-\infty}|}{|I_{2N}^{d,-\infty}|}} \leq \sqrt{\frac{C_1(d)}{c_1(d)}} \sqrt{\frac{2^d N^d 2^{(l+1)d}}{2^d N^d}} = \sqrt{\frac{C_1(d)}{c_1(d)}} 2^{(l+1)\frac{d}{2}}$ by Lemma 4.1, where $c_1(d)$ and $C_1(d)$ are constants which only depend on d , we obtain

$$\begin{aligned} \sum_{l=0}^{\infty} 2^{(l+1)(\frac{\kappa}{2}-\alpha)} \sqrt{\frac{|\mathcal{I}_{N2^{l+1}}|}{|\mathcal{I}_N|}} &\leq \sqrt{\frac{C_1(d)}{c_1(d)}} \sum_{l=0}^{\infty} 2^{(l+1)(-\frac{\alpha}{2}+\frac{d}{4})} \\ &= \sqrt{\frac{C_1(d)}{c_1(d)}} \frac{2^{-\frac{\alpha}{2}+\frac{d}{4}}}{1-2^{-\frac{\alpha}{2}+\frac{d}{4}}} =: \tilde{C}(d, \alpha, 0). \end{aligned}$$

- Case $-\infty < T < 0$, i.e., $\beta > 0$, $\alpha > d(\frac{1}{4} + \frac{1}{4}\sqrt{8\beta+1} - \beta)$. Due to

$$\begin{aligned} \sqrt{\frac{|\mathcal{I}_{N2^{l+1}}|}{|\mathcal{I}_N|}} &= \sqrt{\frac{|I_{2N2^{l+1}}^{d,T}|}{|I_{2N}^{d,T}|}} \leq \sqrt{\frac{C_2(d)}{c_1(d)}} \sqrt{\frac{\left(2^{\frac{d-T}{1-T}} N^{1+\frac{d}{d-T}} 2^{(l+1)(1+\frac{d}{d-T})}\right)^{\frac{T-1}{T/d-1}}}{\left(2^{\frac{d-T}{1-T}} N^{1+\frac{d}{d-T}}\right)^{\frac{T-1}{T/d-1}}}} \\ &= \sqrt{\frac{C_2(d)}{c_1(d)}} 2^{(l+1)(1+\frac{d}{d-T})\frac{T-1}{T/d-1} \frac{1}{2}} \end{aligned}$$

by Lemma 4.1, where $c_1(d)$ and $C_2(d)$ are constants which only depend on d , and since we have $(-\frac{\alpha+\beta}{2} + \frac{1}{4}(1 + \frac{d}{d-T})\frac{T-1}{T/d-1}) < 0$ by property (4.3), we obtain

$$\begin{aligned} \sum_{l=0}^{\infty} 2^{(l+1)(\frac{\kappa}{2}-(\alpha+\beta))} \sqrt{\frac{|\mathcal{I}_{N2^{l+1}}|}{|\mathcal{I}_N|}} &\leq \sqrt{\frac{C_2(d)}{c_1(d)}} \sum_{l=0}^{\infty} 2^{(l+1)(-\frac{\alpha+\beta}{2}+\frac{1}{4}(1+\frac{d}{d-T})\frac{T-1}{T/d-1})} \\ &= \sqrt{\frac{C_2(d)}{c_1(d)}} \frac{2^{-\frac{\alpha+\beta}{2}+\frac{1}{4}(1+\frac{d}{d-T})\frac{T-1}{T/d-1}}}{1-2^{-\frac{\alpha+\beta}{2}+\frac{1}{4}(1+\frac{d}{d-T})\frac{T-1}{T/d-1}}} =: \tilde{C}(d, \alpha, \beta). \end{aligned}$$

- Case $T = 0$, i.e., $\beta > 1$ and $\alpha = 0$. Due to

$$\begin{aligned} \sqrt{\frac{|\mathcal{I}_{N2^{l+1}}|}{|\mathcal{I}_N|}} &\leq \sqrt{\frac{C_3(d)}{c_3(d)}} \sqrt{\frac{2^d N^2 2^{2(l+1)} (\log(2^d N^2 2^{2(l+1)}))^{d-1}}{2^d N^2 (\log(2^d N^2))^{d-1}}} \\ &= \sqrt{\frac{C_3(d)}{c_3(d)}} 2^{l+1} \left(\frac{\log(2^d N^2) + \log(2^{2(l+1)})}{\log(2^d N^2)}\right)^{\frac{d-1}{2}} \\ &\leq \sqrt{\frac{C_3(d)}{c_3(d)}} 2^{l+1} \left(2 \log(2^{2(l+1)})\right)^{\frac{d-1}{2}} = \sqrt{\frac{C_3(d)}{c_3(d)}} (2 \log 2)^{\frac{d-1}{2}} 2^{l+1} (2l+2)^{\frac{d-1}{2}} \end{aligned}$$

by Lemma 4.1, where $c_3(d)$ and $C_3(d)$ are constants which only depend on d , we have

$$\sum_{l=0}^{\infty} 2^{(l+1)(\frac{\kappa}{2}-(\alpha+\beta))} \sqrt{\frac{|\mathcal{I}_{N2^{l+1}}|}{|\mathcal{I}_N|}} \leq \sqrt{\frac{C_3(d)}{c_3(d)}} (2 \log 2)^{\frac{d-1}{2}} \sum_{l=0}^{\infty} \frac{(2l+2)^{\frac{d-1}{2}}}{2^{(l+1)\frac{\beta-1}{2}}}.$$

Since $\beta > 1$, the term $\sum_{l=0}^{\infty} \frac{(2l+2)^{\frac{d-1}{2}}}{2^{(l+1)\frac{\beta-1}{2}}} < \infty$ and we are going to estimate this sum. The function $g : [0, \infty) \rightarrow \mathbb{R}$, $g(l) := \frac{(2l+2)^{\frac{d-1}{2}}}{2^{(l+1)\frac{\beta-1}{2}}}$, has its only maximum at

$$l_{\max} := \max\left(0, \frac{d-1}{(\beta-1)\log_e 2} - 1\right)$$

and we estimate

$$\begin{aligned} \sum_{l=0}^{\infty} \frac{(2l+2)^{\frac{d-1}{2}}}{2^{(l+1)\frac{\beta-1}{2}}} &= \sum_{l=0}^{\infty} g(l) \leq \sum_{l=0}^{\lfloor l_{\max} \rfloor} g(l) + \sum_{l=\lceil l_{\max} \rceil}^{\infty} g(l) \\ &\leq g(l_{\max}) + \int_0^{\lfloor l_{\max} \rfloor} g(l) dl + g(l_{\max}) + \int_{\lceil l_{\max} \rceil}^{\infty} g(l) dl \leq 2g(l_{\max}) + \int_0^{\infty} g(l) dl \\ &\leq 2 \max\left(\frac{2^{\frac{d-1}{2}}}{2^{\frac{\beta-1}{2}}}, \left(\frac{2(d-1)}{(\beta-1)e \log_e 2}\right)^{\frac{d-1}{2}}\right) + \frac{(d-1) \left(\frac{4}{(\beta-1)\log_e 2}\right)^{\frac{d-1}{2}} \Gamma\left(\frac{d-1}{2}\right) + 2^{\frac{d+2-\beta}{2}}}{(\beta-1)\log_e 2} \\ &=: \tilde{C}(d, 0, \beta). \end{aligned}$$

These estimates yield

$$\|S_{I_N} f - \tilde{S}_{I_N} f\|_{L^2(\mathbb{T}^d)} \leq \underbrace{\sqrt{2(2 + (1 - 2^{-\kappa}) C_A(d, T)) \tilde{C}(d, \alpha, \beta) 2^{\alpha+\beta}}}_{:=C(d, \alpha, \beta)} N^{-(\alpha+\beta)} \|f\|_{\mathcal{H}^{\alpha, \beta}(\mathbb{T}^d)}.$$

■

4.2.2 Cases $0 < T < 1$

Lemma 4.8. *Let the dimensionality $d \in \mathbb{N}$, $d \geq 2$, a parameter T , $0 < T < 1$, a parameter $\kappa > 0$, and a number $M \in \mathbb{N}$, $M > \frac{d|\mathcal{D}(I_N^{d, T})|}{1-2^{-\kappa}} + 1$ be given. Then, we have*

$$|\mathcal{D}(I_{N2^{l+1}}^{d, T}) \cap M\mathbb{Z}^d| \leq C_A(d, T) \frac{|\mathcal{D}(I_{N2^{l+1}}^{d, T})|}{M} (l+1)^{d-1} + 1$$

for all refinements $N \in \mathbb{R}$, $N \geq 1$, and levels $l \in \mathbb{N} \cup \{0\}$, where $C_A(d, T) \geq 1$ is a constant which only depends on d and T .

Proof. For $0 \leq T < 1$, we denote

$$\mathcal{A}_{N, t}^{d, T} := \{\mathbf{m} \in \mathcal{D}(I_N^{d, T}) \cap M\mathbb{Z}^d : \text{exactly } t \text{ components of } \mathbf{m} \text{ are non-zero}\}, \quad t = 0, \dots, d.$$

Then, we have $\mathcal{D}(I_N^{d, T}) \cap M\mathbb{Z}^d = \bigcup_{t=0}^d \mathcal{A}_{N, t}^{d, T}$ and $|\mathcal{D}(I_N^{d, T}) \cap M\mathbb{Z}^d| = \sum_{t=0}^d |\mathcal{A}_{N, t}^{d, T}|$. Next, we estimate $|\mathcal{A}_{N, t}^{d, T}|$ for $t = 0, \dots, d$.

- Case $t = 0$. Obviously, we have $\mathcal{A}_{N2^{l+1}, 0}^{d, T} = \{\mathbf{0}\}$ and $|\mathcal{A}_{N2^{l+1}, 0}^{d, T}| = 1$.

- Case $t = 1$. $|\mathcal{A}_{N2^{l+1},1}^{d,T}| \leq d \frac{2N2^{l+1}}{M} < 2d \frac{|I_{N2^{l+1}}^{d,T}|}{M} < 2d |\mathcal{D}(I_{N2^{l+1}}^{d,T})|/M$.
- Case $2 \leq t \leq d$. Due to [14, Lemma 2.3] with $\tilde{T} := 0$, we have $I_N^{d,T} \subset I_{d^{1-T}N}^{d,0}$, $N \in \mathbb{R}$, $N \geq 1$, and consequently, we infer

$$\left(\mathcal{D} \left(I_{N2^{l+1}}^{d,T} \right) \cap M\mathbb{Z}^d \right) \subset \left(\mathcal{D} \left(I_{d^{1-T}N2^{l+1}}^{d,0} \right) \cap M\mathbb{Z}^d \right)$$

as well as

$$\mathcal{A}_{N2^{l+1},t}^{d,T} \subset \mathcal{A}_{d^{1-T}N2^{l+1},t}^{d,0} \subset A_{2^d(d^{1-T}N2^{l+1})^2,t}^{d,0} = A_{2^d d^{1-T} N^2 2^{2(l+1)},t}^{d,0},$$

where $A_{N,t}^{d,0} := \left\{ \mathbf{m} \in I_N^{d,0} \cap M\mathbb{Z}^d : \text{exactly } t \text{ components of } \mathbf{m} \text{ are non-zero} \right\}$.

From the proof of Lemma 4.5 and since $|\mathcal{D}(I_N^{d,T})| \geq (2N+1)^2 > N^2$, we obtain

$$\begin{aligned} & \left| \mathcal{A}_{N2^{l+1},t}^{d,T} \right| \leq \left| A_{2^d d^{1-T} N^2 2^{2(l+1)},t}^{d,0} \right| \\ & \leq C_3(d) \binom{d}{t} \frac{2^d d^{\frac{2T}{1-T}} N^2 2^{(l+1)2}}{M^t} \log_2^{t-1} \left(\frac{2^d d^{\frac{2T}{1-T}} N^2 2^{(l+1)2}}{M^t} \right) \\ & \leq C_3(d) \binom{d}{t} 2^d d^{\frac{2T}{1-T}} \frac{|\mathcal{D}(I_{N2^{l+1}}^{d,T})|}{M M^{t-1}} \log_2^{t-1} \left(2^d d^{\frac{2T}{1-T}} \frac{2^{(l+1)2}}{M^{t-1}} \right) \\ & \leq C_3(d) \binom{d}{t} 2^d d^{\frac{2T}{1-T}} \frac{|\mathcal{D}(I_{N2^{l+1}}^{d,T})|}{M M^{t-1}} \left(\log_2 \left(2^d d^{\frac{2T}{1-T}} \right) + \log_2 2^{(l+1)2} \right)^{t-1} \\ & \leq C_3(d) \binom{d}{t} 2^{d+t-1} d^{\frac{2T}{1-T}} \log_2^{t-1} \left(2^d d^{\frac{2T}{1-T}} \right) \frac{|\mathcal{D}(I_{N2^{l+1}}^{d,T})|}{M} \left(\frac{2(l+1)}{M} \right)^{t-1} \\ & \leq C_3(d) 2^{2d-1} d^{\frac{2T}{1-T}} \log_2^{d-1} \left(2^d d^{\frac{2T}{1-T}} \right) \frac{|\mathcal{D}(I_{N2^{l+1}}^{d,T})|}{M} (l+1)^{d-1} \binom{d}{t}. \end{aligned}$$

Consequently, this yields

$$|\mathcal{D}(I_{N2^{l+1}}^{d,T}) \cap M\mathbb{Z}^d| \leq C_3(d) 2^{2d-1} d^{\frac{2T}{1-T}} \log_2^{d-1} \left(2^d d^{\frac{2T}{1-T}} \right) \frac{|\mathcal{D}(I_{N2^{l+1}}^{d,T})|}{M} (l+1)^{d-1} + 1.$$

■

Lemma 4.9. *Let the dimensionality $d \in \mathbb{N}$, $d \geq 2$ and a function $f \in \mathcal{H}^{\alpha,\beta}(\mathbb{T}^d)$, where $0 > \alpha > \frac{1}{2} - \beta$. Then, the function f has an absolutely converging Fourier series,*

$$\sum_{\mathbf{k} \in \mathbb{Z}^d} |\hat{f}_{\mathbf{k}}| < \infty.$$

Proof. As in the proof of Lemma 4.6, we apply the Cauchy-Schwarz inequality and obtain

$$\sum_{\mathbf{k} \in \mathbb{Z}^d} |\hat{f}_{\mathbf{k}}| \leq \sqrt{\sum_{\mathbf{k} \in \mathbb{Z}^d} \frac{1}{\max(1, \|\mathbf{k}\|_1)^{2\alpha} \prod_{s=1}^d \max(1, |k_s|)^{2\beta}}} \|f\|_{\mathcal{H}^{\alpha,\beta}(\mathbb{T}^d)}.$$

Due to $\max(1, \|\mathbf{k}\|_1) \leq 2^d \prod_{s=1}^d \max(1, |k_s|)$ for $\mathbf{k} \in \mathbb{Z}^d$, we infer

$$\begin{aligned} \sum_{\mathbf{k} \in \mathbb{Z}^d} |\hat{f}_{\mathbf{k}}| &\leq \sqrt{\sum_{\mathbf{k} \in \mathbb{Z}^d} 2^{-d\alpha} \prod_{s=1}^d \frac{1}{\max(1, |k_s|)^{2(\beta+\alpha)}}} \|f\|_{\mathcal{H}^{\alpha,\beta}(\mathbb{T}^d)} \\ &= 2^{-\frac{d\alpha}{2}} (1 + 2\zeta(2(\alpha + \beta)))^{\frac{d}{2}} \|f\|_{\mathcal{H}^{\alpha,\beta}(\mathbb{T}^d)}. \end{aligned}$$

Since we have $2(\alpha + \beta) > 1$ and $f \in \mathcal{H}^{\alpha,\beta}(\mathbb{T}^d)$, we obtain $\sum_{\mathbf{k} \in \mathbb{Z}^d} |\hat{f}_{\mathbf{k}}| < \infty$. \blacksquare

Theorem 4.10. *Let the dimensionality $d \in \mathbb{N}$, $d \geq 2$, a function $f \in \mathcal{H}^{\alpha,\beta}(\mathbb{T}^d)$ and a refinement $N \in \mathbb{R}$, $N \geq 2$ be given, where $\alpha < 0$ and $\beta > 1 - \alpha$. Additionally, let a prime number $M \in \mathbb{N}$,*

$$M > \frac{d|\mathcal{D}(I_N^{d,T})|}{1 - 2^{-\kappa}} + 1, \quad (4.5)$$

be given, where the parameter $T := -\alpha/\beta$ and the parameter $\kappa := \alpha + \beta - 1$. Then, there exists a reconstructing rank-1 lattice $\Lambda(\mathbf{z}, M, I_N^{d,T})$ with generating vector $\mathbf{z} := (1, a, \dots, a^{d-1})^\top \in \mathbb{Z}^d$ of Korobov form and nodes $\mathbf{x}_j := \frac{j}{M}\mathbf{z} \bmod 1$, $j = 0, \dots, M - 1$, such that the aliasing error is bounded by

$$\|S_{I_N^{d,T}} f - \tilde{S}_{I_N^{d,T}} f\|_{L^2(\mathbb{T}^d)} \leq C(d, \alpha, \beta) N^{-(\alpha+\beta)} \|f\|_{\mathcal{H}^{\alpha,\beta}(\mathbb{T}^d)},$$

where $C(d, \alpha, \beta) > 0$ is a constant which only depends on d, α, β .

Proof. We are going to apply Theorem 3.4. Therefore, we set $\omega(\mathbf{k}) := \omega^{\alpha,\beta}(\mathbf{k})$, $\nu := \alpha + \beta$, $I_N := I_N^{d,T}$ and $\mathcal{I}_N := \mathcal{D}(I_N^{d,T})$. From Lemma 4.3, we obtain $\mathcal{H}^{\alpha,\beta}(\mathbb{T}^d) \subset L^2(\mathbb{T}^d)$. We apply Lemma 4.8 and this yields

$$\begin{aligned} &|\{\mathbf{m} \in \mathcal{I}_{N2^l} : \exists \mathbf{m}' \in \mathbb{Z}^d \text{ such that } \mathbf{m} = M\mathbf{m}'\}| = |\mathcal{D}(I_{N2^l}^{d,T}) \cap M\mathbb{Z}^d| \\ &\leq C_A(d, T) \frac{|\mathcal{D}(I_{N2^l}^{d,T})|}{M} l^{d-1} + 1 \quad \text{for all } l \in \mathbb{N}. \end{aligned}$$

Furthermore, we need the property that f has a point-wise convergent Fourier series. Since $\alpha > 1 - \beta > \frac{1}{2} - \beta$, we can apply Lemma 4.9 and obtain that f has an absolutely converging Fourier series, whereby f has a point-wise convergent Fourier series.

Next, we apply Theorem 3.4 with $\psi(l) := l^d$ and we obtain

$$\begin{aligned} &\|S_{I_N} f - \tilde{S}_{I_N} f\|_{L^2(\mathbb{T}^d)} \\ &\leq 2^{\alpha+\beta} N^{-(\alpha+\beta)} \|f\|_{\mathcal{H}^{\alpha,\beta}(\mathbb{T}^d)} \\ &\quad \cdot \sum_{l=0}^{\infty} 2^{(l+1)(\frac{\kappa}{2} - (\alpha+\beta))} \sqrt{\frac{|\mathcal{I}_{N2^{l+1}}|}{|\mathcal{I}_N|}} \sqrt{2(2 + (1 - 2^{-\kappa}) C_A(d, T) (l+1)^{d-1})}. \end{aligned}$$

Due to $|\mathcal{I}_{N2^{l+1}}| = |\mathcal{D}(I_{N2^{l+1}}^{d,T})| \leq (C_4(d)N2^{l+1})^2$ and $|\mathcal{I}_N| = |\mathcal{D}(I_N^{d,T})| \geq (2N)^2 > N^2$, where

$C_4(d)$ is a constant which only depends on d , we infer $\sqrt{\frac{|\mathcal{I}_{N2^{l+1}}|}{|\mathcal{I}_N|}} \leq C_4(d) 2^{l+1}$. Then, we obtain

$$\begin{aligned}
& \sum_{l=0}^{\infty} 2^{(l+1)(\frac{\kappa}{2} - (\alpha + \beta))} \sqrt{\frac{|\mathcal{I}_{N2^{l+1}}|}{|\mathcal{I}_N|}} \sqrt{2(2 + (1 - 2^{-\kappa}) C_A(d, T) (l+1)^{d-1})} \\
& < \sum_{l=0}^{\infty} 2^{(l+1)(\frac{\kappa}{2} - (\alpha + \beta))} C_4(d) 2^{l+1} \sqrt{8 C_A(d, T) (l+1)^{d-1}} \\
& = C_4(d) \sqrt{8 C_A(d, T)} \sum_{l=0}^{\infty} 2^{(l+1)(\frac{\alpha + \beta - 1}{2} - (\alpha + \beta) + 1)} (l+1)^{\frac{d-1}{2}} \\
& = C_4(d) \sqrt{8 C_A(d, T)} 2^{-\frac{d-1}{2}} \sum_{l=0}^{\infty} \frac{(2l+2)^{\frac{d-1}{2}}}{2^{(l+1)(\frac{\alpha + \beta - 1}{2})}}
\end{aligned}$$

and the term $\sum_{l=0}^{\infty} \frac{(2l+2)^{\frac{d-1}{2}}}{2^{(l+1)(\frac{\alpha + \beta - 1}{2})}} < \infty$ since $\alpha + \beta > 1$. As in the proof of Theorem 4.7 for the case $T = 0$ replacing β by $\alpha + \beta$, we infer

$$\begin{aligned}
& \frac{C_4(d) \sqrt{8 C_A(d, T)} 2^{-\frac{d-1}{2}}}{\sqrt{2(2 + (1 - 2^{-\kappa}) C_A(d, T))}} \sum_{l=0}^{\infty} \frac{(2l+2)^{\frac{d-1}{2}}}{2^{(l+1)\frac{\alpha + \beta - 1}{2}}} \\
& \leq \frac{C_4(d) \sqrt{8 C_A(d, T)} 2^{-\frac{d-1}{2}}}{\sqrt{2(2 + (1 - 2^{-\kappa}) C_A(d, T))}} 2 \left[\max \left(\frac{2^{\frac{d-1}{2}}}{2^{\frac{\alpha + \beta - 1}{2}}}, \left(\frac{2(d-1)}{(\alpha + \beta - 1) e \log_e 2} \right)^{\frac{d-1}{2}} \right) \right. \\
& \quad \left. + \frac{(d-1) \left(\frac{4}{(\alpha + \beta - 1) \log_e 2} \right)^{\frac{d-1}{2}} \Gamma(\frac{d-1}{2}) + 2^{\frac{d+2-\alpha+\beta}{2}}}{(\alpha + \beta - 1) \log_e 2} \right] \\
& =: \tilde{C}(d, \alpha, \beta).
\end{aligned}$$

These estimates yield

$$\|S_{I_N} f - \tilde{S}_{I_N} f\|_{L^2(\mathbb{T}^d)} \leq \underbrace{\sqrt{2(2 + (1 - 2^{-\kappa}) C_A(d, T))} \tilde{C}(d, \alpha, \beta) 2^{\alpha + \beta}}_{:=C(d, \alpha, \beta)} N^{-(\alpha + \beta)} \|f\|_{\mathcal{H}^{\alpha, \beta}(\mathbb{T}^d)}.$$

■

4.3 Comparison with previous results

In [14], the truncation error $\|f - S_{I_N^{d, T}} f\|_{\mathcal{H}^{r, t}(\mathbb{T}^d)}$ and aliasing error $\|S_{I_N^{d, T}} f - \tilde{S}_{I_N^{d, T}} f\|_{\mathcal{H}^{r, t}(\mathbb{T}^d)}$ were considered for arbitrarily chosen reconstructing rank-1 lattices $\Lambda(\mathbf{z}, M, I_N^{d, T})$ and functions $f \in \mathcal{H}^{\alpha, \beta + \lambda}(\mathbb{T}^d)$, where $r, t \in \mathbb{R}$, $t \geq 0$, $r > -t$, $\beta \geq 0$, $\alpha > -\beta$, $r + t < \alpha + \beta$, $\lambda > 1/2$, and $T := -\frac{\alpha - r}{\beta - t}$. The truncation error was estimated by

$$\|f - S_{I_N^{d, T}} f\|_{\mathcal{H}^{r, t}(\mathbb{T}^d)} \leq N^{-(\alpha - r + \beta - t)} \|f\|_{\mathcal{H}^{\alpha, \beta}(\mathbb{T}^d)}$$

in [14, Lemma 3.5] and for functions f with point-wise convergent Fourier series, the aliasing error was estimated by

$$\|S_{I_N^{d, T}} f - \tilde{S}_{I_N^{d, T}} f\|_{\mathcal{H}^{r, t}(\mathbb{T}^d)} \leq (1 + 2\zeta(2\lambda))^{\frac{d}{2}} N^{-(\alpha - r + \beta - t)} \|f\|_{\mathcal{H}^{\alpha, \beta + \lambda}(\mathbb{T}^d)} \quad (4.6)$$

in [14], which yields

$$\|f - \tilde{S}_{I_N^{d,T}} f|_{\mathcal{H}^{r,t}(\mathbb{T}^d)}\| \leq \left(1 + (1 + 2\zeta(2\lambda))^{\frac{d}{2}}\right) N^{-(\alpha-r+\beta-t)} \|f|_{\mathcal{H}^{\alpha,\beta+\lambda}(\mathbb{T}^d)}\| \quad (4.7)$$

for the approximation error, cf. [14, Corollary 3.8]. We remark that a constructive method for obtaining a reconstructing rank-1 lattice $\Lambda(\mathbf{z}, M, I)$ for a given frequency $I \subset \mathbb{Z}^d$ of finite cardinality is described in [10]. In the present paper, we were able to improve the estimates (4.6) and (4.7). We showed that there exists a reconstructing rank-1 lattice $\Lambda(\mathbf{z}, M, I_N^{d,T})$ with generating vector $\mathbf{z} := (1, a, \dots, a^{d-1})^\top$ of Korobov form such that we do not have the dependence on λ for the special cases $r = t = 0$, $\alpha + \beta > (1 + \frac{d}{d-(T)_-}) \frac{(T)_- - 1}{(T)_- / d - 1} \frac{1}{2}$, where $(T)_- := \min(0, T)$, see Theorem 4.7 and 4.10. However, we do not know a constructive method for obtaining such a reconstructing rank-1 lattice $\Lambda(\mathbf{z}, M, I_N^{d,T})$.

In [7], functions from the spaces of generalized mixed Sobolev smoothness

$$\mathcal{H}_{\text{mix}}^{t,r}(\mathbb{T}^d) := \left\{ f : \sqrt{\sum_{\mathbf{k} \in \mathbb{Z}^d} \prod_{s=1}^d (1 + |k_s|)^{2t} (1 + \|\mathbf{k}\|_\infty)^{2r} |\hat{f}_{\mathbf{k}}|^2} < \infty \right\}.$$

and generalized hyperbolic cross frequency index sets $I = \Gamma_N^T := \{\mathbf{k} \in \mathbb{Z}^d : \prod_{s=1}^d (1 + |k_s|) \cdot (1 + \|\mathbf{k}\|_\infty)^{-T} \leq N^{1-T}\}$ were considered. As sampling nodes \mathbf{x}_j , the nodes of a (generalized) sparse grid with size $M = |\Gamma_N^T|$ were used. We remark that the inclusions $\mathcal{I}_{(N+1)2^{(T-d)/(1-T)}}^{d,T} \subset \Gamma_N^T \subset \mathcal{I}_{(N+1)d^{-T/(1-T)}}^{d,T}$ are valid in the cases $-\infty \leq T \leq 0$ and $\mathcal{I}_{(N+1)d^{-T/(1-T)2^{-d/(1-T)}}}^{d,T} \subset \Gamma_N^T \subset \mathcal{I}_{(N+1)2^{2T/(1-T)}}^{d,T}$ in the cases $0 < T < 1$ for $d \in \mathbb{N}$ and arbitrary refinement $N \in \mathbb{R}$, $N \geq 1$, cf. [14, Lemma 2.5]. Furthermore, we obtain from the proof of [14, Lemma 2.5] that $c(d, r, t) \|f|_{\mathcal{H}^{r,t}(\mathbb{T}^d)}\| \leq \|f|_{\mathcal{H}_{\text{mix}}^{t,r}(\mathbb{T}^d)}\| \leq C(d, r, t) \|f|_{\mathcal{H}^{r,t}(\mathbb{T}^d)}\|$, where

$$c(d, r, t) := \begin{cases} d^{-r} & \text{for } r \geq 0, t \geq 0, \\ 2^r & \text{for } 0 > r > -t, t > 0, \end{cases} \quad C(d, r, t) := \begin{cases} 2^r 2^{dt} & \text{for } r \geq 0, t \geq 0, \\ d^{-r} 2^{dt} & \text{for } 0 > r > -t, t > 0. \end{cases}$$

For the approximation error (and the aliasing error), it was shown, cf. [7, Lemma 8], that

$$\|f - \mathcal{L}_{\Gamma_N^T} f|_{\mathcal{H}_{\text{mix}}^{0,r}(\mathbb{T}^d)}\| \lesssim N^{-(t-r)} (\log N)^{d-1} \|f|_{\mathcal{H}_{\text{mix}}^{t,0}(\mathbb{T}^d)}\|,$$

where $\mathcal{L}_{\Gamma_N^T}$ is the interpolation operator on the (generalized) sparse grid, $0 \leq r < t$, $t > \frac{1}{2}$, $f \in \mathcal{H}_{\text{mix}}^{t,0}(\mathbb{T}^d)$ and $T := \frac{r}{t}$. In particular in the case $r = 0$, the frequency index sets Γ_N^0 are hyperbolic crosses and the above estimate yields

$$\|f - \mathcal{L}_{\Gamma_N^0} f|_{L^2(\mathbb{T}^d)}\| \lesssim N^{-t} (\log N)^{d-1} \|f|_{\mathcal{H}_{\text{mix}}^{t,0}(\mathbb{T}^d)}\|,$$

i.e., there is an additional factor of $(\log N)^{d-1}$ compared to [24, Theorem 2] and (1.4). Similarly in [28, 21], where the case $r = 0$ and sparse grids sampling nodes were considered, it was proven that the approximation error

$$\|f - \mathcal{L}_{\Gamma_N^0} f|_{L^2(\mathbb{T}^d)}\| \leq C(d) N^{-\beta} (\log N)^{\frac{d-1}{2}} \|f|_{\mathcal{H}^{0,\beta}(\mathbb{T}^d)}\|,$$

where $C(d) > 0$ is a constant which only depends on d , see [21, Theorem 1]. This means, there is an additional factor of $(\log N)^{\frac{d-1}{2}}$ compared to [24, Theorem 2] and (1.4). However,

the sampling schemes in [7, 28, 21] only use $M = |I| = \Theta(N \log^{d-1} N)$ many samples, whereas we require $M = \Theta(N^2 \log^{d-1} N)$ many samples, see (1.3). The advantage of our approach is that the computation of the approximated Fourier coefficients $\hat{f}_{\mathbf{k}}$, $\mathbf{k} \in I$, using the sampling method (1.1) is numerically perfectly stable whereas the computation using the sampling schemes from [7, 28, 21] may be numerically unstable, cf. [12].

5 Numerical results

In practice, we do not know a method for verifying if a generating vector $\mathbf{z} := (1, a, \dots, a^{d-1})^\top \in \mathbb{Z}^d$ of Korobov form fulfills property (2.8) in Lemma 2.1 for a given reconstructing rank-1 lattice $\Lambda(\mathbf{z}, M, I)$. Furthermore, we also do not know how to construct a generating vector \mathbf{z} fulfilling property (2.8). However, this special property is crucial for obtaining the estimate (1.4) by Theorem 4.7 and Theorem 4.10. Consequently, we have only the upper bounds from Section 4.3 available. Nevertheless, numerical tests performed in [14, Section 6], which use reconstructing rank-1 lattices $\Lambda(\mathbf{z}, M, I)$ obtained from a constructive method described in [10], showed that the approximation error $\|f - \tilde{S}_{I_N^{d,0}} f\|_{L^2(\mathbb{T}^d)}$ is in $\mathcal{O}(N^{-\beta}) \|f\|_{\mathcal{H}^{0,\beta}(\mathbb{T}^d)}$ for the functions considered there, which is of optimal order. This suggests that the aliasing error can also be

$$\|S_{I_N^{d,0}} f - \tilde{S}_{I_N^{d,0}} f\|_{L^2(\mathbb{T}^d)} \lesssim N^{-\beta} \|f\|_{\mathcal{H}^{0,\beta}(\mathbb{T}^d)}$$

for reconstructing rank-1 lattices $\Lambda(\mathbf{z}, M, I)$ with generating vectors \mathbf{z} which are not necessarily of Korobov form.

Here, we investigate the approximation error more closely and consider the truncation error and the aliasing error. As in [7] and in [14, Example 6.1], we consider the function

$$f(\mathbf{x}) = \prod_{s=1}^d \frac{8\sqrt{6}\sqrt{\pi}}{\sqrt{6369\pi - 4096}} \left[4 + \operatorname{sgn}(x_s - \frac{1}{2}) (\sin(2\pi x_s)^3 + \sin(2\pi x_s)^4) \right], \quad (5.1)$$

where $\|f\|_{L^2(\mathbb{T}^d)} = 1$, $f \in \mathcal{H}^{0, \frac{7}{2} - \epsilon}(\mathbb{T}^d)$, $\epsilon > 0$, $f \notin \mathcal{H}^{0, \frac{7}{2}}(\mathbb{T}^d)$, and the Fourier coefficients

$$\hat{f}_{\mathbf{k}} = \prod_{s=1}^d \frac{8\sqrt{6}\sqrt{\pi}}{\sqrt{6369\pi - 4096}} \begin{cases} \frac{-12}{(k_s-3)(k_s-1)(k_s+1)(k_s+3)\pi} & \text{for } k_s \in 2\mathbb{Z} \setminus \{0\}, \\ \frac{48i}{(k_s-4)(k_s-2)k_s(k_s+2)(k_s+4)\pi} & \text{for } k_s \text{ odd}, \\ 4 - \frac{4}{3\pi} & \text{for } k_s = 0. \end{cases}$$

As frequency index sets I , we use the symmetric hyperbolic cross index sets $I = I_N^{d,0}$ with different refinements N and as sampling nodes \mathbf{x}_j , we use the nodes of the reconstructing rank-1 lattices $\Lambda(\mathbf{z}, M, I_N^{d,0})$ from [14, Table 6.2] for each index set $I_N^{d,0}$. We remark that these reconstructing rank-1 lattices do not fulfill the requirements (2.7) and (2.8) of Lemma (2.1). Nevertheless, we observe for the lattices from [14, Table 6.2] that the truncation errors dominate the aliasing errors, i.e., $\|S_{I_N^{d,0}} f - \tilde{S}_{I_N^{d,0}} f\|_{L^2(\mathbb{T}^d)} \leq \|f - S_{I_N^{d,0}} f\|_{L^2(\mathbb{T}^d)}$. Plots of the $L^2(\mathbb{T}^d)$ approximation error $\|f - \tilde{S}_{I_N^{d,0}} f\|_{L^2(\mathbb{T}^d)}$ are depicted in [14, Figure 6.1a and 6.2a], where it was observed that the approximation error decreases like $\sim N^{-3.45}$ in the one-dimensional case and slightly slower in the multi-dimensional cases. In Figure 5.1 truncation errors $\|f - S_{I_N^{d,0}} f\|_{L^2(\mathbb{T}^d)}$ and aliasing errors $\|S_{I_N^{d,0}} f - \tilde{S}_{I_N^{d,0}} f\|_{L^2(\mathbb{T}^d)}$ are shown for the

cases $d = 2, \dots, 10$. Here, we observe that the aliasing errors $\|S_{I_N^{d,0}} f - \tilde{S}_{I_N^{d,0}} f\|_{L^2(\mathbb{T}^d)}$ are smaller than the truncation error $\|f - S_{I_N^{d,0}} f\|_{L^2(\mathbb{T}^d)}$ and that the aliasing errors $\|S_{I_N^{d,0}} f - \tilde{S}_{I_N^{d,0}} f\|_{L^2(\mathbb{T}^d)}$ also decrease faster. We stress the fact that the truncation errors only depend on the frequency index set $I_N^{d,0}$ and do not depend on the sampling sets. The truncation errors are asymptotically of optimal order. Consequently, the aliasing errors are also of optimal order — assuming that the trends of the plots is maintained.

In Table 5.1, oversampling factors $M/|I_N^{d,0}|$, i.e., ratios of the rank-1 lattice sizes M and the cardinalities of the symmetric hyperbolic cross index sets $I_N^{d,0}$, are shown. These oversampling factors are less than 100 and still moderate compared to the asymptotic statement $\mathcal{O}(N)$ in (1.3) and (1.2), which is $M \sim |I_N^{d,0}|^2 / \log^{d-1} N$. However, we observe that these oversampling factors $M/|I_N^{d,0}|$ grow for increasing refinements N and fixed dimensionality d .

6 Conclusion

In this paper, we generalized the ideas from [24] in order to improve the estimates for the aliasing error $\|S_{I_N^{d,T}} f - \tilde{S}_{I_N^{d,T}} f\|_{L^2(\mathbb{T}^d)}$ from [14] for functions f from the Hilbert spaces $\mathcal{H}^{\alpha,\beta}(\mathbb{T}^d)$ of isotropic and dominating mixed smoothness when using the lattice rule (1.1). We proved the existence of special reconstructing rank-1 lattices $\Lambda(\mathbf{z}, M, I_N^{d,T})$ with generating vectors $\mathbf{z} := (1, a, \dots, a^{d-1})^\top \in \mathbb{Z}^d$ of Korobov form which yield that the order of the aliasing error $\|S_{I_N^{d,T}} f - \tilde{S}_{I_N^{d,T}} f\|_{L^2(\mathbb{T}^d)}$ is bounded by the order of the truncation error $\|f - S_{I_N^{d,T}} f\|_{L^2(\mathbb{T}^d)}$.

The central statement of this paper is Theorem 3.4, which is a generalisation of the ideas of V. N. Temlyakov, see [24]. We stress the fact that our theorem is quite general and applicable to a wide range of frequency index sets I_N . In order to apply Theorem 3.4 to a given sequence of frequency index sets I_N , $N \in \mathbb{R}$, $N \geq 1$, we need to choose a nested sequence of index sets \mathcal{I}_N , see (3.2), such that the inclusion $\mathcal{I}_N \supset \mathcal{D}(I_N)$ is valid, where $\mathcal{D}(I_N)$ is the difference set of I_N , cf. Section 2.1. Thereby, \mathcal{I}_N has to fulfill the following properties:

- The cardinalities $|\mathcal{I}_N|$ should be close to the cardinalities $|\mathcal{D}(I_N)|$. This is crucial for a small size M of the reconstructing rank-1 lattice $\Lambda(\mathbf{z}, M, I_N)$ used as sampling set, see (3.3).
- The upper and lower bound of the cardinalities $|\mathcal{I}_N|$ need to be known and should be almost of the same order, e.g., gaps of logarithmic order between the upper and lower bound are manageable as demonstrated in Section 4.2.2.

Then, the strategy to bound the aliasing error is analog to the approach in Section 4.2. We remark that we dealt with the difference sets themselves in Section 4.2.2 and set $\mathcal{I}_N := \mathcal{D}(I_N^{d,T})$, whereas we covered the difference sets $\mathcal{D}(I_N^{d,T})$ with larger index sets $\mathcal{I}_N := I_{2^{\frac{d-T}{1-T}} N^{1+\frac{d}{d-T}}}$ in Section 4.2.1.

Acknowledgements

DP acknowledges the fruitful discussions and contributions during the Oberwolfach Workshop “Uniform Distribution Theory and Applications”. Especially we thank V. N. Temlyakov for pointing out the valuable paper [24]. We gratefully acknowledge support by the German Research Foundation (DFG) within the Priority Program 1324, project PO 711/10-2.




Figure 5.1: Truncation errors $\|f - S_{I_N^{d,0}} f\|_{L^2(\mathbb{T}^d)}$ and aliasing errors $\|S_{I_N^{d,0}} f - \tilde{S}_{I_N^{d,0}} f\|_{L^2(\mathbb{T}^d)}$ of the function f from (5.1) as a function of the refinement N .

References

- [1] G. Baszenski and F.-J. Delvos. A discrete Fourier transform scheme for Boolean sums of trigonometric operators. In C. K. Chui, W. Schempp, and K. Zeller, editors, *Multivariate Approximation Theory IV*, ISNM 90, pages 15 – 24. Birkhäuser, Basel, 1989.

d	N	$ I_N^{d,0} $	M	$\frac{M}{ I_N^{d,0} }$	$\ f - \tilde{S}_{I_N^{d,0}} f\ _{L^2(\mathbb{T}^d)}$
2	64	1 377	8 451	6.1	1.1e-06
2	128	3 093	33 283	10.8	1.0e-07
2	256	6 889	132 099	19.2	9.0e-09
3	64	10 113	47 463	4.7	7.2e-06
3	128	24 869	176 603	7.1	5.0e-07
3	256	60 217	753 249	12.5	4.8e-08
4	64	61 889	475 829	7.7	2.6e-05
4	128	164 137	2 244 100	13.7	3.5e-06
4	256	426 193	10 561 497	24.8	2.5e-07
5	64	338 305	3 752 318	11.1	5.5e-05
5	128	958 345	20 645 268	21.5	8.6e-06
5	256	2 644 977	136 178 715	51.5	9.8e-07
6	64	1 709 857	31 829 977	18.6	9.5e-05
6	128	5 137 789	192 757 285	37.5	1.6e-05
6	256	14 977 209	1 400 567 254	93.5	2.3e-06
7	8	198 369	1 059 754	5.3	1.7e-02
7	16	716 985	7 798 320	10.9	3.9e-03
7	32	2 465 613	57 114 640	23.2	7.8e-04
8	8	768 609	6 027 975	7.8	2.0e-02
8	16	2 935 521	49 768 670	17.0	4.9e-03
8	32	10 665 297	359 896 131	33.7	1.0e-03
9	4	688 905	6 898 038	10.0	5.8e-02
9	8	2 910 897	34 112 281	11.7	2.3e-02
9	16	11 693 889	320 144 128	27.4	6.0e-03
10	2	452 709	4 315 343	9.5	2.1e-01
10	4	2 421 009	30 780 958	12.7	6.4e-02
10	8	10 819 089	194 144 634	17.9	2.6e-02

Table 5.1: Cardinalities $|I_N^{d,0}|$, rank-1 lattice sizes M , oversampling factors $\frac{M}{|I_N^{d,0}|}$ and approximation errors $\|f - \tilde{S}_{I_N^{d,0}} f\|_{L^2(\mathbb{T}^d)}$ of the function f from (5.1) for various values of d and N .

- [2] H.-J. Bungartz and M. Griebel. A note on the complexity of solving Poisson's equation for spaces of bounded mixed derivatives. *J. Complexity*, 15:167 – 199, 1999.
- [3] H.-J. Bungartz and M. Griebel. Sparse grids. *Acta Numer.*, 13:147 – 269, 2004.
- [4] R. Cools, F. Y. Kuo, and D. Nuyens. Constructing lattice rules based on weighted degree of exactness and worst case error. *Computing*, 87:63 – 89, 2010.
- [5] J. Dick, F. Y. Kuo, and I. H. Sloan. High-dimensional integration: The quasi-monte carlo way. *Acta Numer.*, 22:133 – 288, 2013.
- [6] V. Gradinaru. Fourier transform on sparse grids: Code design and the time dependent Schrödinger equation. *Computing*, 80:1 – 22, 2007.

- [7] M. Griebel and J. Hamaekers. Fast discrete Fourier transform on generalized sparse grids. Univ. Bonn, INS Preprint No. 1305, 2013.
- [8] K. Hallatschek. Fouriertransformation auf dünnen Gittern mit hierarchischen Basen. *Numer. Math.*, 63:83 – 97, 1992.
- [9] Y. Jiang and Y. Xu. Fast discrete algorithms for sparse Fourier expansions of high dimensional functions. *J. Complexity*, 26:51 – 81, 2010.
- [10] L. Kämmerer. Reconstructing hyperbolic cross trigonometric polynomials by sampling along rank-1 lattices. *SIAM J. Numer. Anal.*, 51:2773 – 2796, 2013.
- [11] L. Kämmerer. Reconstructing multivariate trigonometric polynomials from samples along rank-1 lattices. *submitted*, 2013.
- [12] L. Kämmerer and S. Kunis. On the stability of the hyperbolic cross discrete Fourier transform. *Numer. Math.*, 117:581 – 600, 2011.
- [13] L. Kämmerer, S. Kunis, and D. Potts. Interpolation lattices for hyperbolic cross trigonometric polynomials. *J. Complexity*, 28:76 – 92, 2012.
- [14] L. Kämmerer, D. Potts, and T. Volkmer. Approximation of multivariate functions by trigonometric polynomials based on rank-1 lattice sampling. *Preprint 145, DFG Priority Program 1324*, 2013.
- [15] S. Knapek. Approximation und Kompression mit Tensorprodukt-Multiskalenräumen. Dissertation, Universität Bonn, 2000.
- [16] T. Kühn, W. Sickel, and T. Ullrich. Approximation numbers of Sobolev embeddings - sharp constants and tractability. *J. Complexity*, 2013. in print.
- [17] F. Y. Kuo, I. H. Sloan, and H. Woźniakowski. Lattice rules for multivariate approximation in the worst case setting. In *Monte Carlo and quasi-Monte Carlo methods 2004*, pages 289 – 330. Springer, Berlin, 2006.
- [18] D. Li and F. J. Hickernell. Trigonometric spectral collocation methods on lattices. In *Recent Advances in Scientific Computing and Partial Differential Equations, AMS Series in Contemporary Mathematics*, volume 330, pages 121 – 132, 2003.
- [19] H. Munthe-Kaas and T. Sørøvik. Multidimensional pseudo-spectral methods on lattice grids. *Appl. Numer. Math.*, 62:155 – 165, 2012.
- [20] E. Novak and H. Woźniakowski. *Tractability of Multivariate Problems Volume II: Standard Information for Functionals*. Eur. Math. Society, EMS Tracts in Mathematics Vol 12, 2010.
- [21] W. Sickel and T. Ullrich. The Smolyak algorithm, sampling on sparse grids and function spaces of dominating mixed smoothness. *East J. Approx.*, 13:387 – 425, 2007.
- [22] I. H. Sloan and S. Joe. *Lattice methods for multiple integration*. Oxford Science Publications. The Clarendon Press Oxford University Press, New York, 1994.

- [23] F. Sprengel. A class of function spaces and interpolation on sparse grids. *Numer. Funct. Anal. Optim.*, 21:273 – 293, 2000.
- [24] V. N. Temlyakov. Reconstruction of periodic functions of several variables from the values at the nodes of number-theoretic nets. *Anal. Math.*, 12:287 – 305, 1986. In Russian.
- [25] V. N. Temlyakov. Approximation of functions with bounded mixed derivative. *Proc. Steklov Inst. Math.*, 1989. A translation of Trudy Mat. Inst. Steklov 178 (1986).
- [26] V. N. Temlyakov. *Approximation of periodic functions*. Computational Mathematics and Analysis Series. Nova Science Publishers Inc., Commack, NY, 1993.
- [27] V. N. Temlyakov. Cubature formulas, discrepancy, and nonlinear approximation. *J. Complexity*, 19(3):352 – 391, 2003.
- [28] T. Ullrich. Smolyak’s algorithm, sampling on sparse grids and Sobolev spaces of dominating mixed smoothness. *East J. Approx.*, 14:1 – 38, 2008.
- [29] X. Wang. Volumes of Generalized Unit Balls. *Mathematics Magazine*, 78(5):390 – 395, 2005.
- [30] C. Zenger. Sparse grids. In *Parallel algorithms for partial differential equations (Kiel, 1990)*, volume 31 of *Notes Numer. Fluid Mech.*, pages 241 – 251. Vieweg, Braunschweig, Germany, 1991.

Preprint Series DFG-SPP 1324

<http://www.dfg-spp1324.de>

Reports

- [1] R. Ramlau, G. Teschke, and M. Zhariy. A Compressive Landweber Iteration for Solving Ill-Posed Inverse Problems. Preprint 1, DFG-SPP 1324, September 2008.
- [2] G. Plonka. The Easy Path Wavelet Transform: A New Adaptive Wavelet Transform for Sparse Representation of Two-dimensional Data. Preprint 2, DFG-SPP 1324, September 2008.
- [3] E. Novak and H. Woźniakowski. Optimal Order of Convergence and (In-) Tractability of Multivariate Approximation of Smooth Functions. Preprint 3, DFG-SPP 1324, October 2008.
- [4] M. Espig, L. Grasedyck, and W. Hackbusch. Black Box Low Tensor Rank Approximation Using Fibre-Crosses. Preprint 4, DFG-SPP 1324, October 2008.
- [5] T. Bonesky, S. Dahlke, P. Maass, and T. Raasch. Adaptive Wavelet Methods and Sparsity Reconstruction for Inverse Heat Conduction Problems. Preprint 5, DFG-SPP 1324, January 2009.
- [6] E. Novak and H. Woźniakowski. Approximation of Infinitely Differentiable Multivariate Functions Is Intractable. Preprint 6, DFG-SPP 1324, January 2009.
- [7] J. Ma and G. Plonka. A Review of Curvelets and Recent Applications. Preprint 7, DFG-SPP 1324, February 2009.
- [8] L. Denis, D. A. Lorenz, and D. Trede. Greedy Solution of Ill-Posed Problems: Error Bounds and Exact Inversion. Preprint 8, DFG-SPP 1324, April 2009.
- [9] U. Friedrich. A Two Parameter Generalization of Lions' Nonoverlapping Domain Decomposition Method for Linear Elliptic PDEs. Preprint 9, DFG-SPP 1324, April 2009.
- [10] K. Bredies and D. A. Lorenz. Minimization of Non-smooth, Non-convex Functionals by Iterative Thresholding. Preprint 10, DFG-SPP 1324, April 2009.
- [11] K. Bredies and D. A. Lorenz. Regularization with Non-convex Separable Constraints. Preprint 11, DFG-SPP 1324, April 2009.

- [12] M. Döhler, S. Kunis, and D. Potts. Nonequispaced Hyperbolic Cross Fast Fourier Transform. Preprint 12, DFG-SPP 1324, April 2009.
- [13] C. Bender. Dual Pricing of Multi-Exercise Options under Volume Constraints. Preprint 13, DFG-SPP 1324, April 2009.
- [14] T. Müller-Gronbach and K. Ritter. Variable Subspace Sampling and Multi-level Algorithms. Preprint 14, DFG-SPP 1324, May 2009.
- [15] G. Plonka, S. Tenorth, and A. Iske. Optimally Sparse Image Representation by the Easy Path Wavelet Transform. Preprint 15, DFG-SPP 1324, May 2009.
- [16] S. Dahlke, E. Novak, and W. Sickel. Optimal Approximation of Elliptic Problems by Linear and Nonlinear Mappings IV: Errors in L_2 and Other Norms. Preprint 16, DFG-SPP 1324, June 2009.
- [17] B. Jin, T. Khan, P. Maass, and M. Pidcock. Function Spaces and Optimal Currents in Impedance Tomography. Preprint 17, DFG-SPP 1324, June 2009.
- [18] G. Plonka and J. Ma. Curvelet-Wavelet Regularized Split Bregman Iteration for Compressed Sensing. Preprint 18, DFG-SPP 1324, June 2009.
- [19] G. Teschke and C. Borries. Accelerated Projected Steepest Descent Method for Nonlinear Inverse Problems with Sparsity Constraints. Preprint 19, DFG-SPP 1324, July 2009.
- [20] L. Grasedyck. Hierarchical Singular Value Decomposition of Tensors. Preprint 20, DFG-SPP 1324, July 2009.
- [21] D. Rudolf. Error Bounds for Computing the Expectation by Markov Chain Monte Carlo. Preprint 21, DFG-SPP 1324, July 2009.
- [22] M. Hansen and W. Sickel. Best m-term Approximation and Lizorkin-Triebel Spaces. Preprint 22, DFG-SPP 1324, August 2009.
- [23] F.J. Hickernell, T. Müller-Gronbach, B. Niu, and K. Ritter. Multi-level Monte Carlo Algorithms for Infinite-dimensional Integration on \mathbb{R}^N . Preprint 23, DFG-SPP 1324, August 2009.
- [24] S. Dereich and F. Heidenreich. A Multilevel Monte Carlo Algorithm for Lévy Driven Stochastic Differential Equations. Preprint 24, DFG-SPP 1324, August 2009.
- [25] S. Dahlke, M. Fornasier, and T. Raasch. Multilevel Preconditioning for Adaptive Sparse Optimization. Preprint 25, DFG-SPP 1324, August 2009.

- [26] S. Dereich. Multilevel Monte Carlo Algorithms for Lévy-driven SDEs with Gaussian Correction. Preprint 26, DFG-SPP 1324, August 2009.
- [27] G. Plonka, S. Tenorth, and D. Roşca. A New Hybrid Method for Image Approximation using the Easy Path Wavelet Transform. Preprint 27, DFG-SPP 1324, October 2009.
- [28] O. Koch and C. Lubich. Dynamical Low-rank Approximation of Tensors. Preprint 28, DFG-SPP 1324, November 2009.
- [29] E. Faou, V. Gradinaru, and C. Lubich. Computing Semi-classical Quantum Dynamics with Hagedorn Wavepackets. Preprint 29, DFG-SPP 1324, November 2009.
- [30] D. Conte and C. Lubich. An Error Analysis of the Multi-configuration Time-dependent Hartree Method of Quantum Dynamics. Preprint 30, DFG-SPP 1324, November 2009.
- [31] C. E. Powell and E. Ullmann. Preconditioning Stochastic Galerkin Saddle Point Problems. Preprint 31, DFG-SPP 1324, November 2009.
- [32] O. G. Ernst and E. Ullmann. Stochastic Galerkin Matrices. Preprint 32, DFG-SPP 1324, November 2009.
- [33] F. Lindner and R. L. Schilling. Weak Order for the Discretization of the Stochastic Heat Equation Driven by Impulsive Noise. Preprint 33, DFG-SPP 1324, November 2009.
- [34] L. Kämmerer and S. Kunis. On the Stability of the Hyperbolic Cross Discrete Fourier Transform. Preprint 34, DFG-SPP 1324, December 2009.
- [35] P. Cerejeiras, M. Ferreira, U. Kähler, and G. Teschke. Inversion of the noisy Radon transform on $SO(3)$ by Gabor frames and sparse recovery principles. Preprint 35, DFG-SPP 1324, January 2010.
- [36] T. Jahnke and T. Udrescu. Solving Chemical Master Equations by Adaptive Wavelet Compression. Preprint 36, DFG-SPP 1324, January 2010.
- [37] P. Kittipoom, G. Kutyniok, and W.-Q. Lim. Irregular Shearlet Frames: Geometry and Approximation Properties. Preprint 37, DFG-SPP 1324, February 2010.
- [38] G. Kutyniok and W.-Q. Lim. Compactly Supported Shearlets are Optimally Sparse. Preprint 38, DFG-SPP 1324, February 2010.

- [39] M. Hansen and W. Sickel. Best m -Term Approximation and Tensor Products of Sobolev and Besov Spaces – the Case of Non-compact Embeddings. Preprint 39, DFG-SPP 1324, March 2010.
- [40] B. Niu, F.J. Hickernell, T. Müller-Gronbach, and K. Ritter. Deterministic Multi-level Algorithms for Infinite-dimensional Integration on $\mathbb{R}^{\mathbb{N}}$. Preprint 40, DFG-SPP 1324, March 2010.
- [41] P. Kittipoom, G. Kutyniok, and W.-Q Lim. Construction of Compactly Supported Shearlet Frames. Preprint 41, DFG-SPP 1324, March 2010.
- [42] C. Bender and J. Steiner. Error Criteria for Numerical Solutions of Backward SDEs. Preprint 42, DFG-SPP 1324, April 2010.
- [43] L. Grasedyck. Polynomial Approximation in Hierarchical Tucker Format by Vector-Tensorization. Preprint 43, DFG-SPP 1324, April 2010.
- [44] M. Hansen und W. Sickel. Best m -Term Approximation and Sobolev-Besov Spaces of Dominating Mixed Smoothness - the Case of Compact Embeddings. Preprint 44, DFG-SPP 1324, April 2010.
- [45] P. Binev, W. Dahmen, and P. Lamby. Fast High-Dimensional Approximation with Sparse Occupancy Trees. Preprint 45, DFG-SPP 1324, May 2010.
- [46] J. Ballani and L. Grasedyck. A Projection Method to Solve Linear Systems in Tensor Format. Preprint 46, DFG-SPP 1324, May 2010.
- [47] P. Binev, A. Cohen, W. Dahmen, R. DeVore, G. Petrova, and P. Wojtaszczyk. Convergence Rates for Greedy Algorithms in Reduced Basis Methods. Preprint 47, DFG-SPP 1324, May 2010.
- [48] S. Kestler and K. Urban. Adaptive Wavelet Methods on Unbounded Domains. Preprint 48, DFG-SPP 1324, June 2010.
- [49] H. Yserentant. The Mixed Regularity of Electronic Wave Functions Multiplied by Explicit Correlation Factors. Preprint 49, DFG-SPP 1324, June 2010.
- [50] H. Yserentant. On the Complexity of the Electronic Schrödinger Equation. Preprint 50, DFG-SPP 1324, June 2010.
- [51] M. Guillemard and A. Iske. Curvature Analysis of Frequency Modulated Manifolds in Dimensionality Reduction. Preprint 51, DFG-SPP 1324, June 2010.
- [52] E. Herrholz and G. Teschke. Compressive Sensing Principles and Iterative Sparse Recovery for Inverse and Ill-Posed Problems. Preprint 52, DFG-SPP 1324, July 2010.

- [53] L. Kämmerer, S. Kunis, and D. Potts. Interpolation Lattices for Hyperbolic Cross Trigonometric Polynomials. Preprint 53, DFG-SPP 1324, July 2010.
- [54] G. Kutyniok and W.-Q Lim. Shearlets on Bounded Domains. Preprint 54, DFG-SPP 1324, July 2010.
- [55] A. Zeiser. Wavelet Approximation in Weighted Sobolev Spaces of Mixed Order with Applications to the Electronic Schrödinger Equation. Preprint 55, DFG-SPP 1324, July 2010.
- [56] G. Kutyniok, J. Lemvig, and W.-Q Lim. Compactly Supported Shearlets. Preprint 56, DFG-SPP 1324, July 2010.
- [57] A. Zeiser. On the Optimality of the Inexact Inverse Iteration Coupled with Adaptive Finite Element Methods. Preprint 57, DFG-SPP 1324, July 2010.
- [58] S. Jokar. Sparse Recovery and Kronecker Products. Preprint 58, DFG-SPP 1324, August 2010.
- [59] T. Aboiyar, E. H. Georgoulis, and A. Iske. Adaptive ADER Methods Using Kernel-Based Polyharmonic Spline WENO Reconstruction. Preprint 59, DFG-SPP 1324, August 2010.
- [60] O. G. Ernst, A. Mugler, H.-J. Starkloff, and E. Ullmann. On the Convergence of Generalized Polynomial Chaos Expansions. Preprint 60, DFG-SPP 1324, August 2010.
- [61] S. Holtz, T. Rohwedder, and R. Schneider. On Manifolds of Tensors of Fixed TT-Rank. Preprint 61, DFG-SPP 1324, September 2010.
- [62] J. Ballani, L. Grasedyck, and M. Kluge. Black Box Approximation of Tensors in Hierarchical Tucker Format. Preprint 62, DFG-SPP 1324, October 2010.
- [63] M. Hansen. On Tensor Products of Quasi-Banach Spaces. Preprint 63, DFG-SPP 1324, October 2010.
- [64] S. Dahlke, G. Steidl, and G. Teschke. Shearlet Coorbit Spaces: Compactly Supported Analyzing Shearlets, Traces and Embeddings. Preprint 64, DFG-SPP 1324, October 2010.
- [65] W. Hackbusch. Tensorisation of Vectors and their Efficient Convolution. Preprint 65, DFG-SPP 1324, November 2010.
- [66] P. A. Cioica, S. Dahlke, S. Kinzel, F. Lindner, T. Raasch, K. Ritter, and R. L. Schilling. Spatial Besov Regularity for Stochastic Partial Differential Equations on Lipschitz Domains. Preprint 66, DFG-SPP 1324, November 2010.

- [67] E. Novak and H. Woźniakowski. On the Power of Function Values for the Approximation Problem in Various Settings. Preprint 67, DFG-SPP 1324, November 2010.
- [68] A. Hinrichs, E. Novak, and H. Woźniakowski. The Curse of Dimensionality for Monotone and Convex Functions of Many Variables. Preprint 68, DFG-SPP 1324, November 2010.
- [69] G. Kutyniok and W.-Q. Lim. Image Separation Using Shearlets. Preprint 69, DFG-SPP 1324, November 2010.
- [70] B. Jin and P. Maass. An Analysis of Electrical Impedance Tomography with Applications to Tikhonov Regularization. Preprint 70, DFG-SPP 1324, December 2010.
- [71] S. Holtz, T. Rohwedder, and R. Schneider. The Alternating Linear Scheme for Tensor Optimisation in the TT Format. Preprint 71, DFG-SPP 1324, December 2010.
- [72] T. Müller-Gronbach and K. Ritter. A Local Refinement Strategy for Constructive Quantization of Scalar SDEs. Preprint 72, DFG-SPP 1324, December 2010.
- [73] T. Rohwedder and R. Schneider. An Analysis for the DIIS Acceleration Method used in Quantum Chemistry Calculations. Preprint 73, DFG-SPP 1324, December 2010.
- [74] C. Bender and J. Steiner. Least-Squares Monte Carlo for Backward SDEs. Preprint 74, DFG-SPP 1324, December 2010.
- [75] C. Bender. Primal and Dual Pricing of Multiple Exercise Options in Continuous Time. Preprint 75, DFG-SPP 1324, December 2010.
- [76] H. Harbrecht, M. Peters, and R. Schneider. On the Low-rank Approximation by the Pivoted Cholesky Decomposition. Preprint 76, DFG-SPP 1324, December 2010.
- [77] P. A. Cioica, S. Dahlke, N. Döhring, S. Kinzel, F. Lindner, T. Raasch, K. Ritter, and R. L. Schilling. Adaptive Wavelet Methods for Elliptic Stochastic Partial Differential Equations. Preprint 77, DFG-SPP 1324, January 2011.
- [78] G. Plonka, S. Tenorth, and A. Iske. Optimal Representation of Piecewise Hölder Smooth Bivariate Functions by the Easy Path Wavelet Transform. Preprint 78, DFG-SPP 1324, January 2011.

- [79] A. Mugler and H.-J. Starkloff. On Elliptic Partial Differential Equations with Random Coefficients. Preprint 79, DFG-SPP 1324, January 2011.
- [80] T. Müller-Gronbach, K. Ritter, and L. Yaroslavtseva. A Derandomization of the Euler Scheme for Scalar Stochastic Differential Equations. Preprint 80, DFG-SPP 1324, January 2011.
- [81] W. Dahmen, C. Huang, C. Schwab, and G. Welper. Adaptive Petrov-Galerkin methods for first order transport equations. Preprint 81, DFG-SPP 1324, January 2011.
- [82] K. Grella and C. Schwab. Sparse Tensor Spherical Harmonics Approximation in Radiative Transfer. Preprint 82, DFG-SPP 1324, January 2011.
- [83] D.A. Lorenz, S. Schiffler, and D. Trede. Beyond Convergence Rates: Exact Inversion With Tikhonov Regularization With Sparsity Constraints. Preprint 83, DFG-SPP 1324, January 2011.
- [84] S. Dereich, M. Scheutzow, and R. Schottstedt. Constructive quantization: Approximation by empirical measures. Preprint 84, DFG-SPP 1324, January 2011.
- [85] S. Dahlke and W. Sickel. On Besov Regularity of Solutions to Nonlinear Elliptic Partial Differential Equations. Preprint 85, DFG-SPP 1324, January 2011.
- [86] S. Dahlke, U. Friedrich, P. Maass, T. Raasch, and R.A. Ressel. An adaptive wavelet method for parameter identification problems in parabolic partial differential equations. Preprint 86, DFG-SPP 1324, January 2011.
- [87] A. Cohen, W. Dahmen, and G. Welper. Adaptivity and Variational Stabilization for Convection-Diffusion Equations. Preprint 87, DFG-SPP 1324, January 2011.
- [88] T. Jahnke. On Reduced Models for the Chemical Master Equation. Preprint 88, DFG-SPP 1324, January 2011.
- [89] P. Binev, W. Dahmen, R. DeVore, P. Lamby, D. Savu, and R. Sharpley. Compressed Sensing and Electron Microscopy. Preprint 89, DFG-SPP 1324, March 2011.
- [90] P. Binev, F. Blanco-Silva, D. Blom, W. Dahmen, P. Lamby, R. Sharpley, and T. Vogt. High Quality Image Formation by Nonlocal Means Applied to High-Angle Annular Dark Field Scanning Transmission Electron Microscopy (HAADF-STEM). Preprint 90, DFG-SPP 1324, March 2011.
- [91] R. A. Ressel. A Parameter Identification Problem for a Nonlinear Parabolic Differential Equation. Preprint 91, DFG-SPP 1324, May 2011.

- [92] G. Kutyniok. Data Separation by Sparse Representations. Preprint 92, DFG-SPP 1324, May 2011.
- [93] M. A. Davenport, M. F. Duarte, Y. C. Eldar, and G. Kutyniok. Introduction to Compressed Sensing. Preprint 93, DFG-SPP 1324, May 2011.
- [94] H.-C. Kreuzler and H. Yserentant. The Mixed Regularity of Electronic Wave Functions in Fractional Order and Weighted Sobolev Spaces. Preprint 94, DFG-SPP 1324, June 2011.
- [95] E. Ullmann, H. C. Elman, and O. G. Ernst. Efficient Iterative Solvers for Stochastic Galerkin Discretizations of Log-Transformed Random Diffusion Problems. Preprint 95, DFG-SPP 1324, June 2011.
- [96] S. Kunis and I. Melzer. On the Butterfly Sparse Fourier Transform. Preprint 96, DFG-SPP 1324, June 2011.
- [97] T. Rohwedder. The Continuous Coupled Cluster Formulation for the Electronic Schrödinger Equation. Preprint 97, DFG-SPP 1324, June 2011.
- [98] T. Rohwedder and R. Schneider. Error Estimates for the Coupled Cluster Method. Preprint 98, DFG-SPP 1324, June 2011.
- [99] P. A. Cioica and S. Dahlke. Spatial Besov Regularity for Semilinear Stochastic Partial Differential Equations on Bounded Lipschitz Domains. Preprint 99, DFG-SPP 1324, July 2011.
- [100] L. Grasedyck and W. Hackbusch. An Introduction to Hierarchical (H-) Rank and TT-Rank of Tensors with Examples. Preprint 100, DFG-SPP 1324, August 2011.
- [101] N. Chegini, S. Dahlke, U. Friedrich, and R. Stevenson. Piecewise Tensor Product Wavelet Bases by Extensions and Approximation Rates. Preprint 101, DFG-SPP 1324, September 2011.
- [102] S. Dahlke, P. Oswald, and T. Raasch. A Note on Quarkonial Systems and Multi-level Partition of Unity Methods. Preprint 102, DFG-SPP 1324, September 2011.
- [103] A. Uschmajew. Local Convergence of the Alternating Least Squares Algorithm For Canonical Tensor Approximation. Preprint 103, DFG-SPP 1324, September 2011.
- [104] S. Kvaal. Multiconfigurational time-dependent Hartree method for describing particle loss due to absorbing boundary conditions. Preprint 104, DFG-SPP 1324, September 2011.

- [105] M. Guillemard and A. Iske. On Groupoid C^* -Algebras, Persistent Homology and Time-Frequency Analysis. Preprint 105, DFG-SPP 1324, September 2011.
- [106] A. Hinrichs, E. Novak, and H. Woźniakowski. Discontinuous information in the worst case and randomized settings. Preprint 106, DFG-SPP 1324, September 2011.
- [107] M. Espig, W. Hackbusch, A. Litvinenko, H. Matthies, and E. Zander. Efficient Analysis of High Dimensional Data in Tensor Formats. Preprint 107, DFG-SPP 1324, September 2011.
- [108] M. Espig, W. Hackbusch, S. Handschuh, and R. Schneider. Optimization Problems in Contracted Tensor Networks. Preprint 108, DFG-SPP 1324, October 2011.
- [109] S. Dereich, T. Müller-Gronbach, and K. Ritter. On the Complexity of Computing Quadrature Formulas for SDEs. Preprint 109, DFG-SPP 1324, October 2011.
- [110] D. Belomestny. Solving optimal stopping problems by empirical dual optimization and penalization. Preprint 110, DFG-SPP 1324, November 2011.
- [111] D. Belomestny and J. Schoenmakers. Multilevel dual approach for pricing American style derivatives. Preprint 111, DFG-SPP 1324, November 2011.
- [112] T. Rohwedder and A. Uschmajew. Local convergence of alternating schemes for optimization of convex problems in the TT format. Preprint 112, DFG-SPP 1324, December 2011.
- [113] T. Görner, R. Hielscher, and S. Kunis. Efficient and accurate computation of spherical mean values at scattered center points. Preprint 113, DFG-SPP 1324, December 2011.
- [114] Y. Dong, T. Görner, and S. Kunis. An iterative reconstruction scheme for photoacoustic imaging. Preprint 114, DFG-SPP 1324, December 2011.
- [115] L. Kämmerer. Reconstructing hyperbolic cross trigonometric polynomials by sampling along generated sets. Preprint 115, DFG-SPP 1324, February 2012.
- [116] H. Chen and R. Schneider. Numerical analysis of augmented plane waves methods for full-potential electronic structure calculations. Preprint 116, DFG-SPP 1324, February 2012.
- [117] J. Ma, G. Plonka, and M.Y. Hussaini. Compressive Video Sampling with Approximate Message Passing Decoding. Preprint 117, DFG-SPP 1324, February 2012.

- [118] D. Heinen and G. Plonka. Wavelet shrinkage on paths for scattered data denoising. Preprint 118, DFG-SPP 1324, February 2012.
- [119] T. Jahnke and M. Kreim. Error bound for piecewise deterministic processes modeling stochastic reaction systems. Preprint 119, DFG-SPP 1324, March 2012.
- [120] C. Bender and J. Steiner. A-posteriori estimates for backward SDEs. Preprint 120, DFG-SPP 1324, April 2012.
- [121] M. Espig, W. Hackbusch, A. Litvinenkoy, H.G. Matthiesy, and P. Wähnert. Efficient low-rank approximation of the stochastic Galerkin matrix in tensor formats. Preprint 121, DFG-SPP 1324, May 2012.
- [122] O. Bokanowski, J. Garcke, M. Griebel, and I. Klompmaker. An adaptive sparse grid semi-Lagrangian scheme for first order Hamilton-Jacobi Bellman equations. Preprint 122, DFG-SPP 1324, June 2012.
- [123] A. Mugler and H.-J. Starkloff. On the convergence of the stochastic Galerkin method for random elliptic partial differential equations. Preprint 123, DFG-SPP 1324, June 2012.
- [124] P.A. Cioica, S. Dahlke, N. Döhring, U. Friedrich, S. Kinzel, F. Lindner, T. Raasch, K. Ritter, and R.L. Schilling. On the convergence analysis of Rothe’s method. Preprint 124, DFG-SPP 1324, July 2012.
- [125] P. Binev, A. Cohen, W. Dahmen, and R. DeVore. Classification Algorithms using Adaptive Partitioning. Preprint 125, DFG-SPP 1324, July 2012.
- [126] C. Lubich, T. Rohwedder, R. Schneider, and B. Vandereycken. Dynamical approximation of hierarchical Tucker and Tensor-Train tensors. Preprint 126, DFG-SPP 1324, July 2012.
- [127] M. Kovács, S. Larsson, and K. Urban. On Wavelet-Galerkin methods for semilinear parabolic equations with additive noise. Preprint 127, DFG-SPP 1324, August 2012.
- [128] M. Bachmayr, H. Chen, and R. Schneider. Numerical analysis of Gaussian approximations in quantum chemistry. Preprint 128, DFG-SPP 1324, August 2012.
- [129] D. Rudolf. Explicit error bounds for Markov chain Monte Carlo. Preprint 129, DFG-SPP 1324, August 2012.
- [130] P.A. Cioica, K.-H. Kim, K. Lee, and F. Lindner. On the $L_q(L_p)$ -regularity and Besov smoothness of stochastic parabolic equations on bounded Lipschitz domains. Preprint 130, DFG-SPP 1324, December 2012.

- [131] M. Hansen. n -term Approximation Rates and Besov Regularity for Elliptic PDEs on Polyhedral Domains. Preprint 131, DFG-SPP 1324, December 2012.
- [132] R. E. Bank and H. Yserentant. On the H^1 -stability of the L_2 -projection onto finite element spaces. Preprint 132, DFG-SPP 1324, December 2012.
- [133] M. Gnewuch, S. Mayer, and K. Ritter. On Weighted Hilbert Spaces and Integration of Functions of Infinitely Many Variables. Preprint 133, DFG-SPP 1324, December 2012.
- [134] D. Crisan, J. Diehl, P.K. Friz, and H. Oberhauser. Robust Filtering: Correlated Noise and Multidimensional Observation. Preprint 134, DFG-SPP 1324, January 2013.
- [135] Wolfgang Dahmen, Christian Plesken, and Gerrit Welper. Double Greedy Algorithms: Reduced Basis Methods for Transport Dominated Problems. Preprint 135, DFG-SPP 1324, February 2013.
- [136] Aicke Hinrichs, Erich Novak, Mario Ullrich, and Henryk Wozniakowski. The Curse of Dimensionality for Numerical Integration of Smooth Functions. Preprint 136, DFG-SPP 1324, February 2013.
- [137] Markus Bachmayr, Wolfgang Dahmen, Ronald DeVore, and Lars Grasedyck. Approximation of High-Dimensional Rank One Tensors. Preprint 137, DFG-SPP 1324, March 2013.
- [138] Markus Bachmayr and Wolfgang Dahmen. Adaptive Near-Optimal Rank Tensor Approximation for High-Dimensional Operator Equations. Preprint 138, DFG-SPP 1324, April 2013.
- [139] Felix Lindner. Singular Behavior of the Solution to the Stochastic Heat Equation on a Polygonal Domain. Preprint 139, DFG-SPP 1324, May 2013.
- [140] Stephan Dahlke, Dominik Lellek, Shiu Hong Lui, and Rob Stevenson. Adaptive Wavelet Schwarz Methods for the Navier-Stokes Equation. Preprint 140, DFG-SPP 1324, May 2013.
- [141] Jonas Ballani and Lars Grasedyck. Tree Adaptive Approximation in the Hierarchical Tensor Format. Preprint 141, DFG-SPP 1324, June 2013.
- [142] Harry Yserentant. A short theory of the Rayleigh-Ritz method. Preprint 142, DFG-SPP 1324, July 2013.
- [143] M. Hefter and K. Ritter. On Embeddings of Weighted Tensor Product Hilbert Spaces. Preprint 143, DFG-SPP 1324, August 2013.

- [144] M. Altmayer and A. Neuenkirch. Multilevel Monte Carlo Quadrature of Discontinuous Payoffs in the Generalized Heston Model using Malliavin Integration by Parts. Preprint 144, DFG-SPP 1324, August 2013.
- [145] L. Kämmerer, D. Potts, and T. Volkmer. Approximation of multivariate functions by trigonometric polynomials based on rank-1 lattice sampling. Preprint 145, DFG-SPP 1324, September 2013.
- [146] C. Bender, N. Schweizer, and J. Zhuo. A primal-dual algorithm for BSDEs. Preprint 146, DFG-SPP 1324, October 2013.
- [147] D. Rudolf. Hit-and-run for numerical integration. Preprint 147, DFG-SPP 1324, October 2013.
- [148] D. Rudolf and M. Ullrich. Positivity of hit-and-run and related algorithms. Preprint 148, DFG-SPP 1324, October 2013.
- [149] L. Grasedyck, M. Kluge, and S. Krämer. Alternating Directions Fitting (ADF) of Hierarchical Low Rank Tensors. Preprint 149, DFG-SPP 1324, October 2013.
- [150] F. Filbir, S. Kunis, and R. Seyfried. Effective discretization of direct reconstruction schemes for photoacoustic imaging in spherical geometries. Preprint 150, DFG-SPP 1324, November 2013.
- [151] E. Novak, M. Ullrich, and H. Woźniakowski. Complexity of Oscillatory Integration for Univariate Sobolev Spaces. Preprint 151, DFG-SPP 1324, November 2013.
- [152] A. Hinrichs, E. Novak, and M. Ullrich. A Tractability Result for the Clenshaw Curtis Smolyak Algorithm. Preprint 152, DFG-SPP 1324, November 2013.
- [153] M. Hein, S. Setzer, L. Jost, and S. Rangapuram. The Total Variation on Hypergraphs - Learning on Hypergraphs Revisited. Preprint 153, DFG-SPP 1324, November 2013.
- [154] M. Kovács, S. Larsson, and F. Lindgren. On the Backward Euler Approximation of the Stochastic Allen-Chan Equation. Preprint 154, DFG-SPP 1324, November 2013.
- [155] S. Dahlke, M. Fornasier, U. Friedrich, and T. Raasch. Multilevel preconditioning for sparse optimization of functionals with nonconvex fidelity terms. Preprint 155, DFG-SPP 1324, December 2013.
- [156] T. Müller-Gronbach, K. Ritter, and L. Yaroslavtseva. On the complexity of computing quadrature formulas for marginal distributions of SDEs. Preprint 156, DFG-SPP 1324, January 2014.

- [157] M. Giles, T. Nagapetyan, and K. Ritter. Multi-Level Monte Carlo Approximation of Distribution Functions and Densities. Preprint 157, DFG-SPP 1324, February 2014.
- [158] F. Dickmann and N. Schweizer. Faster Comparison of Stopping Times by Nested Conditional Monte Carlo . Preprint 158, DFG-SPP 1324, February 2014.
- [159] L. Kämmerer, D. Potts, and T. Volkmer. Approximation of multivariate periodic functions by trigonometric polynomials based on sampling along rank-1 lattice with generating vector of Korobov form . Preprint 159, DFG-SPP 1324, February 2014.